

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number  
**WO 02/22605 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 403/12**,  
401/14, 409/14, A61K 31/497, 31/53, A61P 35/00

(21) International Application Number: PCT/US01/28793

(22) International Filing Date:  
14 September 2001 (14.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
(4)/232,795 15 September 2000 (15.09.2000) US  
(4)/257,887 21 December 2000 (21.12.2000) US  
(4)/286,949 27 April 2001 (27.04.2001) US

(71) Applicant (for all designated States except US): **VERTEX PHARMACEUTICALS INCORPORATED** [US/US]; Patent Department, 130 Waverly Street, Cambridge, MA 02139-4242 (US).

(72) Inventors: and

(75) Inventors/Applicants (for US only): **GOLEC, Julian**, M.C. [GB/GB]; 8 Manor Farm, Chapel Road, Ashbury, Wiltshire SN6 8LS (GB). **CHARRIER, Jean-Damien** [FR/GB], Vertex Pharmaceuticals Inc., Cottage Wing, Station Road, Southam, Bishops Itchington, Oxfordshire CV47 2QB (GB). **KNEGTEL, Ronald** [GB/GB]; 3 Bath Court, Bath Street, Abingdon, Oxfordshire OX1X1EE (GB). **BEBBINGTON, David** [GB/GB]; 6 Linden Close,

Newbury, Berkshire R6141QA (GB). **DAVIES, Robert** [GB/US]; 65 Orient Avenue, Arlington, MA 02474 (US). **LI, Pan** [CN/US]; 15 Mystic View Terrace, Arlington, MA 02474 (US).

(74) Agents: **GOVINDASWAMY, Nandakumar**; Vertex Pharmaceuticals Inc., 130 Waverly Street, Cambridge, MA 02139-4242 et al. (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRAZOLE COMPOUNDS USEFUL AS PROTEIN KINASE INHIBITORS



(VIII)

(57) Abstract: This invention describes novel pyrazole compositions comprising a pharmaceutically acceptable carrier and a compound of formula VIII, wherein Z<sup>1</sup> is N or C-R<sup>9</sup>, Z<sup>2</sup> is N or CH, and Z<sup>3</sup> is N or C-R<sup>5</sup>, provided that one of Z<sup>1</sup> and Z<sup>3</sup> is nitrogen; G is Ring C or Ring D; Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from -R<sup>1</sup>; Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl; and R<sup>5</sup>, R<sup>1</sup>, R<sup>2</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>9</sup> are as described in the specification. The compounds are useful as protein kinase inhibitors, especially as inhibitors of aurora-2 and GSK-3, for treating diseases such as cancer, diabetes and Alzheimer's disease.

WO 02/22605 A1

PYRAZOLE COMPOUNDS USEFUL AS PROTEIN KINASE INHIBITORSCROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to US Provisional Patent Application 60/232,795 filed September 15, 2000, US Provisional Patent Application 60/257,887 filed December 21, 2000 and US Provisional Patent Application 60/286,949 filed April 27, 2001, the contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

The present invention is in the field of medicinal chemistry and relates to compounds that are protein kinase inhibitors, compositions containing such compounds and methods of use. More particularly, this invention relates to compounds that are inhibitors of GSK-3 and Aurora-2 protein kinases. The invention also relates to methods of treating diseases associated with these protein kinases, such as diabetes, cancer and Alzheimer's disease.

BACKGROUND OF THE INVENTION

The search for new therapeutic agents has been greatly aided in recent years by better understanding of the structure of enzymes and other biomolecules associated with target diseases. One important class of enzymes that has been the subject of extensive study is the protein kinases.

Protein kinases mediate intracellular signal transduction. They do this by effecting a phosphoryl

extracellular and other stimuli cause a variety of cellular responses to occur inside the cell. Examples of such stimuli include environmental and chemical stress signals (e.g. osmotic shock, heat shock, ultraviolet radiation, bacterial endotoxin, H<sub>2</sub>O<sub>2</sub>), cytokines (e.g. interleukin-1 (IL-1) and tumor necrosis factor  $\alpha$  (TNF- $\alpha$ )), and growth factors (e.g. granulocyte macrophage-colony-stimulating factor (GM-CSF), and fibroblast growth factor (FGF)). An extracellular stimulus may effect one or more cellular responses related to cell growth, migration, differentiation, secretion of hormones, activation of transcription factors, muscle contraction, glucose metabolism, control of protein synthesis and regulation of cell cycle.

Many diseases are associated with abnormal cellular responses triggered by protein kinase-mediated events. These diseases include autoimmune diseases, inflammatory diseases, neurological and neurodegenerative diseases, cancer, cardiovascular diseases, allergies and asthma, Alzheimer's disease or hormone-related diseases. Accordingly, there has been a substantial effort in medicinal chemistry to find protein kinase inhibitors that are effective as therapeutic agents.

Aurora-2 is a serine/threonine protein kinase that has been implicated in human cancer, such as colon, breast and other solid tumors. This kinase is believed to be involved in protein phosphorylation events that regulate the cell cycle. Specifically, Aurora-2 may play a role in controlling the accurate segregation of chromosomes during mitosis. Misregulation of the cell cycle can lead to cellular proliferation and other abnormalities. In human colon cancer tissue, the aurora-2 protein has been found to be overexpressed. See Bischoff et al., *EMBO J.*, 1998, 17, 3052-3065; Schumacher

et al., *J. Cell Biol.*, 1998, 143, 1635-1646; Kimura et al., *J. Biol. Chem.*, 1997, 272, 13766-13771.

Glycogen synthase kinase-3 (GSK-3) is a serine/threonine protein kinase comprised of  $\alpha$  and  $\beta$  isoforms that are each encoded by distinct genes [Coghlan et al., *Chemistry & Biology*, 7, 793-803 (2000); Kim and Kimmel, *Curr. Opinion Genetics Dev.*, 10, 508-514 (2000)]. GSK-3 has been implicated in various diseases including diabetes, Alzheimer's disease, CNS disorders such as manic depressive disorder and neurodegenerative diseases, and cardiomyocyte hypertrophy [WO 99/65897; WO 00/38675; and Haq et al., *J. Cell Biol.* (2000) 151, 117]. These diseases may be caused by, or result in, the abnormal operation of certain cell signaling pathways in which GSK-3 plays a role. GSK-3 has been found to phosphorylate and modulate the activity of a number of regulatory proteins. These proteins include glycogen synthase which is the rate limiting enzyme necessary for glycogen synthesis, the microtubule associated protein Tau, the gene transcription factor  $\beta$ -catenin, the translation initiation factor eIF2B, as well as ATP citrate lyase, axin, heat shock factor-1, c-Jun, c-Myc, c-Myb, CREB, and CEPB $\alpha$ . These diverse protein targets implicate GSK-3 in many aspects of cellular metabolism, proliferation, differentiation and development.

In a GSK-3 mediated pathway that is relevant for the treatment of type II diabetes, insulin-induced signaling leads to cellular glucose uptake and glycogen synthesis. Along this pathway, GSK-3 is a negative regulator of the insulin-induced signal. Normally, the presence of insulin causes inhibition of GSK-3 mediated phosphorylation and deactivation of glycogen synthase. The inhibition of GSK-3 leads to increased glycogen synthesis and glucose uptake [Klein et al., *PNAS*, 93,



8455-9 (1996); Cross et al., *Biochem. J.*, 303, 21-26 (1994); Cohen, *Biochem. Soc. Trans.*, 21, 555-567 (1993); Massillon et al., *Biochem J.* 299, 123-128 (1994)].

However, in a diabetic patient where the insulin response is impaired, glycogen synthesis and glucose uptake fail to increase despite the presence of relatively high blood levels of insulin. This leads to abnormally high blood levels of glucose with acute and long term effects that may ultimately result in cardiovascular disease, renal failure and blindness. In such patients, the normal insulin-induced inhibition of GSK-3 fails to occur. It has also been reported that in patients with type II diabetes, GSK-3 is overexpressed [WO 00/38675].

Therapeutic inhibitors of GSK-3 are therefore potentially useful for treating diabetic patients suffering from an impaired response to insulin.

GSK-3 activity has also been associated with Alzheimer's disease. This disease is characterized by the well-known  $\beta$ -amyloid peptide and the formation of intracellular neurofibrillary tangles. The neurofibrillary tangles contain hyperphosphorylated Tau protein where Tau is phosphorylated on abnormal sites. GSK-3 has been shown to phosphorylate these abnormal sites in cell and animal models. Furthermore, inhibition of GSK-3 has been shown to prevent hyperphosphorylation of Tau in cells [Lovestone et al., *Current Biology* 4, 1077-86 (1994); Brownlees et al., *Neuroreport* 8, 3251-55 (1997)]. Therefore, it is believed that GSK-3 activity may promote generation of the neurofibrillary tangles and the progression of Alzheimer's disease.

Another substrate of GSK-3 is  $\beta$ -catenin which is degraded after phosphorylation by GSK-3. Reduced levels of  $\beta$ -catenin have been reported in schizophrenic patients and have also been associated with other

diseases related to increase in neuronal cell death [Zhong et al., *Nature*, 395, 698-702 (1998); Takashima et al., *PNAS*, 90, 7789-93 (1993); Pei et al., *J. Neuropathol. Exp*, 56, 70-78 (1997)].


5           As a result of the biological importance of GSK-3, there is current interest in therapeutically effective GSK-3 inhibitors. Small molecules that inhibit GSK-3 have recently been reported [WO 99/65897 (Chiron) and WO 00/38675 (SmithKline Beecham)].

10           For many of the aforementioned diseases associated with abnormal GSK-3 activity, other protein kinases have also been targeted for treating the same diseases. However, the various protein kinases often act through different biological pathways. For example,  
15   certain quinazoline derivatives have been reported recently as inhibitors of p38 kinase (WO 00/12497 to Scios). The compounds are reported to be useful for treating conditions characterized by enhanced p38- $\alpha$  activity and/or enhanced TGF- $\beta$  activity. While p38  
20   activity has been implicated in a wide variety of diseases, including diabetes, p38 kinase is not reported to be a constituent of an insulin signaling pathway that regulates glycogen synthesis or glucose uptake. Therefore, unlike GSK-3, p38 inhibition would not be  
25   expected to enhance glycogen synthesis and/or glucose uptake.

          There is a continued need to find new therapeutic agents to treat human diseases. The protein kinases aurora-2 and GSK-3 are especially attractive  
30   targets for the discovery of new therapeutics due to their important role in cancer, diabetes, Alzheimer's disease and other diseases.

DESCRIPTION OF THE INVENTION

It has now been found that compounds of this invention and pharmaceutical compositions thereof are effective as protein kinase inhibitors, particularly as inhibitors of aurora-2 and GSK-3. These compounds have the general formula I:



I


10

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:


Z<sup>1</sup> to Z<sup>4</sup> are as described below;

Ring A is selected from the group consisting of:

15



20



G is Ring C or Ring D;

- 5 Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from  $-R^1$ , any substitutable non-ortho carbon position on Ring C is independently substituted by  $-R^5$ , and two adjacent substituents on
- 10 Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo,
- 15 oxo, or  $-R^8$ ;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or

20 heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or  $-R^5$ , and at any substitutable ring nitrogen by  $-R^4$ , provided that when Ring D is a six-membered aryl or

25 heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^1$  is selected from -halo, -CN,  $-\text{NO}_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $\text{C}_{1-6}$  aliphatic group, said phenyl, heteroaryl,

30 and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo,

oxo, or  $-R^8$ , said  $C_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

$R^x$  and  $R^y$  are independently selected from  $T-R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 0-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring formed by  $R^x$  and  $R^y$  is substituted by oxo or  $T-R^3$ , and any substitutable nitrogen on said ring formed by  $R^x$  and  $R^y$  is substituted by  $R^4$ ;

T is a valence bond or a  $C_{1-4}$  alkylidene chain;

5  $R^2$  and  $R^{2'}$  are independently selected from  $-R$ ,  $-T-W-R^6$ , or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each  
10 substitutable carbon on said fused ring formed by  $R^2$  and  $R^{2'}$  is substituted by halo, oxo,  $-CN$ ,  $-NO_2$ ,  $-R^7$ , or  $-V-R^6$ , and any substitutable nitrogen on said ring formed by  $R^2$  and  $R^{2'}$  is substituted by  $R^4$ ;

$R^3$  is selected from  $-R$ ,  $-halo$ ,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-COCH_2COR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-S(O)_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^7)_2$ ,  $-SO_2N(R^7)_2$ ,  $-OC(=O)R$ ,  $-N(R^7)COR$ ,  $-N(R^7)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^7)CON(R^7)_2$ ,  $-N(R^7)SO_2N(R^7)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^7)_2$ ;

each R is independently selected from hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-\text{COR}^7$ ,  $-\text{CO}_2(\text{C}_{1-6}$  aliphatic),  $-\text{CON}(\text{R}^7)_2$ , or  $-\text{SO}_2\text{R}^7$ , or two  $R^4$  on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or heteroaryl ring;

- 5 each  $R^5$  is independently selected from  $-R$ , halo,  $-\text{OR}$ ,  $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{SO}_2\text{R}$ ,  $-\text{SR}$ ,  $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^4)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^4)\text{COR}$ ,  $-\text{N}(\text{R}^4)\text{CO}_2$  (optionally substituted  $\text{C}_{1-6}$  aliphatic),  $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$ ,  
 10  $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$ , or  $R^5$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

V is  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2-$ ,  $-\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{N}(\text{R}^6)\text{CO}-$ ,  $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  
 15  $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ , or  
 20  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ;

W is  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$ ,  $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  
 25  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ , or  $-\text{CON}(\text{R}^6)-$ ;

each  $R^6$  is independently selected from hydrogen or an optionally substituted  $\text{C}_{1-4}$  aliphatic group, or two  $R^6$  groups on the same nitrogen atom are taken together  
 30 with the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;

each  $R^7$  is independently selected from hydrogen or an optionally substituted  $\text{C}_{1-6}$  aliphatic group, or two  $R^7$  on the same nitrogen are taken together with the

nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring;

each R<sup>8</sup> is independently selected from an optionally

substituted C<sub>1-4</sub> aliphatic group, -OR<sup>6</sup>, -SR<sup>6</sup>, -COR<sup>6</sup>,  
5 -SO<sub>2</sub>R<sup>6</sup>, -N(R<sup>6</sup>)<sub>2</sub>, -N(R<sup>6</sup>)N(R<sup>6</sup>)<sub>2</sub>, -CN, -NO<sub>2</sub>, -CON(R<sup>6</sup>)<sub>2</sub>, or  
-CO<sub>2</sub>R<sup>6</sup>; and

R<sup>9</sup> is selected from -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR,

-NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>,

-SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally  
10 substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>,  
-C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or  
-OC(=O)N(R<sup>4</sup>)<sub>2</sub>.

As used herein, the following definitions shall  
apply unless otherwise indicated. The phrase "optionally  
15 substituted" is used interchangeably with the phrase  
"substituted or unsubstituted" or with the term  
"(un)substituted." Unless otherwise indicated, an  
optionally substituted group may have a substituent at  
each substitutable position of the group, and each  
20 substitution is independent of the other.

The term "aliphatic" as used herein means  
straight-chain, branched or cyclic C<sub>1</sub>-C<sub>12</sub> hydrocarbons  
which are completely saturated or which contain one or  
more units of unsaturation but which are not aromatic.

25 For example, suitable aliphatic groups include  
substituted or unsubstituted linear, branched or cyclic  
alkyl, alkenyl, alkynyl groups and hybrids thereof such  
as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or  
(cycloalkyl)alkenyl. The terms "alkyl", "alkoxy",  
30 "hydroxyalkyl", "alkoxyalkyl", and "alkoxycarbonyl", used  
alone or as part of a larger moiety includes both  
straight and branched chains containing one to twelve  
carbon atoms. The terms "alkenyl" and "alkynyl" used  
alone or as part of a larger moiety shall include both

straight and branched chains containing two to twelve carbon atoms. The term "cycloalkyl" used alone or as part of a larger moiety shall include cyclic C<sub>3</sub>-C<sub>12</sub> hydrocarbons which are completely saturated or which  
5 contain one or more units of unsaturation, but which are not aromatic.

The terms "haloalkyl", "haloalkenyl" and "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. The  
10 term "halogen" means F, Cl, Br, or I.

The term "heteroatom" means nitrogen, oxygen, or sulfur and includes any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen. Also the term "nitrogen" includes a substitutable  
15 nitrogen of a heterocyclic ring. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR<sup>+</sup> (as in N-substituted  
20 pyrrolidinyl).

The terms "carbocycle", "carbocyclyl", "carbocyclo", or "carbocyclic" as used herein means an aliphatic ring system having three to fourteen members. The terms "carbocycle", "carbocyclyl", "carbocyclo", or  
25 "carbocyclic" whether saturated or partially unsaturated, also refers to rings that are optionally substituted. The terms "carbocycle", "carbocyclyl", "carbocyclo", or "carbocyclic" also include aliphatic rings that are fused to one or more aromatic or nonaromatic rings, such as in  
30 a decahydronaphthyl or tetrahydronaphthyl, where the radical or point of attachment is on the aliphatic ring.

The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to aromatic ring groups having



five to fourteen members, such as phenyl, benzyl, phenethyl, 1-naphthyl, 2-naphthyl, 1-anthracyl and 2-anthracyl. The term "aryl" also refers to rings that are optionally substituted. The term "aryl" may be used interchangeably with the term "aryl ring". "Aryl" also includes fused polycyclic aromatic ring systems in which an aromatic ring is fused to one or more rings. Examples include 1-naphthyl, 2-naphthyl, 1-anthracyl and 2-anthracyl. Also included within the scope of the term "aryl", as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings, such as in an indanyl, phenanthridinyl, or tetrahydronaphthyl, where the radical or point of attachment is on the aromatic ring.

The term "heterocycle", "heterocyclyl", or "heterocyclic" as used herein includes non-aromatic ring systems having five to fourteen members, preferably five to ten, in which one or more ring carbons, preferably one to four, are each replaced by a heteroatom such as N, O, or S. Examples of heterocyclic rings include 3-1H-benzimidazol-2-one, (1-substituted)-2-oxo-benzimidazol-3-yl, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydropyranyl, 3-tetrahydropyranyl, 4-tetrahydropyranyl, [1,3]-dioxalanyl, [1,3]-dithiolanyl, [1,3]-dioxanyl, 2-tetrahydrothiophenyl, 3-tetrahydrothiophenyl, 2-morpholinyl, 3-morpholinyl, 4-morpholinyl, 2-thiomorpholinyl, 3-thiomorpholinyl, 4-thiomorpholinyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 4-thiazolidinyl, diazolonyl, N-substituted diazolonyl, 1-phthalimidinyl, benzoxanyl, benzopyrrolidinyl, benzopiperidinyl, benzoxolanyl, benzothiolanyl, and benzothianyl. Also included within the scope of the term

"heterocyclyl" or "heterocyclic", as it is used herein, is a group in which a non-aromatic heteroatom-containing ring is fused to one or more aromatic or non-aromatic rings, such as in an indolinyl, chromanyl, phenanthridinyl, or tetrahydroquinolinyl, where the radical or point of attachment is on the non-aromatic heteroatom-containing ring. The term "heterocycle", "heterocyclyl", or "heterocyclic" whether saturated or partially unsaturated, also refers to rings that are optionally substituted.

The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to heteroaromatic ring groups having five to fourteen members. Examples of heteroaryl rings include 2-furanyl, 3-furanyl, N-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxadiazolyl, 5-oxadiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 3-pyridazinyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 5-tetrazolyl, 2-triazolyl, 5-triazolyl, 2-thienyl, 3-thienyl, carbazolyl, benzimidazolyl, benzothienyl, benzofuranyl, indolyl, quinolinyl, benzotriazolyl, benzothiazolyl, benzooxazolyl, benzimidazolyl, isoquinolinyl, indolyl, isoindolyl, acridinyl, or benzoisoxazolyl. Also included within the scope of the term "heteroaryl", as it is used herein, is a group in which a heteroatomic ring is fused to one or more aromatic or nonaromatic rings where the radical or point of attachment is on the heteroaromatic ring. Examples include tetrahydroquinolinyl, tetrahydroisoquinolinyl, and pyrido[3,4-d]pyrimidinyl. The term "heteroaryl" also refers to rings that are optionally substituted. The term "heteroaryl" may be

used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents. Examples of suitable substituents on the unsaturated carbon atom of an aryl, heteroaryl, aralkyl, or heteroaralkyl group include a halogen,  $-R^o$ ,  $-OR^o$ ,  $-SR^o$ , 1,2-methylene-dioxy, 1,2-ethylenedioxy, protected OH (such as acyloxy), phenyl (Ph), substituted Ph,  $-O(Ph)$ , substituted  $-O(Ph)$ ,  $-CH_2(Ph)$ , substituted  $-CH_2(Ph)$ ,  $-CH_2CH_2(Ph)$ , substituted  $-CH_2CH_2(Ph)$ ,  $-NO_2$ ,  $-CN$ ,  $-N(R^o)_2$ ,  $-NR^oC(O)R^o$ ,  $-NR^oC(O)N(R^o)_2$ ,  $-NR^oCO_2R^o$ ,  $-NR^oNR^oC(O)R^o$ ,  $-NR^oNR^oC(O)N(R^o)_2$ ,  $-NR^oNR^oCO_2R^o$ ,  $-C(O)C(O)R^o$ ,  $-C(O)CH_2C(O)R^o$ ,  $-CO_2R^o$ ,  $-C(O)R^o$ ,  $-C(O)N(R^o)_2$ ,  $-OC(O)N(R^o)_2$ ,  $-S(O)_2R^o$ ,  $-SO_2N(R^o)_2$ ,  $-S(O)R^o$ ,  $-NR^oSO_2N(R^o)_2$ ,  $-NR^oSO_2R^o$ ,  $-C(=S)N(R^o)_2$ ,  $-C(=NH)-N(R^o)_2$ ,  $-(CH_2)_yNHC(O)R^o$ ,  $-(CH_2)_yNHC(O)CH(V-R^o)(R^o)$ ; wherein  $R^o$  is hydrogen, a substituted or unsubstituted aliphatic group, an unsubstituted heteroaryl or heterocyclic ring, phenyl (Ph), substituted Ph,  $-O(Ph)$ , substituted  $-O(Ph)$ ,  $-CH_2(Ph)$ , or substituted  $-CH_2(Ph)$ ;  $y$  is 0-6; and  $V$  is a linker group. Examples of substituents on the aliphatic group or the phenyl ring of  $R^o$  include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, haloalkoxy, or haloalkyl.

An aliphatic group or a non-aromatic heterocyclic ring may contain one or more substituents. Examples of suitable substituents on the saturated carbon of an aliphatic group or of a non-aromatic heterocyclic

ring include those listed above for the unsaturated carbon of an aryl or heteroaryl group and the following: =O, =S, =NNHR<sup>+</sup>, =NN(R<sup>+</sup>)<sub>2</sub>, =N-, =NNHC(O)R<sup>+</sup>, =NNHCO<sub>2</sub>(alkyl), =NNHSO<sub>2</sub>(alkyl), or =NR<sup>+</sup>, where each R<sup>+</sup> is independently  
5 selected from hydrogen, an unsubstituted aliphatic group or a substituted aliphatic group. Examples of substituents on the aliphatic group include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl,  
10 alkylaminocarbonyloxy, dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, haloalkoxy, or haloalkyl.

Suitable substituents on the nitrogen of a non-aromatic heterocyclic ring include -R<sup>+</sup>, -N(R<sup>+</sup>)<sub>2</sub>, -C(O)R<sup>+</sup>,  
15 -CO<sub>2</sub>R<sup>+</sup>, -C(O)C(O)R<sup>+</sup>, -C(O)CH<sub>2</sub>C(O)R<sup>+</sup>, -SO<sub>2</sub>R<sup>+</sup>, -SO<sub>2</sub>N(R<sup>+</sup>)<sub>2</sub>, -C(=S)N(R<sup>+</sup>)<sub>2</sub>, -C(=NH)-N(R<sup>+</sup>)<sub>2</sub>, and -NR<sup>+</sup>SO<sub>2</sub>R<sup>+</sup>; wherein R<sup>+</sup> is hydrogen, an aliphatic group, a substituted aliphatic group, phenyl (Ph), substituted Ph,  
-O(Ph), substituted -O(Ph), CH<sub>2</sub>(Ph), substituted CH<sub>2</sub>(Ph),  
20 or an unsubstituted heteroaryl or heterocyclic ring. Examples of substituents on the aliphatic group or the phenyl ring include amino, alkylamino, dialkylamino, aminocarbonyl, halogen, alkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylaminocarbonyloxy,  
25 dialkylaminocarbonyloxy, alkoxy, nitro, cyano, carboxy, alkoxycarbonyl, alkylcarbonyl, hydroxy, haloalkoxy, or haloalkyl.

The term "linker group" or "linker" means an organic moiety that connects two parts of a compound.  
30 Linkers are typically comprised of an atom such as oxygen or sulfur, a unit such as -NH-, -CH<sub>2</sub>-, -C(O)-, -C(O)NH-, or a chain of atoms, such as an alkylidene chain. The molecular mass of a linker is typically in the range of about 14 to 200, preferably in the range of 14 to 96 with

a length of up to about six atoms. Examples of linkers include a saturated or unsaturated C<sub>1-6</sub> alkylidene chain which is optionally substituted, and wherein one or two saturated carbons of the chain are optionally replaced by  
5 -C(O)-, -C(O)C(O)-, -CONH-, -CONHNH-, -CO<sub>2</sub>-, -OC(O)-, -NHCO<sub>2</sub>-, -O-, -NHCONH-, -OC(O)NH-, -NHNH-, -NHCO-, -S-, -SO-, -SO<sub>2</sub>-, -NH-, -SO<sub>2</sub>NH-, or -NHSO<sub>2</sub>-.

The term "alkylidene chain" refers to an optionally substituted, straight or branched carbon chain  
10 that may be fully saturated or have one or more units of unsaturation. The optional substituents are as described above for an aliphatic group.

A combination of substituents or variables is permissible only if such a combination results in a  
15 stable or chemically feasible compound. A stable compound or chemically feasible compound is one in which the chemical structure is not substantially altered when kept at a temperature of 40 °C or less, in the absence of moisture or other chemically reactive conditions, for at  
20 least a week.

Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical  
25 isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically  
30 enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a <sup>13</sup>C- or <sup>14</sup>C-enriched carbon are within the scope of this invention.

Compounds of formula I or salts thereof may be formulated into compositions. In a preferred embodiment, the composition is a pharmaceutical composition. In one embodiment, the composition comprises an amount of the protein kinase inhibitor effective to inhibit a protein kinase, particularly GSK-3, in a biological sample or in a patient. In another embodiment, compounds of this invention and pharmaceutical compositions thereof, which comprise an amount of the protein kinase inhibitor effective to treat or prevent a GSK-3-mediated condition and a pharmaceutically acceptable carrier, adjuvant, or vehicle, may be formulated for administration to a patient.

The term "GSK-3-mediated condition" or "disease", as used herein, means any disease or other deleterious condition or state in which GSK-3 is known to play a role. Such diseases or conditions include, without limitation, diabetes, Alzheimer's disease, Huntington's Disease, Parkinson's Disease, AIDS-associated dementia, amyotrophic lateral sclerosis (AML), multiple sclerosis (MS), schizophrenia, cardiomyocyte hypertrophy, reperfusion/ischemia, and baldness.

One aspect of this invention relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, which method comprises administering to the patient a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof. This method is especially useful for diabetic patients. Another method relates to inhibiting the production of hyperphosphorylated Tau protein, which is useful in halting or slowing the progression of Alzheimer's disease. Another method relates to inhibiting the

phosphorylation of  $\beta$ -catenin, which is useful for treating schizophrenia.

Another aspect of the invention relates to inhibiting GSK-3 activity in a biological sample, which method comprises contacting the biological sample with a  
5 GSK-3 inhibitor of formula I.

Another aspect of this invention relates to a method of inhibiting Aurora-2 activity in a patient, which method comprises administering to the patient a  
10 compound of formula I or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing an Aurora-2-mediated disease with an Aurora-2 inhibitor, which method  
15 comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

The term "Aurora-2-mediated condition" or  
20 "disease", as used herein, means any disease or other deleterious condition in which Aurora is known to play a role. The term "Aurora-2-mediated condition" or "disease" also means those diseases or conditions that are alleviated by treatment with an Aurora-2 inhibitor.  
25 Such conditions include, without limitation, cancer. The term "cancer" includes, but is not limited to the following cancers: colon and ovarian.

Another aspect of the invention relates to inhibiting Aurora-2 activity in a biological sample,  
30 which method comprises contacting the biological sample with the Aurora-2 inhibitor of formula I, or a composition thereof.

Another aspect of this invention relates to a method of treating or preventing a CDK-2-mediated

diseases with a CDK-2 inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

5           The term "CDK-2-mediated condition" or "disease", as used herein, means any disease or other deleterious condition in which CDK-2 is known to play a role. The term "CDK-2-mediated condition" or "disease" also means those diseases or conditions that are  
10 alleviated by treatment with a CDK-2 inhibitor. Such conditions include, without limitation, cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune  
15 diseases such as rheumatoid arthritis. See Fischer, P.M. and Lane, D.P., *Current Medicinal Chemistry*, 7, 1213-1245 (2000); Mani, S., Wang, C., Wu, K., Francis, R. and Pestell, R., *Exp. Opin. Invest. Drugs*, 9, 1849 (2000); Fry, D.W. and Garrett, M.D., *Current Opinion in*  
20 *Oncologic, Endocrine & Metabolic Investigational Drugs*, 2, 40-59 (2000).

Another aspect of the invention relates to inhibiting CDK-2 activity in a biological sample or a patient, which method comprises administering to the  
25 patient a compound of formula I or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing an ERK-2-mediated diseases with an ERK-2 inhibitor, which method comprises  
30 administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

The term "ERK-mediated condition", as used herein means any disease state or other deleterious



condition in which ERK is known to play a role. The term "ERK-2-mediated condition" or "disease" also means those diseases or conditions that are alleviated by treatment with a ERK-2 inhibitor. Such conditions include, without  
5 limitation, cancer, stroke, diabetes, hepatomegaly, cardiovascular disease including cardiomegaly, Alzheimer's disease, cystic fibrosis, viral disease, autoimmune diseases, atherosclerosis, restenosis, psoriasis, allergic disorders including asthma,  
10 inflammation, neurological disorders and hormone-related diseases. The term "cancer" includes, but is not limited to the following cancers: breast, ovary, cervix, prostate, testis, genitourinary tract, esophagus, larynx, glioblastoma, neuroblastoma, stomach, skin,  
15 keratoacanthoma, lung, epidermoid carcinoma, large cell carcinoma, small cell carcinoma, lung adenocarcinoma, bone, colon, adenoma, pancreas, adenocarcinoma, thyroid, follicular carcinoma, undifferentiated carcinoma, papillary carcinoma, seminoma, melanoma, sarcoma, bladder  
20 carcinoma, liver carcinoma and biliary passages, kidney carcinoma, myeloid disorders, lymphoid disorders, Hodgkin's, hairy cells, buccal cavity and pharynx (oral), lip, tongue, mouth, pharynx, small intestine, colon-rectum, large intestine, rectum, brain and central  
25 nervous system, and leukemia. ERK-2 protein kinase and its implication in various diseases has been described [Bokemeyer et al. 1996, *Kidney Int.* 49, 1187; Anderson et al., 1990, *Nature* 343, 651; Crews et al., 1992, *Science* 258, 478; Bjorbaek et al., 1995, *J. Biol. Chem.* 270, 18848; Rouse et al., 1994, *Cell* 78, 1027; Raingeaud et al., 1996, *Mol. Cell Biol.* 16, 1247; Raingeaud et al. 1996; Chen et al., 1993 *Proc. Natl. Acad. Sci. USA* 90, 10952; Oliver et al., 1995, *Proc. Soc. Exp. Biol. Med.* 210, 162; Moodie et al., 1993, *Science* 260, 1658; Frey

and Mulder, 1997, *Cancer Res.* 57, 628; Sivaraman et al., 1997, *J Clin. Invest.* 99, 1478; Whelchel et al., 1997, *Am. J. Respir. Cell Mol. Biol.* 16, 589].

Another aspect of the invention relates to  
5 inhibiting ERK-2 activity in a biological sample or a patient, which method comprises administering to the patient a compound of formula I or a composition comprising said compound.

Another aspect of this invention relates to a  
10 method of treating or preventing an AKT-mediated diseases with an AKT inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

15 The term "AKT-mediated condition", as used herein, means any disease state or other deleterious condition in which AKT is known to play a role. The term "AKT-mediated condition" or "disease" also means those diseases or conditions that are alleviated by treatment  
20 with a AKT inhibitor. AKT-mediated diseases or conditions include, but are not limited to, proliferative disorders, cancer, and neurodegenerative disorders. The association of AKT, also known as protein kinase B, with various diseases has been described [Khwaja, A., *Nature*,  
25 pp. 33-34, 1990; Zang, Q. Y., et al, *Oncogene*, 19 2000; Kazuhiko, N., et al, *The Journal of Neuroscience*, 20 2000].

Another aspect of the invention relates to  
inhibiting AKT activity in a biological sample or a  
30 patient, which method comprises administering to the patient a compound of formula I or a composition comprising said compound.

Another aspect of this invention relates to a method of treating or preventing a Src-mediated disease

with a Src inhibitor, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a compound of formula I or a pharmaceutical composition thereof.

5           The term "Src-mediated condition", as used herein means any disease state or other deleterious condition in which Src is known to play a role. The term "Src-mediated condition" or "disease" also means those diseases or conditions that are alleviated by treatment  
10 with a Src inhibitor. Such conditions include, without limitation, hypercalcemia, osteoporosis, osteoarthritis, cancer, symptomatic treatment of bone metastasis, and Paget's disease. Src protein kinase and its implication in various diseases has been described [Soriano, *Cell*,  
15 69, 551 (1992); Soriano et al., *Cell*, 64, 693 (1991); Takayanagi, *J. Clin. Invest.*, 104, 137 (1999); Boschelli, *Drugs of the Future* 2000, 25(7), 717, (2000); Talamonti, *J. Clin. Invest.*, 91, 53 (1993); Lutz, *Biochem. Biophys. Res.* 243, 503 (1998); Rosen, *J. Biol. Chem.*, 261, 13754  
20 (1986); Bolen, *Proc. Natl. Acad. Sci. USA*, 84, 2251 (1987); Masaki, *Hepatology*, 27, 1257 (1998); Biscardi, *Adv. Cancer Res.*, 76, 61 (1999); Lynch, *Leukemia*, 7, 1416 (1993); Wiener, *Clin. Cancer Res.*, 5, 2164 (1999); Staley, *Cell Growth Diff.*, 8, 269 (1997)].

25           Another aspect of the invention relates to inhibiting Src activity in a biological sample or a patient, which method comprises administering to the patient a compound of formula I or a composition comprising said compound.

30           The term "pharmaceutically acceptable carrier, adjuvant, or vehicle" refers to a non-toxic carrier, adjuvant, or vehicle that may be administered to a patient, together with a compound of this invention, and

which does not destroy the pharmacological activity thereof.

The term "patient" includes human and veterinary subjects.

5           The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; preparations of an enzyme suitable for *in vitro* assay; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen,  
10       tears, or other body fluids or extracts thereof.

          The amount effective to inhibit protein kinase, for example, GSK-3 and Aurora-2, is one that measurably inhibits the kinase activity where compared to the activity of the enzyme in the absence of an inhibitor.  
15       Any method may be used to determine inhibition, such as, for example, the Biological Testing Examples described below.

          Pharmaceutically acceptable carriers that may be used in these pharmaceutical compositions include, but  
20       are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts  
25       or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium  
30       carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

          The compositions of the present invention may be administered orally, parenterally, by inhalation

spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously.

Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers

which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

5           The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used  
10 include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active  
15 ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

          Alternatively, the pharmaceutical compositions of this invention may be administered in the form of  
20 suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials  
25 include cocoa butter, beeswax and polyethylene glycols.

          The pharmaceutical compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including  
30 diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

          Topical application for the lower intestinal tract can be effected in a rectal suppository formulation

(see above) or in a suitable enema formulation.

Topically-transdermal patches may also be used.

For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability,

fluorocarbons, and/or other conventional solubilizing or dispersing agents.

In addition to the compounds of this invention, pharmaceutically acceptable derivatives or prodrugs of  
5 the compounds of this invention may also be employed in compositions to treat or prevent the above-identified diseases or disorders.

A "pharmaceutically acceptable derivative or prodrug" means any pharmaceutically acceptable salt,  
10 ester, salt of an ester or other derivative of a compound of this invention which, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.  
15 Particularly favored derivatives or prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a patient (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or  
20 which enhance delivery of the parent compound to a biological compartment (e.g., the brain or lymphatic system) relative to the parent species.

Pharmaceutically acceptable prodrugs of the compounds of this invention include, without limitation,  
25 esters, amino acid esters, phosphate esters, metal salts and sulfonate esters.

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids  
30 and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate,



fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., magnesium), ammonium and  $N^+(C_{1-4} \text{ alkyl})_4$  salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

The amount of the protein kinase inhibitor that may be combined with the carrier materials to produce a single dosage form will vary depending upon the patient treated and the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and

the judgment of the treating physician and the severity of the particular disease being treated. The amount of the inhibitor will also depend upon the particular compound in the composition.

5            Depending upon the particular protein kinase-mediated condition to be treated or prevented, additional therapeutic agents, which are normally administered to treat or prevent that condition, may be administered together with the inhibitors of this invention. For  
10    example, in the treatment of diabetes other anti-diabetic agents may be combined with the GSK-3 inhibitors of this invention to treat diabetes. These agents include, without limitation, insulin or insulin analogues, in injectable or inhalation form, glitazones, alpha  
15    glucosidase inhibitors, biguanides, insulin sensitizers, and sulfonyl ureas.


          Other examples of agents the inhibitors of this invention may also be combined with include, without limitation, chemotherapeutic agents or other anti-  
20    proliferative agents such as adriamycin, dexamethasone, vincristine, cyclophosphamide, fluorouracil, topotecan, taxol, interferons, and platinum derivatives; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and  
25    sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase  
30    inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents

for treating liver disease such as corticosteroids,  
cholestyramine, interferons, and anti-viral agents;  
agents for treating blood disorders such as  
corticosteroids, anti-leukemic agents, and growth  
5 factors; and agents for treating immunodeficiency  
disorders such as gamma globulin.

Those additional agents may be administered  
separately from the protein kinase inhibitor-containing  
composition, as part of a multiple dosage regimen.

10 Alternatively, those agents may be part of a single  
dosage form, mixed together with the protein kinase  
inhibitor of this invention in a single composition.

Compounds of this invention may exist in  
alternative tautomeric forms, as in tautomers 1 and 2  
15 shown below. Unless otherwise indicated, the  
representation of either tautomer is meant to include the  
other.




20


1

2


$R^x$  and  $R^y$  (at positions  $Z^3$  and  $Z^4$ , respectively)  
may be taken together to form a fused ring, providing a  
bicyclic ring system containing Ring A. Preferred  $R^x/R^y$   
25 rings include a 5-, 6-, 7-, or 8-membered unsaturated or  
partially unsaturated ring having 0-2 heteroatoms,  
wherein said  $R^x/R^y$  ring is optionally substituted.  
Examples of Ring A systems are shown below by compounds  
I-A through I-DD, wherein  $Z^1$  is nitrogen or C( $R^9$ ) and  $Z^2$   
30 is nitrogen or C(H).




I-A




I-B




I-C




I-D




I-E




I-F




I-G




I-H




I-I




I-J




I-K




I-L




I-M




I-N




I-O




I-P




I-Q




I-R




I-S




I-T




I-U




I-V




I-W




I-X




I-Y




I-Z




I-AA



I-BB



I-CC



I-DD


Preferred bicyclic Ring A systems include I-A,  
 15 I-B, I-C, I-D, I-E, I-F, I-G, I-H, I-I, I-J, I-K, I-L,  
 and I-M, more preferably I-A, I-B, I-C, I-F, and I-H, and  
 most preferably I-A, I-B, and I-H.

In the monocyclic Ring A system, preferred  $R^x$  groups, when present, include hydrogen, alkyl- or dialkylamino, acetamido, or a  $C_{1-4}$  aliphatic group such as methyl, ethyl, cyclopropyl, isopropyl or t-butyl.

- 5 Preferred  $R^y$  groups, when present, include T- $R^3$  wherein T is a valence bond or a methylene, and  $R^3$  is -R,  $-N(R^4)_2$ , or -OR. Examples of preferred  $R^y$  include 2-pyridyl, 4-pyridyl, piperidinyl, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, alkyl- or dialkylamino, acetamido, optionally substituted phenyl such as phenyl or halo-substituted phenyl, and methoxymethyl.
- 10

- In the bicyclic Ring A system, the ring formed when  $R^x$  and  $R^y$  are taken together may be substituted or unsubstituted. Suitable substituents include -R, halo, -OR,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$ (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ , wherein R and  $R^4$  are as defined above. Preferred  $R^x/R^y$  ring substituents include -halo, -R, -OR, -COR,  $-CO_2R$ ,  $-CON(R^4)_2$ ,  $-CN$ , or  $-N(R^4)_2$  wherein R is hydrogen or an optionally substituted  $C_{1-6}$  aliphatic group.
- 15
- 20


- $R^2$  and  $R^{2'}$  may be taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring. Preferred fused rings include benzo, pyrido, pyrimido, and a partially unsaturated 6-membered carbocyclo ring, wherein said fused ring is optionally substituted. These are exemplified in the following formula I compounds having a pyrazole-containing bicyclic ring system:
- 25
- 30



Preferred substituents on the  $R^2/R^{2'}$  fused ring include one or more of the following: -halo,  $-N(R^4)_2$ ,  $-C_{1-3}$  alkyl,  $-C_{1-3}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-3}$  alkyl),  $-CO_2(C_{1-3}$  alkyl),  $-CN$ ,  $-SO_2(C_{1-3}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-3}$  alkyl),  $-NHC(O)(C_{1-3}$  alkyl),  $-C(O)NH_2$ , and  $-CO(C_{1-3}$  alkyl), wherein the  $(C_{1-3}$  alkyl) is most preferably methyl.

When the pyrazole ring system is monocyclic, preferred  $R^2$  groups include hydrogen,  $C_{1-4}$  aliphatic, alkoxy carbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred  $R^2$  substituents include methyl, cyclopropyl, ethyl, isopropyl, propyl, t-butyl, cyclopentyl, phenyl,  $CO_2H$ ,  $CO_2CH_3$ ,  $CH_2OH$ ,  $CH_2OCH_3$ ,  $CH_2CH_2CH_2OH$ ,  $CH_2CH_2CH_2OCH_3$ ,  $CH_2CH_2CH_2OCH_2Ph$ ,  $CH_2CH_2CH_2NH_2$ ,  $CH_2CH_2CH_2NHCOOC(CH_3)_3$ ,  $CONHCH(CH_3)_2$ ,  $CONHCH_2CH=CH_2$ ,  $CONHCH_2CH_2OCH_3$ ,  $CONHCH_2Ph$ ,  $CONH(cyclohexyl)$ ,  $CON(Et)_2$ ,  $CON(CH_3)CH_2Ph$ ,  $CONH(n-C_3H_7)$ ,  $CON(Et)CH_2CH_2CH_3$ ,  $CONHCH_2CH(CH_3)_2$ ,  $CON(n-C_3H_7)_2$ ,  $CO(3-methoxymethylpyrrolidin-1-yl)$ ,  $CONH(3-tolyl)$ ,  $CONH(4-tolyl)$ ,  $CONHCH_3$ ,  $CO(morpholin-1-yl)$ ,  $CO(4-methylpiperazin-1-yl)$ ,  $CONHCH_2CH_2OH$ ,  $CONH_2$ , and  $CO(piperidin-1-yl)$ . A preferred  $R^{2'}$  group is hydrogen.

An embodiment that is particularly useful for treating GSK3-mediated diseases relates to compounds of formula II:



II

or a pharmaceutically acceptable derivative or prodrug thereof, wherein;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from -R<sup>1</sup>, any substitutable non-ortho carbon position on Ring C is independently substituted by -R<sup>5</sup>, and two adjacent substituents on Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo, oxo, or -R<sup>8</sup>;

R<sup>1</sup> is selected from -halo, -CN, -NO<sub>2</sub>, T-V-R<sup>6</sup>, phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or C<sub>1-6</sub> aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or -R<sup>8</sup>, said C<sub>1-6</sub> aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or R<sup>1</sup> and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;



$R^x$  and  $R^y$  are independently selected from  $T-R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 0-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring formed by  $R^x$  and  $R^y$  is substituted by oxo or  $T-R^3$ , and any substitutable nitrogen on said ring formed by  $R^x$  and  $R^y$  is substituted by  $R^4$ ;

$T$  is a valence bond or a  $C_{1-4}$  alkylidene chain;

$R^2$  and  $R^{2'}$  are independently selected from  $-R$ ,  $-T-W-R^6$ , or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable carbon on said fused ring formed by  $R^2$  and  $R^{2'}$  is substituted by halo, oxo,  $-CN$ ,  $-NO_2$ ,  $-R^7$ , or  $-V-R^6$ , and any substitutable nitrogen on said ring formed by  $R^2$  and  $R^{2'}$  is substituted by  $R^4$ ;

$R^3$  is selected from  $-R$ ,  $-\text{halo}$ ,  $-\text{OR}$ ,  $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{COCH}_2\text{COR}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{S}(\text{O})_2\text{R}$ ,  $-\text{SR}$ ,  $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^7)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^7)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^7)\text{COR}$ ,  $-\text{N}(\text{R}^7)\text{CO}_2$ (optionally substituted  $C_{1-6}$  aliphatic),  $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^7)\text{CON}(\text{R}^7)_2$ ,  $-\text{N}(\text{R}^7)\text{SO}_2\text{N}(\text{R}^7)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^7)_2$ ;

each  $R$  is independently selected from hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-\text{COR}^7$ ,  $-\text{CO}_2$ (optionally substituted  $C_{1-6}$  aliphatic),  $-\text{CON}(\text{R}^7)_2$ , or  $-\text{SO}_2\text{R}^7$ , or two  $R^4$  on the same nitrogen are taken

together to form a 5-8 membered heterocyclyl or heteroaryl ring;

each  $R^5$  is independently selected from -R, halo, -OR,

-C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR,  
 5 -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR,  
 -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic),  
 -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>,  
 -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>, or  $R^5$  and

10 intervening atoms form said ring fused to Ring C;

V is -O-, -S-, -SO-, -SO<sub>2</sub>-, -N(R<sup>6</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>6</sup>)-,

-N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -N(R<sup>6</sup>)CO-, -N(R<sup>6</sup>)C(O)O-,

-N(R<sup>6</sup>)CON(R<sup>6</sup>)-, -N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)N(R<sup>6</sup>)-,

-C(O)N(R<sup>6</sup>)-, -OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-,

15 -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-,

-C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-,

-C(R<sup>6</sup>)=N-O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, or

-C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-;

W is -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-,

20 -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-,

-C(R<sup>6</sup>)OC(O)-, -C(R<sup>6</sup>)OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CO-,

-C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-, -C(R<sup>6</sup>)=N-O-,

-C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-,


-C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-, or -CON(R<sup>6</sup>)-;

25 each  $R^6$  is independently selected from hydrogen, an optionally substituted C<sub>1-4</sub> aliphatic group, or two  $R^6$  groups on the same nitrogen atom are taken together with the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;

30 each  $R^7$  is independently selected from hydrogen or an optionally substituted C<sub>1-6</sub> aliphatic group, or two  $R^7$  on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring; and


each  $R^8$  is independently selected from an optionally substituted  $C_{1-4}$  aliphatic group,  $-OR^6$ ,  $-SR^6$ ,  $-COR^6$ ,  $-SO_2R^6$ ,  $-N(R^6)_2$ ,  $-N(R^6)N(R^6)_2$ ,  $-CN$ ,  $-NO_2$ ,  $-CON(R^6)_2$ , or  $-CO_2R^6$ .

5 When the  $R^x$  and  $R^y$  groups of formula II are taken together to form a fused ring, preferred  $R^x/R^y$  rings include a 5-, 6-, 7-, or 8-membered unsaturated or partially unsaturated ring having 0-2 heteroatoms, wherein said  $R^x/R^y$  ring is optionally substituted. This  
10 provides a bicyclic ring system containing a pyrimidine ring. Examples of preferred pyrimidine ring systems of formula II are the mono- and bicyclic systems shown below.




15


II-A




II-B




II-C




II-D




II-E




II-F



II-G




II-H




II-I


20




II-J




II-K




II-L



II-M




II-N



II-O

5



II-P

10

More preferred pyrimidine ring systems of formula II include II-A, II-B, II-C, II-F, and II-H, most preferably II-A, II-B, and II-H.

In the monocyclic pyrimidine ring system of formula II, preferred  $R^x$  groups include hydrogen, alkyl- or dialkylamino, acetamido, or a  $C_{1-4}$  aliphatic group such as methyl, ethyl, cyclopropyl, isopropyl or t-butyl. Preferred  $R^y$  groups include  $T-R^3$  wherein T is a valence bond or a methylene, and  $R^3$  is  $-R$ ,  $-N(R^4)_2$ , or  $-OR$ . When  $R^3$  is  $-R$  or  $-OR$ , a preferred R is an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring. Examples of preferred  $R^y$  include 2-pyridyl, 4-pyridyl, piperidinyl, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, alkyl- or dialkylamino, acetamido, optionally substituted phenyl


15

20

such as phenyl or halo-substituted phenyl, and methoxymethyl.

In the bicyclic pyrimidine ring system of formula II, the ring formed when  $R^x$  and  $R^y$  are taken together may be substituted or unsubstituted. Suitable substituents include -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>, wherein R and R<sup>4</sup> are as defined above. Preferred  $R^x/R^y$  ring substituents include -halo, -R, -OR, -COR, -CO<sub>2</sub>R, -CON(R<sup>4</sup>)<sub>2</sub>, -CN, or -N(R<sup>4</sup>)<sub>2</sub> wherein R is an optionally substituted C<sub>1-6</sub> aliphatic group.

The  $R^2$  and  $R^{2'}$  groups of formula II may be taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring. Preferred fused rings include benzo, pyrido, pyrimido, and a partially unsaturated 6-membered carbocyclo ring. These are exemplified in the following formula II compounds having a pyrazole-containing bicyclic ring system:




Preferred substituents on the  $R^2/R^{2'}$  fused ring of formula II include one or more of the following: -halo, -N(R<sup>4</sup>)<sub>2</sub>, -C<sub>1-4</sub> alkyl, -C<sub>1-4</sub> haloalkyl, -NO<sub>2</sub>, -O(C<sub>1-4</sub> alkyl), -CO<sub>2</sub>(C<sub>1-4</sub> alkyl), -CN, -SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -OC(O)NH<sub>2</sub>, -NH<sub>2</sub>SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -NHC(O)(C<sub>1-4</sub> alkyl),

-C(O)NH<sub>2</sub>, and -CO(C<sub>1-4</sub> alkyl), wherein the (C<sub>1-4</sub> alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the (C<sub>1-4</sub> alkyl) group is methyl.

When the pyrazole ring system of formula II is monocyclic, preferred R<sup>2</sup> groups include hydrogen, a substituted or unsubstituted group selected from aryl, heteroaryl, or a C<sub>1-6</sub> aliphatic group. Examples of such preferred R<sup>2</sup> groups include methyl, t-butyl, -CH<sub>2</sub>OCH<sub>3</sub>, cyclopropyl, furanyl, thienyl, and phenyl. A preferred R<sup>2'</sup> group is hydrogen.

More preferred ring systems of formula II are the following, which may be substituted as described above, wherein R<sup>2</sup> and R<sup>2'</sup> are taken together with the pyrazole ring to form an indazole ring; and R<sup>x</sup> and R<sup>y</sup> are each methyl, or R<sup>x</sup> and R<sup>y</sup> are taken together with the pyrimidine ring to form a quinazoline or tetrahydroquinazoline ring:



Particularly preferred are those compounds of formula II-Aa, II-Ba, or II-Ha wherein ring C is a phenyl ring and R<sup>1</sup> is halo, methyl, or trifluoromethyl.

Preferred formula II Ring C groups are phenyl and pyridinyl. When two adjacent substituents on Ring C are taken together to form a fused ring, Ring C is contained in a bicyclic ring system. Preferred fused rings include a benzo or pyrido ring. Such rings

preferably are fused at ortho and meta positions of Ring C. Examples of preferred bicyclic Ring C systems include naphthyl, quinolinyl and isoquinolinyl.

An important feature of the formula II

- 5 compounds is the  $R^1$  ortho substituent on Ring C. An ortho position on Ring C or Ring D is defined relative to the position where Ring A is attached. Preferred  $R^1$  groups include -halo, an optionally substituted  $C_{1-6}$  aliphatic group, phenyl,  $-COR^6$ ,  $-OR^6$ ,  $-CN$ ,  $-SO_2R^6$ ,  $-SO_2NH_2$ ,  $-N(R^6)_2$ ,  
10  $-CO_2R^6$ ,  $-CONH_2$ ,  $-NHCOR^6$ ,  $-OC(O)NH_2$ , or  $-NHSO_2R^6$ . When  $R^1$  is an optionally substituted  $C_{1-6}$  aliphatic group, the most preferred optional substituents are halogen. Examples of preferred  $R^1$  groups include  $-CF_3$ ,  $-Cl$ ,  $-F$ ,  $-CN$ ,  $-COCH_3$ ,  
15  $-OCH_3$ ,  $-OH$ ,  $-CH_2CH_3$ ,  $-OCH_2CH_3$ ,  $-CH_3$ ,  $-CF_2CH_3$ , cyclohexyl, t-butyl, isopropyl, cyclopropyl,  $-C\equiv CH$ ,  $-C\equiv C-CH_3$ ,  $-SO_2CH_3$ ,  
 $-SO_2NH_2$ ,  $-N(CH_3)_2$ ,  $-CO_2CH_3$ ,  $-CONH_2$ ,  $-NHCOCH_3$ ,  $-OC(O)NH_2$ ,  
 $-NHSO_2CH_3$ , and  $-OCF_3$ .

- On Ring C of formula II, preferred  $R^5$  substituents, when present, include -halo,  $-CN$ ,  $-NO_2$ ,  
20  $-N(R^4)_2$ , optionally substituted  $C_{1-6}$  aliphatic group,  $-OR$ ,  $-C(O)R$ ,  $-CO_2R$ ,  $-CONH(R^4)$ ,  $-N(R^4)COR$ ,  $-SO_2N(R^4)_2$ , and  $-N(R^4)SO_2R$ . More preferred  $R^5$  substituents include  $-Cl$ ,  
 $-F$ ,  $-CN$ ,  $-CF_3$ ,  $-NH_2$ ,  $-NH(C_{1-4} \text{ aliphatic})$ ,  $-N(C_{1-4} \text{ aliphatic})_2$ ,  $-O(C_{1-4} \text{ aliphatic})$ ,  $C_{1-4} \text{ aliphatic}$ , and  
25  $-CO_2(C_{1-4} \text{ aliphatic})$ . Examples of such preferred  $R^5$  substituents include  $-Cl$ ,  $-F$ ,  $-CN$ ,  $-CF_3$ ,  $-NH_2$ ,  $-NHMe$ ,  
 $-NMe_2$ ,  $-OEt$ , methyl, ethyl, cyclopropyl, isopropyl, t-butyl, and  $-CO_2Et$ .

- Preferred formula II compounds have one or  
30 more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring

system, the bicyclic ring system is selected from a naphthyl, quinolinyl or isoquinolinyl ring;

(b)  $R^x$  is hydrogen or  $C_{1-4}$  aliphatic and  $R^y$  is  $T-R^3$ , or  $R^x$  and  $R^y$  are taken together with their  
5 intervening atoms to form an optionally substituted 5-7 membered unsaturated or partially unsaturated ring having 0-2 ring nitrogens;

(c)  $R^1$  is -halo, an optionally substituted  $C_{1-6}$  aliphatic group, phenyl,  $-COR^6$ ,  $-OR^6$ ,  $-CN$ ,  $-SO_2R^6$ ,  $-SO_2NH_2$ ,  
10  $-N(R^6)_2$ ,  $-CO_2R^6$ ,  $-CONH_2$ ,  $-NHCOR^6$ ,  $-OC(O)NH_2$ , or  $-NHCO_2R^6$ ; and

(d)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are  
15 taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring.

More preferred compounds of formula II have one or more, and more preferably all, of the features  
20 selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring;

(b)  $R^x$  is hydrogen or methyl and  $R^y$  is  $-R$ ,  $N(R^4)_2$ , or  $-OR$ , or  $R^x$  and  $R^y$  are taken together with their  
25 intervening atoms to form a 5-7 membered unsaturated or partially unsaturated carbocyclo ring optionally substituted with  $-R$ , halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  
30  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;



(c)  $R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or -CN;

(d)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, or  
5 a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring; and

(e) each  $R^5$  is independently selected from  
10 -halo, -CN, -NO<sub>2</sub>, -N( $R^4$ )<sub>2</sub>, optionally substituted  $C_{1-6}$  aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH( $R^4$ ), -N( $R^4$ )COR, -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, or -N( $R^4$ )SO<sub>2</sub>R.

Even more preferred compounds of formula II have one or more, and more preferably all, of the  
15 features selected from the group consisting of:

(a) Ring C is a phenyl ring optionally substituted by - $R^5$ ;

(b)  $R^x$  is hydrogen or methyl and  $R^y$  is methyl, methoxymethyl, ethyl, cyclopropyl, isopropyl, t-butyl,  
20 alkyl- or an optionally substituted group selected from 2-pyridyl, 4-pyridyl, piperidinyl, or phenyl, or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted benzo ring or partially unsaturated 6-membered carbocyclo ring;

25 (c)  $R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN;


(d)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring  
30 optionally substituted with -halo, -N( $R^4$ )<sub>2</sub>, - $C_{1-4}$  alkyl, - $C_{1-4}$  haloalkyl, -NO<sub>2</sub>, -O( $C_{1-4}$  alkyl), -CO<sub>2</sub>( $C_{1-4}$  alkyl), -CN, -SO<sub>2</sub>( $C_{1-4}$  alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -OC(O)NH<sub>2</sub>, -NH<sub>2</sub>SO<sub>2</sub>( $C_{1-4}$  alkyl), -NHC(O)( $C_{1-4}$  alkyl), -C(O)NH<sub>2</sub>, or -CO( $C_{1-4}$  alkyl), wherein

the (C<sub>1-4</sub> alkyl) is a straight, branched, or cyclic alkyl group; and


- (e) each R<sup>5</sup> is independently selected from -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic).

Representative compounds of formula II are shown below in Table 1.


10 Table 1.




II-1




II-2




II-3




II-4




II-5




II-6




II-7




II-8




II-9




II-10




II-11




II-12




II-13




II-14




II-15




II-16




II-17




II-18




II-19




II-20




II-21




II-22




II-23




II-24




II-25




II-26




II-27




II-28




II-29




II-30




II-31




II-32




II-33




II-34




II-35




II-36




II-37




II-38




II-39




II-40




II-41




II-42




II-43




II-44



II-45




II-46




II-47




II-48




II-49




II-50




II-51




II-52




II-53




II-54



II-55




II-56




II-57




II-58




II-59




II-60




II-61



II-62




II-63



II-64



II-65




II-66


5




II-67




II-68



II-69



II-70




II-71




II-72


10




II-73




II-74




II-75



II-76




II-77




II-78


5




II-79



II-80




II-81



II-82




II-83




II-84

10







II-85




II-86




II-87




II-88




II-89




II-90




II-91



II-92



II-93




II-94




II-95




II-96




II-97




II-98




II-99



II-100




II-101




II-102


5




II-103




II-104




II-105



II-106




II-107




II-108


10




II-109




II-110




II-111



II-112




II-113




II-114


5




II-115




II-116




II-117



II-118




II-119




II-120


10




II-121




II-122




II-123




II-124




II-125




II-126




II-127




II-128




II-129



II-130




II-131




II-132


10




II-133




II-134




II-135



II-136




II-137




II-138


5



II-139




II-140




II-141


10




II-142




II-143




II-144




II-145




II-146




II-147




II-148




II-149




II-150




II-151




II-152




II-153




II-154




II-155




II-156




II-157




II-158




II-159




II-160




II-161




II-162




II-163




II-164




II-165




II-166




II-167




II-168




II-169




II-170




II-171




II-172




II-173




II-174




II-175




II-176




II-177



II-178




II-179




II-180







II-181




II-182.




II-183



II-184




II-185




II-186


5




II-187




II-188




II-189



II-190




II-191




II-192


10




II-193




II-194




II-195




II-196




II-197




II-198




II-199




II-200




II-201




II-202




II-203




II-204




II-205




II-206




II-207




II-208




II-209




II-210




II-211




II-212




II-213




II-214




II-215




II-216




II-217




II-218




II-219




II-220




II-221



II-222




II-223




II-224




II-225



II-226




II-227




II-228

5


10



II-229




II-230



II-231



II-232




II-233




II-234


5




II-235




II-236




II-237



II-238




II-239




II-240

10




II-241




II-242




II-243




II-244




II-245




II-246




II-247




II-248



II-249



II-250



II-251

5

10

In another embodiment, this invention provides a composition comprising a compound of formula II and a pharmaceutically acceptable carrier.

One aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient,

comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula II.

Another aspect relates to a method of treating  
5 a disease that is alleviated by treatment with a GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula II.

10 Another aspect relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula  
15 II. This method is especially useful for diabetic patients.

Another aspect relates to a method of inhibiting the production of hyperphosphorylated Tau protein in a patient in need thereof, comprising  
20 administering to said patient a therapeutically effective amount of a composition comprising a compound of formula II. This method is especially useful in halting or slowing the progression of Alzheimer's disease.

Another aspect relates to a method of  
25 inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula II. This method is especially useful for treating schizophrenia.

30 One aspect of this invention relates to a method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula II.

Another aspect relates to a method of treating a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula II. This method is especially useful for treating cancer, such as colon, ovarian, and breast cancer.

One aspect of this invention relates to a method of inhibiting CDK-2 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula II.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a CDK-2 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula II. This method is especially useful for treating cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis.


Another method relates to inhibiting GSK-3, Aurora, or CDK-2 activity in a biological sample, which method comprises contacting the biological sample with the GSK-3 or Aurora inhibitor of formula II, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3, Aurora or CDK-2.

Each of the aforementioned methods directed to the inhibition of GSK-3, Aurora or CDK-2, or the treatment of a disease alleviated thereby, is preferably



carried out with a preferred compound of formula II, as described above.

Another embodiment of this invention relates to compounds of formula III:



III

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or -R<sup>5</sup>, and at any substitutable ring nitrogen by -R<sup>4</sup>, provided that when Ring D is a six-membered aryl or heteroaryl ring, -R<sup>5</sup> is hydrogen at each ortho carbon position of Ring D;

R<sup>x</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a fused, benzo ring or a 5-8 membered carbocyclo ring, wherein any substitutable carbon on said fused ring formed by R<sup>x</sup> and R<sup>y</sup> is substituted by oxo or T-R<sup>3</sup>;

T is a valence bond or a C<sub>1-4</sub> alkylidene chain;

R<sup>2</sup> and R<sup>2'</sup> are independently selected from -R, -T-W-R<sup>6</sup>, or R<sup>2</sup> and R<sup>2'</sup> are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each

substitutable carbon on said fused ring formed by  $R^2$  and  $R^{2'}$  is substituted by halo, oxo,  $-CN$ ,  $-NO_2$ ,  $-R^7$ , or  $-V-R^6$ , and any substitutable nitrogen on said ring formed by  $R^2$  and  $R^{2'}$  is substituted by  $R^4$ ;

- 5  $R^3$  is selected from  $-R$ ,  $-\text{halo}$ ,  $=O$ ,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-COCH_2COR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-S(O)_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  
 10  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;

each  $R$  is independently selected from hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-COR^7$ ,  $-CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-CON(R^7)_2$ , or  $-SO_2R^7$ , or two  $R^4$  on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or  
 15 heteroaryl ring;

each  $R^5$  is independently selected from  $-R$ , halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  
 20  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;

$V$  is  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-N(R^6)SO_2-$ ,  $-SO_2N(R^6)-$ ,  $-N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  $-N(R^6)CO-$ ,  $-N(R^6)C(O)O-$ ,  $-N(R^6)CON(R^6)-$ ,  $-N(R^6)SO_2N(R^6)-$ ,  $-N(R^6)N(R^6)-$ ,  
 25  $-C(O)N(R^6)-$ ,  $-OC(O)N(R^6)-$ ,  $-C(R^6)_2O-$ ,  $-C(R^6)_2S-$ ,  $-C(R^6)_2SO-$ ,  $-C(R^6)_2SO_2-$ ,  $-C(R^6)_2SO_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)C(O)-$ ,  $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  $-C(R^6)=N-O-$ ,  $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ , or  $-C(R^6)_2N(R^6)CON(R^6)-$ ;

W is  $-C(R^6)_2O-$ ,  $-C(R^6)_2S-$ ,  $-C(R^6)_2SO-$ ,  $-C(R^6)_2SO_2-$ ,  
 $-C(R^6)_2SO_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  
 $-C(R^6)OC(O)-$ ,  $-C(R^6)OC(O)N(R^6)-$ ,  $-C(R^6)_2N(R^6)CO-$ ,  
 $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  $-C(R^6)=N-O-$ ,  
5  $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ ,  
 $-C(R^6)_2N(R^6)CON(R^6)-$ , or  $-CON(R^6)-$ ;

each  $R^6$  is independently selected from hydrogen or an  
optionally substituted  $C_{1-4}$  aliphatic group, or two  $R^6$   
groups on the same nitrogen atom are taken together  
10 with the nitrogen atom to form a 5-6 membered  
heterocyclyl or heteroaryl ring; and

each  $R^7$  is independently selected from hydrogen or an  
optionally substituted  $C_{1-6}$  aliphatic group, or two  $R^7$   
on the same nitrogen are taken together with the  
15 nitrogen to form a 5-8 membered heterocyclyl or  
heteroaryl ring.


Preferred formula III Ring D monocyclic rings  
include substituted and unsubstituted phenyl, pyridinyl,  
piperidinyl, piperazinyl, pyrrolidinyl, thienyl,  
20 azepanyl, and morpholinyl rings. When two adjacent  
substituents on Ring D are taken together to form a fused  
ring, the Ring D system is bicyclic. Preferred formula  
III Ring D bicyclic rings include 1,2,3,4-  
tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl,  
25 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl,  
isoquinolinyl, quinolinyl, and naphthyl. Examples of  
more preferred bicyclic Ring D systems include naphthyl  
and isoquinolinyl.

Preferred  $R^5$  substituents on Ring D of formula  
30 III include halo, oxo, CN,  $-NO_2$ ,  $-N(R^4)_2$ ,  $-CO_2R$ ,  $-CONH(R^4)$ ,  
 $-N(R^4)COR$ ,  $-SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ ,  $-SR$ ,  $-OR$ ,  $-C(O)R$ , or  
substituted or unsubstituted group selected from 5-6  
membered heterocyclyl,  $C_{6-10}$  aryl, or  $C_{1-6}$  aliphatic. More  
preferred  $R^5$  substituents include -halo, -CN, -oxo, -SR,


-OR, -N(R<sup>4</sup>)<sub>2</sub>, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. Examples of Ring D substituents include -OH, phenyl, methyl, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH,

- 5 pyrrolidinyl, OPh, CF<sub>3</sub>, C≡CH, Cl, Br, F, I, NH<sub>2</sub>, C(O)CH<sub>3</sub>, i-propyl, tert-butyl, SEt, OMe, N(Me)<sub>2</sub>, methylene dioxy, and ethylene dioxy.


Preferred rings formed when the R<sup>x</sup> and R<sup>y</sup> groups of formula III are taken together to form a fused ring include a 5-, 6-, or 7-membered unsaturated or partially unsaturated carbocyclo ring, wherein any substitutable carbon on said fused ring is substituted by oxo or T-R<sup>3</sup>. Examples of preferred bicyclic ring systems are shown below.




III-A




III-B



III-C



III-F



III-I

20

Preferred substituents on the R<sup>x</sup>/R<sup>y</sup> fused ring of formula III include -R, oxo, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub> (optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or

-OC(=O)N(R<sup>4</sup>)<sub>2</sub>, wherein R and R<sup>4</sup> are as defined above.

More preferred substituents on the R<sup>x</sup>/R<sup>y</sup> fused ring include halo, CN, oxo, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, (C<sub>1-6</sub> alkyl)carbonyl, (C<sub>1-6</sub> alkyl)sulfonyl, mono- or

5 dialkylamino, mono- or dialkylaminocarbonyl, mono- or dialkylaminocarbonyloxy, or 5-6 membered heteroaryl.

Examples of such preferred substituents include methoxy, methyl, isopropyl, methylsulfonyl, cyano, chloro, pyrrolyl, methoxy, ethoxy, ethylamino, acetyl, and

10 acetamido.

Preferred R<sup>2</sup> substituents of formula III include hydrogen, C<sub>1-4</sub> aliphatic, alkoxycarbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl,

15 dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred R<sup>2</sup>

substituents include methyl, cyclopropyl, ethyl, isopropyl, propyl, t-butyl, cyclopentyl, phenyl, CO<sub>2</sub>H, CO<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>OH, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>,

20 CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>Ph, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHCOOC(CH<sub>3</sub>)<sub>3</sub>,


CONHCH(CH<sub>3</sub>)<sub>2</sub>, CONHCH<sub>2</sub>CH=CH<sub>2</sub>, CONHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CONHCH<sub>2</sub>Ph, CONH(cyclohexyl), CON(Et)<sub>2</sub>, CON(CH<sub>3</sub>)CH<sub>2</sub>Ph, CONH(n-C<sub>3</sub>H<sub>7</sub>), CON(Et)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, CONHCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, CON(n-C<sub>3</sub>H<sub>7</sub>)<sub>2</sub>, CO(3-

methoxymethylpyrrolidin-1-yl), CONH(3-tolyl), CONH(4-

25 tollyl), CONHCH<sub>3</sub>, CO(morpholin-1-yl), CO(4-methylpiperazin-1-yl), CONHCH<sub>2</sub>CH<sub>2</sub>OH, CONH<sub>2</sub>, and CO(piperidin-1-yl).

When the R<sup>2</sup> and R<sup>2'</sup> groups of formula III are taken together to form a ring, preferred R<sup>2</sup>/R<sup>2'</sup> ring systems containing the pyrazole ring include benzo,

30 pyrido, pyrimido, 3-oxo-2H-pyridazino, and a partially unsaturated 6-membered carbocyclo ring. Examples of such preferred R<sup>2</sup>/R<sup>2'</sup> ring systems containing the pyrazole ring include the following:



5 Preferred substituents on the  $R^2/R^{2'}$  fused ring of formula III include one or more of the following:

-halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl),  $-CN$ ,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  
 10  $-C(O)NH_2$ , and  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the  $(C_{1-4}$  alkyl) group is methyl.

Preferred formula III compounds have one or more, and more preferably all, of the features selected  
 15 from the group consisting of:

(a) Ring D is an optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-  
 20 tetrahydroquinolinyl, 2,3-dihydro-1*H*-isoindolyl, 2,3-dihydro-1*H*-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted benzo  
 25 ring or a 5-7 membered carbocyclo ring; and

(c)  $R^{2'}$  is hydrogen or methyl and  $R^2$  is T-W- $R^6$  or R, wherein W is  $-C(R^6)_2O-$ ,  $-C(R^6)_2N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  $-C(R^6)OC(O)-$ ,  $-C(R^6)_2N(R^6)CO-$ ,  $-C(R^6)_2N(R^6)C(O)O-$ , or  $-CON(R^6)-$ , and R is an optionally substituted group

selected from C<sub>1-6</sub> aliphatic or phenyl, or R<sup>2</sup> and R<sup>2'</sup> are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido, or partially unsaturated 6-membered carbocyclo ring.

5 More preferred compounds of formula III have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl,  
10 piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b) R<sup>x</sup> and R<sup>y</sup> are taken together with their  
15 intervening atoms to form a benzo ring or a 5-7 membered carbocyclo ring optionally substituted with -R, oxo, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic),  
20 -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>; and

(c) each R<sup>5</sup> is independently selected from halo, oxo, CN, NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -SR, -OR, -C(O)R, or a substituted  
25 or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic.

Even more preferred compounds of formula III have one or more, and more preferably all, of the features selected from the group consisting of:

30 (a) R<sup>x</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a benzo or 6-membered partially unsaturated carbocyclo ring optionally substituted with halo, CN, oxo, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, (C<sub>1-6</sub> alkyl)carbonyl, (C<sub>1-6</sub> alkyl)sulfonyl, mono- or dialkylamino, mono- or


dialkylaminocarbonyl, mono- or dialkylaminocarbonyloxy, or 5-6 membered heteroaryl;

(b) each  $R^5$  is independently selected from -halo, -CN, -oxo, -SR, -OR, -N( $R^4$ )<sub>2</sub>, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic; and


(c)  $R^{2'}$  is hydrogen and  $R^2$  is selected from  $R^{2'}$  is hydrogen or methyl and  $R^2$  is T-W- $R^6$  or R, wherein W is -C( $R^6$ )<sub>2</sub>O-, -C( $R^6$ )<sub>2</sub>N( $R^6$ )-, -CO-, -CO<sub>2</sub>-, -C( $R^6$ )OC(O)-, -C( $R^6$ )<sub>2</sub>N( $R^6$ )CO-, or -CON( $R^6$ )-, and R is an optionally substituted group selected from C<sub>1-6</sub> aliphatic or phenyl, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, or partially unsaturated 6-membered carbocyclo ring optionally substituted with -halo, -N( $R^4$ )<sub>2</sub>, -C<sub>1-4</sub> alkyl, -C<sub>1-4</sub> haloalkyl, -NO<sub>2</sub>, -O(C<sub>1-4</sub> alkyl), -CO<sub>2</sub>(C<sub>1-4</sub> alkyl), -CN, -SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -OC(O)NH<sub>2</sub>, -NH<sub>2</sub>SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -NHC(O)(C<sub>1-4</sub> alkyl), -C(O)NH<sub>2</sub>, or -CO(C<sub>1-4</sub> alkyl), wherein the (C<sub>1-4</sub> alkyl) is a straight, branched, or cyclic alkyl group.

Representative compounds of formula III are set forth in Table 2 below.


Table 2



III-1




III-2




III-3







III-4




III-5




III-6




III-7



III-8




III-9




III-10




III-11




III-12




III-13



III-14




III-15




III-16




III-17




III-18




III-19




III-20




III-21




III-22




III-23




III-24




III-25




III-26




III-27




III-28




III-29




III-30




III-31




III-32




III-33




III-34



III-35




III-36




III-37




III-38




III-39




III-40




III-41




III-42




III-43




III-44




III-45




III-46




III-47




III-48




III-49




III-50




III-51




III-52




III-53




III-54




III-55




III-56




III-57




III-58




III-59




III-60




III-61




III-62




III-63




III-64




III-65




III-66




III-67




III-68




III-69




III-70




III-71




III-72




III-73




III-74




III-75




IIII-76




IIII-77




IIII-78




IIII-79




IIII-80




IIII-81




IIII-82




IIII-83




IIII-84




IIII-85




IIII-86




IIII-87




III-88




III-89




III-90




III-91




III-92




III-93




III-94




III-95




III-96




III-97



III-98




III-99




III-100




III-101




III-102




III-103




III-104




III-105




III-106




III-107




III-108




III-109




III-110




III-111



III-112




III-113




III-114







III-115




III-116




III-117




III-118




III-119




III-120




III-121




III-122




III-123




III-124




III-125




III-126




III-127




III-128




III-129




III-130




III-131




III-132




III-133




III-134




III-135




III-136




III-137




III-138



III-139




III-140




III-141

5


10




III-142




III-143



III-144



III-145



III-146

5

In another embodiment, this invention provides a composition comprising a compound of formula III and a pharmaceutically acceptable carrier.

10

One aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula III.

15

Another aspect relates to a method of treating a disease that is alleviated by treatment with a GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition

20

comprising a compound of formula III.

Another aspect relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula

25

III. This method is especially useful for diabetic patients.

Another aspect relates to a method of inhibiting the production of hyperphosphorylated Tau protein in a patient in need thereof, comprising  
5 administering to said patient a therapeutically effective amount of a composition comprising a compound of formula III. This method is especially useful in halting or slowing the progression of Alzheimer's disease.

10 Another aspect relates to a method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula III. This method is  
15 especially useful for treating schizophrenia.

One aspect of this invention relates to a method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound  
20 of formula III.

Another aspect relates to a method of treating a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of administering to a patient in need of such a treatment a  
25 therapeutically effective amount of a composition comprising a compound of formula III. This method is especially useful for treating cancer, such as colon, ovarian, and breast cancer.

One aspect of this invention relates to a  
30 method of inhibiting CDK-2 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula III.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a CDK-2 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula III. This method is especially useful for treating cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis.

One aspect of this invention relates to a method of inhibiting Src activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula III.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a Src inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula III. This method is especially useful for treating hypercalcemia, osteoporosis, osteoarthritis, cancer, symptomatic treatment of bone metastasis, and Paget's disease.

Another method relates to inhibiting GSK-3, Aurora, CDK-2, or Src activity in a biological sample, which method comprises contacting the biological sample with the GSK-3, Aurora, CDK-2, or Src inhibitor of formula III, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3, Aurora, CDK-2, or Src.

Each of the aforementioned methods directed to the inhibition of GSK-3, Aurora, CDK-2, or Src, or the


treatment of a disease alleviated thereby, is preferably carried out with a preferred compound of formula III, as described above.

Compounds of formula III, wherein  $R^{2'}$  is  
5 hydrogen and  $R^x$  and  $R^y$  are taken together with the pyrimidine ring to form an optionally substituted quinazoline ring system, are also inhibitors of ERK-2 and AKT protein kinases.

Accordingly, another method of this invention  
10 relates to a method of inhibiting ERK-2 or AKT activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula III, wherein  $R^{2'}$  is  
15 hydrogen and  $R^x$  and  $R^y$  are taken together with the pyrimidine ring to form an optionally substituted quinazoline ring system.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a ERK-2 or AKT inhibitor, said method comprising the step of  
20 administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula III, wherein  $R^{2'}$  is hydrogen and  $R^x$  and  $R^y$  are taken together with the pyrimidine ring to form an optionally substituted  
25 quinazoline ring system. This method is especially useful for treating cancer, stroke, hepatomegaly, cardiovascular disease, Alzheimer's disease, cystic fibrosis, viral disease, autoimmune diseases, restenosis, psoriasis, allergic disorders including asthma,  
30 inflammation, and neurological disorders.

Another embodiment of this invention relates to compounds of formula IV:



IV

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or -R<sup>5</sup>, and at any substitutable ring nitrogen by -R<sup>4</sup>, provided that when Ring D is a six-membered aryl or heteroaryl ring, -R<sup>5</sup> is hydrogen at each ortho carbon position of Ring D;

R<sup>x</sup> and R<sup>y</sup> are independently selected from T-R<sup>3</sup>, or R<sup>x</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 1-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring is optionally and independently substituted by T-R<sup>3</sup>, and any substitutable nitrogen on said ring is substituted by R<sup>4</sup>;

T is a valence bond or a C<sub>1-4</sub> alkylidene chain;

R<sup>2</sup> and R<sup>2'</sup> are independently selected from -R, -T-W-R<sup>6</sup>, or R<sup>2</sup> and R<sup>2'</sup> are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring containing 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur,

wherein said fused ring is optionally substituted by up to three groups independently selected from halo, oxo, -CN, -NO<sub>2</sub>, -R<sup>7</sup>, or -V-R<sup>6</sup>;

R<sup>3</sup> is selected from -R, -halo, =O, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -COCH<sub>2</sub>COR, -NO<sub>2</sub>, -CN, -S(O)R, -S(O)<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>;

each R is independently selected from hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each R<sup>4</sup> is independently selected from -R<sup>7</sup>, -COR<sup>7</sup>,  
 5 -CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -CON(R<sup>7</sup>)<sub>2</sub>, or -SO<sub>2</sub>R<sup>7</sup>, or two R<sup>4</sup> on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or heteroaryl ring;

each R<sup>5</sup> is independently selected from -R, halo, -OR,  
 10 -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>;

15 V is -O-, -S-, -SO-, -SO<sub>2</sub>-, -N(R<sup>6</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -N(R<sup>6</sup>)CO-, -N(R<sup>6</sup>)C(O)O-, -N(R<sup>6</sup>)CON(R<sup>6</sup>)-, -N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(O)N(R<sup>6</sup>)-, -OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-,  
 20 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-, -C(R<sup>6</sup>)=N-O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, or -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-;



W is  $-C(R^6)_2O-$ ,  $-C(R^6)_2S-$ ,  $-C(R^6)_2SO-$ ,  $-C(R^6)_2SO_2-$ ,  
 $-C(R^6)_2SO_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  
 $-C(R^6)OC(O)-$ ,  $-C(R^6)OC(O)N(R^6)-$ ,  $-C(R^6)_2N(R^6)CO-$ ,  
 $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  $-C(R^6)=N-O-$ ,  
5  $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ ,  
 $-C(R^6)_2N(R^6)CON(R^6)-$ , or  $-CON(R^6)-$ ;

each  $R^6$  is independently selected from hydrogen or an  
optionally substituted  $C_{1-4}$  aliphatic group, or two  $R^6$   
groups on the same nitrogen atom are taken together  
10 with the nitrogen atom to form a 5-6 membered  
heterocyclyl or heteroaryl ring; and


each  $R^7$  is independently selected from hydrogen or an  
optionally substituted  $C_{1-6}$  aliphatic group, or two  $R^7$   
on the same nitrogen are taken together with the  
15 nitrogen to form a 5-8 membered heterocyclyl ring or  
heteroaryl.

Preferred formula IV Ring D monocyclic rings  
include substituted and unsubstituted phenyl, pyridinyl,  
piperidinyl, piperazinyl, pyrrolidinyl, thienyl,  
20 azepanyl, and morpholinyl rings. Preferred formula IV  
Ring D bicyclic rings include 1,2,3,4-  
tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl,  
2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl,  
isoquinolinyl, quinolinyl, and naphthyl. Examples of  
25 more preferred Ring D bicyclic rings include naphthyl and  
isoquinolinyl.


Preferred substituents on Ring D of formula IV  
include halo, oxo, CN,  $-NO_2$ ,  $-N(R^4)_2$ ,  $-CO_2R$ ,  $-CONH(R^4)$ ,  
 $-N(R^4)COR$ ,  $-SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ ,  $-SR$ ,  $-OR$ ,  $-C(O)R$ , or  
30 substituted or unsubstituted group selected from 5-6  
membered heterocyclyl,  $C_{6-10}$  aryl, or  $C_{1-6}$  aliphatic. More  
preferred  $R^5$  substituents include -halo, -CN, -oxo, -SR,  
-OR,  $-N(R^4)_2$ ,  $-C(O)R$ , or a substituted or unsubstituted  
group selected from 5-6 membered heterocyclyl,  $C_{6-10}$  aryl,

or C<sub>1-6</sub> aliphatic. Examples of Ring D substituents include -OH, phenyl, methyl, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH, pyrrolidinyl, OPh, CF<sub>3</sub>, C≡CH, Cl, Br, F, I, NH<sub>2</sub>, C(O)CH<sub>3</sub>, *i*-propyl, *tert*-butyl, SEt, OMe, N(Me)<sub>2</sub>, methylene dioxy, and ethylene dioxy.


When the R<sup>x</sup> and R<sup>y</sup> groups of formula IV are taken together to form a fused ring, preferred R<sup>x</sup>/R<sup>y</sup> rings include a 5-, 6-, 7-, or 8-membered unsaturated or partially unsaturated ring having 1-2 heteroatoms. This provides a bicyclic ring system containing the pyrimidine ring. Examples of preferred pyrimidine ring systems of formula IV are the mono- and bicyclic systems shown below.




IV-D




IV-E




IV-G




IV-H




IV-J




IV-K




IV-L




IV-M




IV-N




IV-O




IV-P




IV-Q




IV-R




IV-S




IV-T




IV-U




IV-V




IV-W




IV-X




IV-Y




IV-Z




IV-AA



IV-BB



IV-CC



IV-DD


More preferred pyrimidine ring systems of  
 5 formula IV include IV-E, IV-G, IV-H, IV-J, IV-K, IV-L,  
 IV-M, IV-T, and IV-U.

In the monocyclic pyrimidine ring system of  
 formula IV, preferred  $R^x$  groups include hydrogen, amino,  
 nitro, alkyl- or dialkylamino, acetamido, or a  $C_{1-4}$   
 10 aliphatic group such as methyl, ethyl, cyclopropyl,  
 isopropyl or t-butyl. Preferred  $R^y$  groups include T- $R^3$   
 wherein T is a valence bond or a methylene, and  $R^3$  is -R,  
 - $N(R^4)_2$ , or -OR. When  $R^3$  is -R or -OR, a preferred R is  
 an optionally substituted group selected from  $C_{1-6}$   
 15 aliphatic, phenyl, or a 5-6 membered heteroaryl or  
 heterocyclyl ring. Examples of preferred  $R^y$  groups  
 include 2-pyridyl, 4-pyridyl, piperidinyl, methyl, ethyl,  
 cyclopropyl, isopropyl, t-butyl, alkyl- or dialkylamino,  
 acetamido, optionally substituted phenyl such as phenyl,  
 20 methoxyphenyl, trimethoxyphenyl, or halo-substituted  
 phenyl, and methoxymethyl.

In the bicyclic pyrimidine ring system of  
 formula IV, the ring formed when  $R^x$  and  $R^y$  are taken  
 together may be substituted or unsubstituted. Suitable  
 25 substituents include -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R,  
 -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, - $N(R^4)_2$ , -CON( $R^4$ )<sub>2</sub>,  
 -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, -OC(=O)R, - $N(R^4)$ COR, - $N(R^4)$ CO<sub>2</sub> (optionally  
 substituted  $C_{1-6}$  aliphatic), - $N(R^4)$ N( $R^4$ )<sub>2</sub>, -C=NN( $R^4$ )<sub>2</sub>,  
 -C=N-OR, - $N(R^4)$ CON( $R^4$ )<sub>2</sub>, - $N(R^4)$ SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, - $N(R^4)$ SO<sub>2</sub>R, or  
 30 -OC(=O)N( $R^4$ )<sub>2</sub>, wherein R and  $R^4$  are as defined above for  
 compounds of formula IV. Preferred  $R^x/R^y$  ring

substituents include -halo, -R, -OR, -COR, -CO<sub>2</sub>R, -CON(R<sup>4</sup>)<sub>2</sub>, -CN, or -N(R<sup>4</sup>)<sub>2</sub> wherein R is a substituted or unsubstituted C<sub>1-6</sub> aliphatic group.

The R<sup>2</sup> and R<sup>2'</sup> groups of formula IV may be taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring. Preferred fused rings include benzo, pyrido, pyrimido, and a partially unsaturated 6-membered carbocyclo ring. These are exemplified in the following formula IV compounds having a pyrazole-containing bicyclic ring system:



Preferred substituents on the R<sup>2</sup>/R<sup>2'</sup> fused ring of formula IV include one or more of the following:  
 -halo, -N(R<sup>4</sup>)<sub>2</sub>, -C<sub>1-4</sub> alkyl, -C<sub>1-4</sub> haloalkyl, -NO<sub>2</sub>, -O(C<sub>1-4</sub> alkyl), -CO<sub>2</sub>(C<sub>1-4</sub> alkyl), -CN, -SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -OC(O)NH<sub>2</sub>, -NH<sub>2</sub>SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -NHC(O)(C<sub>1-4</sub> alkyl), -C(O)NH<sub>2</sub>, and -CO(C<sub>1-4</sub> alkyl), wherein the (C<sub>1-4</sub> alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the (C<sub>1-4</sub> alkyl) group is methyl.

When the pyrazole ring system of formula IV is monocyclic, preferred R<sup>2</sup> groups include hydrogen, a substituted or unsubstituted group selected from aryl, heteroaryl, or a C<sub>1-6</sub> aliphatic group. Examples of such preferred R<sup>2</sup> groups include methyl, t-butyl, -CH<sub>2</sub>OCH<sub>3</sub>, cyclopropyl, furanyl, thienyl, and phenyl. A preferred R<sup>2'</sup> group is hydrogen.

Preferred formula IV compounds have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring D is an optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^x$  is hydrogen or  $C_{1-4}$  aliphatic and  $R^y$  is T- $R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted 5-7 membered unsaturated or partially unsaturated ring having 1-2 ring heteroatoms; and

(c)  $R^{2'}$  is hydrogen or methyl and  $R^2$  is T-W- $R^6$  or R, wherein W is  $-C(R^6)_2O-$ ,  $-C(R^6)_2N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  $-C(R^6)OC(O)-$ ,  $-C(R^6)_2N(R^6)CO-$ ,  $-C(R^6)_2N(R^6)C(O)O-$ , or  $-CON(R^6)-$ , and R is an optionally substituted group selected from  $C_{1-6}$  aliphatic or phenyl, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido, or partially unsaturated 6-membered carbocyclo ring.

More preferred compounds of formula IV have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^x$  is hydrogen or methyl and  $R^y$  is -R,  $N(R^4)_2$ , or -OR, or  $R^x$  and  $R^y$  are taken together with their

intervening atoms to form a 5-7 membered unsaturated or partially unsaturated ring having 1-2 ring nitrogens, wherein said ring is optionally substituted with -R, halo, oxo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>; and

(c) each R<sup>5</sup> is independently selected from halo, oxo, CN, NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -SR, -OR, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic.

Even more preferred compounds of formula IV have one or more, and more preferably all, of the features selected from the group consisting of:

(a) R<sup>x</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a 6-membered unsaturated or partially unsaturated ring having 1-2 ring nitrogens, optionally substituted with halo, CN, oxo, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, (C<sub>1-6</sub> alkyl)carbonyl, (C<sub>1-6</sub> alkyl)sulfonyl, mono- or dialkylamino, mono- or dialkylaminocarbonyl, mono- or dialkylaminocarbonyloxy, or 5-6 membered heteroaryl;


(b) each R<sup>5</sup> is independently selected from -halo, -CN, -oxo, -SR, -OR, -N(R<sup>4</sup>)<sub>2</sub>, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic; and

(c) R<sup>2'</sup> is hydrogen and R<sup>2</sup> is T-W-R<sup>6</sup> or R, wherein W is -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -C(R<sup>6</sup>)OC(O)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CO-, or -CON(R<sup>6</sup>)-, and R is an optionally substituted group selected from C<sub>1-6</sub> aliphatic or phenyl, or R<sup>2</sup> and R<sup>2'</sup> are taken together with their intervening atoms to form a benzo, pyrido, or partially unsaturated 6-membered carbocyclo ring optionally


substituted with -halo, oxo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl),  $-CN$ ,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , or  $-CO(C_{1-4}$  alkyl), wherein  
 5 the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group.

Representative compounds of formula IV are set forth in Table 3 below.


10 Table 3.




IV-1




IV-2




IV-3




IV-4




IV-5




IV-6



IV-7




IV-8




IV-9







IV-10




IV-11




IV-12



IV-13




IV-14




IV-15


5




IV-16




IV-17




IV-18



IV-19




IV-20




IV-21


10




IV-22




IV-23




IV-24




IV-25




IV-26




IV-27




IV-28




IV-29




IV-30



IV-31



IV-32



IV-33

In another embodiment, this invention provides a composition comprising a compound of formula IV and a pharmaceutically acceptable carrier.

One aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, comprising administering to the patient a therapeutically

effective amount of a composition comprising a compound of formula IV.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula IV.

Another aspect relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula IV. This method is especially useful for diabetic patients.

Another aspect relates to a method of inhibiting the production of hyperphosphorylated Tau protein in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula IV. This method is especially useful in halting or slowing the progression of Alzheimer's disease.

Another aspect relates to a method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula IV. This method is especially useful for treating schizophrenia.

One aspect of this invention relates to a method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula IV.

Another aspect relates to a method of treating a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula IV. This method is especially useful for treating cancer, such as colon, ovarian, and breast cancer.

One aspect of this invention relates to a method of inhibiting CDK-2 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula IV.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a CDK-2 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula IV. This method is especially useful for treating cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis.


Another method relates to inhibiting GSK-3, Aurora, or CDK-2 activity in a biological sample, which method comprises contacting the biological sample with the GSK-3 or Aurora inhibitor of formula IV, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3, Aurora or CDK-2.

Each of the aforementioned methods directed to the inhibition of GSK-3, Aurora or CDK-2, or the treatment of a disease alleviated thereby, is preferably

carried out with a preferred compound of formula IV, as described above.

Another embodiment of this invention relates to compounds of formula V:

5



V

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

$Z^1$  is N,  $CR^a$ , or CH and  $Z^2$  is N or CH, provided that one of  $Z^1$  and  $Z^2$  is nitrogen;

G is Ring C or Ring D;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from  $-R^1$ , any substitutable non-ortho carbon position on Ring C is independently substituted by  $-R^5$ , and two adjacent substituents on Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo, oxo, or  $-R^8$ ;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected

from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or -R<sup>5</sup>, and at any substitutable ring nitrogen by -R<sup>4</sup>, provided that when Ring D is a six-membered aryl or heteroaryl ring, -R<sup>5</sup> is hydrogen at each ortho carbon position of Ring D;

R<sup>1</sup> is selected from -halo, -CN, -NO<sub>2</sub>, T-V-R<sup>6</sup>, phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or C<sub>1-6</sub> aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or -R<sup>8</sup>, said C<sub>1-6</sub> aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or R<sup>1</sup> and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

R<sup>x</sup> and R<sup>y</sup> are independently selected from T-R<sup>3</sup>, or R<sup>x</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 0-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring formed by R<sup>x</sup> and R<sup>y</sup> is substituted by oxo or T-R<sup>3</sup>, and any substitutable nitrogen on said ring formed by R<sup>x</sup> and R<sup>y</sup> is substituted by R<sup>4</sup>;

T is a valence bond or a C<sub>1-4</sub> alkylidene chain;

R<sup>2</sup> and R<sup>2'</sup> are independently selected from -R, -T-W-R<sup>6</sup>, or R<sup>2</sup> and R<sup>2'</sup> are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable carbon on said fused ring formed by R<sup>2</sup> and R<sup>2'</sup> is substituted by halo, oxo, -CN, -NO<sub>2</sub>, -R<sup>7</sup>, or -V-R<sup>6</sup>, and any substitutable nitrogen on said ring formed by R<sup>2</sup> and R<sup>2'</sup> is substituted by R<sup>4</sup>;

$R^3$  is selected from -R, -halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -COCH<sub>2</sub>COR, -NO<sub>2</sub>, -CN, -S(O)R, -S(O)<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>7</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>7</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>7</sup>)COR, -N(R<sup>7</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>7</sup>)CON(R<sup>7</sup>)<sub>2</sub>, -N(R<sup>7</sup>)SO<sub>2</sub>N(R<sup>7</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>7</sup>)<sub>2</sub>;

each R is independently selected from hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each R<sup>4</sup> is independently selected from -R<sup>7</sup>, -COR<sup>7</sup>, -CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -CON(R<sup>7</sup>)<sub>2</sub>, or -SO<sub>2</sub>R<sup>7</sup>, or two R<sup>4</sup> on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or heteroaryl ring;

each R<sup>5</sup> is independently selected from -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>, or R<sup>5</sup> and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

V is -O-, -S-, -SO-, -SO<sub>2</sub>-, -N(R<sup>6</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -N(R<sup>6</sup>)CO-, -N(R<sup>6</sup>)C(O)O-, -N(R<sup>6</sup>)CON(R<sup>6</sup>)-, -N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(O)N(R<sup>6</sup>)-, -OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-, -C(R<sup>6</sup>)=N-O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, or -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-;

W is -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-,


-C(R<sup>6</sup>)OC(O)-, -C(R<sup>6</sup>)OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CO-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-, -C(R<sup>6</sup>)=N-O-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-, or -CON(R<sup>6</sup>)-;

- 5 each R<sup>6</sup> is independently selected from hydrogen, an optionally substituted C<sub>1-4</sub> aliphatic group, or two R<sup>6</sup> groups on the same nitrogen atom are taken together with the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;
- 10 each R<sup>7</sup> is independently selected from hydrogen or an optionally substituted C<sub>1-6</sub> aliphatic group, or two R<sup>7</sup> on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring;
- 15 each R<sup>8</sup> is independently selected from an optionally substituted C<sub>1-4</sub> aliphatic group, -OR<sup>6</sup>, -SR<sup>6</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>R<sup>6</sup>, -N(R<sup>6</sup>)<sub>2</sub>, -N(R<sup>6</sup>)N(R<sup>6</sup>)<sub>2</sub>, -CN, -NO<sub>2</sub>, -CON(R<sup>6</sup>)<sub>2</sub>, or -CO<sub>2</sub>R<sup>6</sup>; and
- R<sup>a</sup> is selected from halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -OC(=O)N(R<sup>4</sup>)<sub>2</sub>, or an optionally substituted group
- 20 selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms.
- 25


Compounds of formula V may be represented by specifying Z<sup>1</sup> and Z<sup>2</sup> as shown below:

30






Va




Vb




Vc


When the  $R^x$  and  $R^y$  groups of formula V are taken  
 5 together to form a fused ring, preferred  $R^x/R^y$  rings  
 include a 5-, 6-, 7-, or 8-membered unsaturated or  
 partially unsaturated ring having 0-2 heteroatoms,  
 wherein said  $R^x/R^y$  ring is optionally substituted. This  
 provides a bicyclic ring system containing a pyridine  
 10 ring. Examples of preferred bicyclic ring systems of  
 formula V are shown below.




Va-A




Vb-A




Vc-A




Va-B




Vb-B




Vc-B




Va-C




Vb-C




Vc-C




Va-D




Vb-D




Vc-D




Va-E




Vb-E




Vc-E




Va-F




Vb-F




Vc-F




Va-J




Vb-J




Vc-J




Va-K




Vb-K




Vc-K



Va-L




Vb-L




Vc-L


5




Va-M




Vb-M




Vc-M



Va-N




Vb-N




Vc-N


10




Va-O




Vb-O




Vc-O



Va-P



Vb-P



Vc-P


More preferred bicyclic ring systems of formula  
 5 V include Va-A, Vb-A, Vc-A, Va-B, Vb-B, Vc-B, Va-D, Vb-D, Vc-D, Va-E, Vb-E, Vc-E, Va-J, Vb-J, Vc-J, Va-K, Vb-K, Vc-K, Va-L, Vb-L, Vc-L, Va-M, Vb-M, and Vc-M, most preferably Va-A, Vb-A, Vc-A, Va-B, Vb-B, and Vc-B.

In the monocyclic pyridine ring system of  
 10 formula V, preferred  $R^x$  groups include hydrogen, alkyl- or dialkylamino, acetamido, or a  $C_{1-4}$  aliphatic group such as methyl, ethyl, cyclopropyl, isopropyl or t-butyl. Preferred  $R^y$  groups include  $T-R^3$  wherein T is a valence bond or a methylene, and  $R^3$  is  $-R$ ,  $-N(R^4)_2$ , or  $-OR$ . When  
 15  $R^3$  is  $-R$  or  $-OR$ , a preferred R is an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring. Examples of preferred  $R^y$  include 2-pyridyl, 4-pyridyl, piperidinyl, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, alkyl- or  
 20 dialkylamino, acetamido, optionally substituted phenyl such as phenyl or halo-substituted phenyl, and methoxymethyl.

In the bicyclic ring system of formula V, the ring formed when  $R^x$  and  $R^y$  are taken together may be  
 25 substituted or unsubstituted. Suitable substituents include  $-R$ , halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  
 30  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ , wherein R and  $R^4$  are as defined above.

Preferred  $R^x/R^y$  ring substituents include -halo, -R, -OR, -COR, -CO<sub>2</sub>R, -CON(R<sup>4</sup>)<sub>2</sub>, -CN, or -N(R<sup>4</sup>)<sub>2</sub> wherein R is an optionally substituted C<sub>1-6</sub> aliphatic group.

The R<sup>2</sup> and R<sup>2'</sup> groups of formula V may be taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring. Preferred fused rings include benzo, pyrido, pyrimido, and a partially unsaturated 6-membered carbocyclo ring. These are exemplified in the following formula V compounds having a pyrazole-containing bicyclic ring system:




Preferred substituents on the R<sup>2</sup>/R<sup>2'</sup> fused ring of formula V include one or more of the following: -halo, -N(R<sup>4</sup>)<sub>2</sub>, -C<sub>1-4</sub> alkyl, -C<sub>1-4</sub> haloalkyl, -NO<sub>2</sub>, -O(C<sub>1-4</sub> alkyl), -CO<sub>2</sub>(C<sub>1-4</sub> alkyl), -CN, -SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -OC(O)NH<sub>2</sub>, -NH<sub>2</sub>SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -NHC(O)(C<sub>1-4</sub> alkyl), -C(O)NH<sub>2</sub>, and -CO(C<sub>1-4</sub> alkyl), wherein the (C<sub>1-4</sub> alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the (C<sub>1-4</sub> alkyl) group is methyl.


When the pyrazole ring system is monocyclic, preferred R<sup>2</sup> groups include hydrogen, C<sub>1-4</sub> aliphatic, alkoxy carbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred R<sup>2</sup> substituents include methyl, cyclopropyl, ethyl,

isopropyl, propyl, t-butyl, cyclopentyl, phenyl, CO<sub>2</sub>H,  
 CO<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>OH, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>,  
 CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>Ph, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHCOOC(CH<sub>3</sub>)<sub>3</sub>,  
 CONHCH(CH<sub>3</sub>)<sub>2</sub>, CONHCH<sub>2</sub>CH=CH<sub>2</sub>, CONHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CONHCH<sub>2</sub>Ph,  
 5 CONH(cyclohexyl), CON(Et)<sub>2</sub>, CON(CH<sub>3</sub>)CH<sub>2</sub>Ph, CONH(n-C<sub>3</sub>H<sub>7</sub>),  
 CON(Et)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, CONHCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, CON(n-C<sub>3</sub>H<sub>7</sub>)<sub>2</sub>, CO(3-  
 methoxymethylpyrrolidin-1-yl), CONH(3-tolyl), CONH(4-  
 tolyl), CONHCH<sub>3</sub>, CO(morpholin-1-yl), CO(4-methylpiperazin-  
 1-yl), CONHCH<sub>2</sub>CH<sub>2</sub>OH, CONH<sub>2</sub>, and CO(piperidin-1-yl). A  
 10 preferred R<sup>2'</sup> group is hydrogen.


More preferred ring systems of formula V are  
 the following, which may be substituted as described  
 above, wherein R<sup>2</sup> and R<sup>2'</sup> are taken together with the  
 pyrazole ring to form an optionally substituted indazole  
 15 ring; and R<sup>x</sup> and R<sup>y</sup> are each methyl, or R<sup>x</sup> and R<sup>y</sup> are taken  
 together with the pyridine ring to form an optionally  
 substituted quinoline, isoquinoline, tetrahydroquinoline  
 or tetrahydroisoquinoline ring:



V-Aa



V-Ba



V-Ha

20

When G is Ring C, preferred formula V Ring C  
 groups are phenyl and pyridinyl. When two adjacent  
 substituents on Ring C are taken together to form a fused  
 25 ring, Ring C is contained in a bicyclic ring system.  
 Preferred fused rings include a benzo or pyrido ring.  
 Such rings preferably are fused at ortho and meta  
 positions of Ring C. Examples of preferred bicyclic Ring  
 C systems include naphthyl and isoquinolinyl. Preferred

R<sup>1</sup> groups include -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>.

When R<sup>1</sup> is an optionally substituted C<sub>1-6</sub> aliphatic group,

5 the most preferred optional substituents are halogen.

Examples of preferred R<sup>1</sup> groups include -CF<sub>3</sub>, -Cl, -F, -CN, -COCH<sub>3</sub>, -OCH<sub>3</sub>, -OH, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -CH<sub>3</sub>, -CF<sub>2</sub>CH<sub>3</sub>, cyclohexyl, t-butyl, isopropyl, cyclopropyl, -C≡CH, -C≡C-CH<sub>3</sub>, -SO<sub>2</sub>CH<sub>3</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, -CO<sub>2</sub>CH<sub>3</sub>, -CONH<sub>2</sub>,

10 -NHCOCH<sub>3</sub>, -OC(O)NH<sub>2</sub>, -NHSO<sub>2</sub>CH<sub>3</sub>, and -OCF<sub>3</sub>.

On Ring C preferred R<sup>5</sup> substituents, when present, include -halo, -CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, optionally substituted C<sub>1-6</sub> aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, and -N(R<sup>4</sup>)SO<sub>2</sub>R. More preferred R<sup>5</sup> substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic).

15 Examples of such preferred R<sup>5</sup> substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHMe, -NMe<sub>2</sub>, -OEt, methyl, ethyl, 20 cyclopropyl, isopropyl, t-butyl, and -CO<sub>2</sub>Et.

When G is Ring D, preferred formula V Ring D monocyclic rings include substituted and unsubstituted phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, and morpholinyl rings.

25 When two adjacent substituents on Ring D are taken together to form a fused ring, the Ring D system is bicyclic. Preferred formula V Ring D bicyclic rings include 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, and 30 naphthyl. Examples of more preferred bicyclic Ring D systems include naphthyl and isoquinolinyl.

Preferred substituents on Ring D of formula V include one or more of the following: halo, oxo, CN, -NO<sub>2</sub>,

-N(R<sup>4</sup>)<sub>2</sub>, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -SR, -OR, -C(O)R, or substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. More preferred Ring D substituents include

5 -halo, -CN, -oxo, -SR, -OR, -N(R<sup>4</sup>)<sub>2</sub>, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. Examples of Ring D substituents include -OH, phenyl, methyl, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH, pyrrolidinyl, OPh, CF<sub>3</sub>, C≡CH, Cl,

10 Br, F, I, NH<sub>2</sub>, C(O)CH<sub>3</sub>, *i*-propyl, *tert*-butyl, Set, OMe, N(Me)<sub>2</sub>, methylene dioxy, and ethylene dioxy.

Preferred formula V compounds have one or more, and more preferably all, of the features selected from the group consisting of:

15 (a) Ring C is a phenyl or pyridinyl ring, optionally substituted by -R<sup>5</sup>, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a naphthyl, quinolinyl or isoquinolinyl ring, and R<sup>1</sup> is

20 -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>; or Ring D is an optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl,

25 thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-indolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b) R<sup>x</sup> is hydrogen or C<sub>1-4</sub> aliphatic and R<sup>y</sup> is T-  
30 R<sup>3</sup>, or R<sup>x</sup> and R<sup>y</sup> are taken together with their intervening atoms to form an optionally substituted 5-7 membered unsaturated or partially unsaturated ring having 0-2 ring nitrogens; and



(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring.

More preferred compounds of formula V have one or more, and more preferably all, of the features selected from the group consisting of:

10 (a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^x$  is hydrogen or methyl and  $R^y$  is -R,  $N(R^4)_2$ , or -OR, or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a benzo ring or a 5-7 membered partially unsaturated carbocyclo ring, said benzo or carbocyclo ring optionally substituted with -R, halo, -OR,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ , -CN,  $-S(O)R$ ,  $-SO_2R$ , -SR,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together

with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring; and

- (d) Ring D is substituted by oxo or  $R^5$ , wherein  
5 each  $R^5$  is independently selected from -halo, -CN, -NO<sub>2</sub>, -N( $R^4$ )<sub>2</sub>, optionally substituted C<sub>1-6</sub> aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH( $R^4$ ), -N( $R^4$ )COR, -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, or -N( $R^4$ )SO<sub>2</sub>R.

Even more preferred compounds of formula V have  
10 one or more, and more preferably all, of the features selected from the group consisting of:

- (a) Ring C is a phenyl or pyridinyl ring, optionally substituted by - $R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring  
15 system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a C<sub>1-4</sub> aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl,  
20 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or naphthyl;


- (b)  $R^x$  is hydrogen or methyl and  $R^y$  is methyl, methoxymethyl, ethyl, cyclopropyl, isopropyl, t-butyl,  
25 alkyl- or an optionally substituted group selected from 2-pyridyl, 4-pyridyl, piperidinyl, or phenyl, or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a benzo ring or a 6-membered partially unsaturated carbocyclo ring optionally substituted with halo, CN,  
30 oxo, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, (C<sub>1-6</sub> alkyl)carbonyl, (C<sub>1-6</sub> alkyl)sulfonyl, mono- or dialkylamino, mono- or dialkylaminocarbonyl, mono- or dialkylaminocarbonyloxy, or 5-6 membered heteroaryl;

(c)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring optionally substituted with -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl),  $-CN$ ,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , or  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group; and


(d) Ring D is substituted by oxo or  $R^5$ , wherein each  $R^5$  is independently selected from  $-Cl$ ,  $-F$ ,  $-CN$ ,  $-CF_3$ ,  $-NH_2$ ,  $-NH(C_{1-4}$  aliphatic),  $-N(C_{1-4}$  aliphatic) $_2$ ,  $-O(C_{1-4}$  aliphatic),  $C_{1-4}$  aliphatic, and  $-CO_2(C_{1-4}$  aliphatic).

Representative compounds of formula V are set forth in Table 4 below.


Table 4.




V-1




V-2




V-3




V-4




V-5




V-6




V-7




V-8




V-9




V-10




V-11




V-12




V-13




V-14




V-15




V-16




V-17




V-18




V-19




V-20




V-21




V-22




V-23




V-24




V-25




V-26




V-27




V-28




V-29




V-30




V-31




V-32




V-33



V-34




V-35




V-36


5




V-37




V-38




V-39



V-40




V-41




V-42


10




V-43




V-44




V-45



V-46




V-47




V-48


5




V-49




V-50




V-51



V-52




V-53




V-54


10




V-55




V-56




V-57




V-58




V-59




V-60




V-61




V-62




V-63



V-64




V-65




V-66





V-67



V-68

In another embodiment, this invention provides  
5 a composition comprising a compound of formula V and a  
pharmaceutically acceptable carrier.

One aspect of this invention relates to a  
method of inhibiting GSK-3 activity in a patient,  
comprising administering to the patient a therapeutically  
10 effective amount of a composition comprising a compound  
of formula V.

Another aspect relates to a method of treating  
a disease that is alleviated by treatment with a GSK-3  
inhibitor, said method comprising the step of  
15 administering to a patient in need of such a treatment a  
therapeutically effective amount of a composition  
comprising a compound of formula V.

Another aspect relates to a method of enhancing  
glycogen synthesis and/or lowering blood levels of  
20 glucose in a patient in need thereof, comprising  
administering to said patient a therapeutically effective  
amount of a composition comprising a compound of formula  
V. This method is especially useful for diabetic  
patients.

25 Another aspect relates to a method of  
inhibiting the production of hyperphosphorylated Tau  
protein in a patient in need thereof, comprising  
administering to said patient a therapeutically effective  
amount of a composition comprising a compound of formula

V. This method is especially useful in halting or slowing the progression of Alzheimer's disease.

Another aspect relates to a method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient  
5 in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula V. This method is especially useful for treating schizophrenia.

One aspect of this invention relates to a  
10 method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula V.

Another aspect relates to a method of treating  
15 a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula V. This method is  
20 especially useful for treating cancer, such as colon, ovarian, and breast cancer.

One aspect of this invention relates to a method of inhibiting CDK-2 activity in a patient, comprising administering to the patient a therapeutically  
25 effective amount of a composition comprising a compound of formula V.


Another aspect relates to a method of treating a disease that is alleviated by treatment with a CDK-2 inhibitor, said method comprising the step of  
30 administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula V. This method is especially useful for treating cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis,

cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis.

Another method relates to inhibiting GSK-3, Aurora, or CDK-2 activity in a biological sample, which method comprises contacting the biological sample with the GSK-3 or Aurora inhibitor of formula V, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3, Aurora or CDK-2.

Each of the aforementioned methods directed to the inhibition of GSK-3, Aurora or CDK-2, or the treatment of a disease alleviated thereby, is preferably carried out with a preferred compound of formula V, as described above.

Another embodiment of this invention relates to compounds of formula VI:



VI

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

G is Ring C or Ring D;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from -R<sup>1</sup>, any substitutable non-ortho carbon position on Ring C is independently substituted by -R<sup>5</sup>, and two adjacent substituents on Ring C are optionally taken together with their

intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo, oxo, or  $-R^8$ ;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or  $-R^5$ , and at any substitutable ring nitrogen by  $-R^4$ , provided that when Ring D is a six-membered aryl or heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^1$  is selected from -halo, -CN,  $-NO_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $C_{1-6}$  aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or  $-R^8$ , said  $C_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

$R^Y$  is T- $R^{3'}$ ;

T is a valence bond or a  $C_{1-4}$  alkylidene chain;

$R^2$  and  $R^{2'}$  are independently selected from -R, -T-W- $R^6$ , or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable carbon on said fused ring formed by  $R^2$  and  $R^{2'}$  is substituted by halo, oxo, -CN,  $-NO_2$ ,  $-R^7$ , or

-V-R<sup>6</sup>, and any substitutable nitrogen on said ring formed by R<sup>2</sup> and R<sup>2'</sup> is substituted by R<sup>4</sup>;

R<sup>3'</sup> is an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>3-10</sub> carbocyclyl, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each R is independently selected from hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each R<sup>4</sup> is independently selected from -R<sup>7</sup>, -COR<sup>7</sup>, -CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -CON(R<sup>7</sup>)<sub>2</sub>,  
 5 or -SO<sub>2</sub>R<sup>7</sup>, or two R<sup>4</sup> on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or heteroaryl ring;

each R<sup>5</sup> is independently selected from -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR,  
 10 -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>, or R<sup>5</sup> and an adjacent substituent taken together with their  
 15 intervening atoms form said ring fused to Ring C;

V is -O-, -S-, -SO-, -SO<sub>2</sub>-, -N(R<sup>6</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>6</sup>)-,  
 -N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -N(R<sup>6</sup>)CO-, -N(R<sup>6</sup>)C(O)O-,  
 -N(R<sup>6</sup>)CON(R<sup>6</sup>)-, -N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)N(R<sup>6</sup>)-,  
 -C(O)N(R<sup>6</sup>)-, -OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-,  
 20 -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-,  
 -C(R<sup>6</sup>)=N-O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, or  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-;

W is -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-,  
 25 -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-,


$-C(R^6)OC(O)-$ ,  $-C(R^6)OC(O)N(R^6)-$ ,  $-C(R^6)_2N(R^6)CO-$ ,  
 $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  $-C(R^6)=N-O-$ ,  
 $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ ,  
 $-C(R^6)_2N(R^6)CON(R^6)-$ , or  $-CON(R^6)-$ ;

- 5 each  $R^6$  is independently selected from hydrogen, an optionally substituted  $C_{1-4}$  aliphatic group, or two  $R^6$  groups on the same nitrogen atom are taken together with the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;
- 10 each  $R^7$  is independently selected from hydrogen or an optionally substituted  $C_{1-6}$  aliphatic group, or two  $R^7$  on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring; and
- 15 each  $R^8$  is independently selected from an optionally substituted  $C_{1-4}$  aliphatic group,  $-OR^6$ ,  $-SR^6$ ,  $-COR^6$ ,  $-SO_2R^6$ ,  $-N(R^6)_2$ ,  $-N(R^6)N(R^6)_2$ ,  $-CN$ ,  $-NO_2$ ,  $-CON(R^6)_2$ , or  $-CO_2R^6$ .

- Preferred  $R^Y$  groups of formula VI include  $T-R^{3'}$
- 20 wherein T is a valence bond or a methylene, and  $R^{3'}$  is an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{3-10}$  carbocyclyl,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms. A preferred  $R^{3'}$  group is an optionally substituted
- 25 group selected from  $C_{3-6}$  carbocyclyl, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring. Examples of preferred  $R^Y$  include 2-pyridyl, 4-pyridyl, piperidinyl, morpholinyl, cyclopropyl, cyclohexyl, and optionally substituted phenyl such as phenyl or halo-substituted
- 30 phenyl.

The  $R^2$  and  $R^{2'}$  groups of formula VI may be taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring. Preferred fused rings include benzo, pyrido, pyrimido, and a partially

unsaturated 6-membered carbocyclo ring. These are exemplified in the following formula VI compounds having a pyrazole-containing bicyclic ring system:



Preferred substituents on the  $R^2/R^{2'}$  fused ring include one or more of the following: -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl),  $-CN$ ,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , and  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the  $(C_{1-4}$  alkyl) group is methyl.

When the pyrazole ring system is monocyclic, preferred  $R^2$  groups of formula VI include hydrogen,  $C_{1-4}$  aliphatic, alkoxycarbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred  $R^2$  substituents include methyl, cyclopropyl, ethyl, isopropyl, propyl, t-butyl, cyclopentyl, phenyl,  $CO_2H$ ,  $CO_2CH_3$ ,  $CH_2OH$ ,  $CH_2OCH_3$ ,  $CH_2CH_2CH_2OH$ ,  $CH_2CH_2CH_2OCH_3$ ,  $CH_2CH_2CH_2OCH_2Ph$ ,  $CH_2CH_2CH_2NH_2$ ,  $CH_2CH_2CH_2NHCOOC(CH_3)_3$ ,  $CONHCH(CH_3)_2$ ,  $CONHCH_2CH=CH_2$ ,  $CONHCH_2CH_2OCH_3$ ,  $CONHCH_2Ph$ ,  $CONH(cyclohexyl)$ ,  $CON(Et)_2$ ,  $CON(CH_3)CH_2Ph$ ,  $CONH(n-C_3H_7)$ ,  $CON(Et)CH_2CH_2CH_3$ ,  $CONHCH_2CH(CH_3)_2$ ,  $CON(n-C_3H_7)_2$ ,  $CO(3-methoxymethylpyrrolidin-1-yl)$ ,  $CONH(3-tolyl)$ ,  $CONH(4-$

tolyl), CONHCH<sub>3</sub>, CO(morpholin-1-yl), CO(4-methylpiperazin-1-yl), CONHCH<sub>2</sub>CH<sub>2</sub>OH, CONH<sub>2</sub>, and CO(piperidin-1-yl). A preferred R<sup>2'</sup> group is hydrogen.

When G is Ring C, preferred formula VI Ring C groups are phenyl and pyridinyl. When two adjacent substituents on Ring C are taken together to form a fused ring, Ring C is contained in a bicyclic ring system. Preferred fused rings include a benzo or pyrido ring. Such rings preferably are fused at ortho and meta positions of Ring C. Examples of preferred bicyclic Ring C systems include naphthyl and isoquinolinyl. Preferred R<sup>1</sup> groups include -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>. When R<sup>1</sup> is an optionally substituted C<sub>1-6</sub> aliphatic group, the most preferred optional substituents are halogen. Examples of preferred R<sup>1</sup> groups include -CF<sub>3</sub>, -Cl, -F, -CN, -COCH<sub>3</sub>, -OCH<sub>3</sub>, -OH, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -CH<sub>3</sub>, -CF<sub>2</sub>CH<sub>3</sub>, cyclohexyl, t-butyl, isopropyl, cyclopropyl, -C≡CH, -C≡C-CH<sub>3</sub>, -SO<sub>2</sub>CH<sub>3</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, -CO<sub>2</sub>CH<sub>3</sub>, -CONH<sub>2</sub>, -NHCOCH<sub>3</sub>, -OC(O)NH<sub>2</sub>, -NHSO<sub>2</sub>CH<sub>3</sub>, and -OCF<sub>3</sub>.

On Ring C preferred R<sup>5</sup> substituents, when present, include -halo, -CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, optionally substituted C<sub>1-6</sub> aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, and -N(R<sup>4</sup>)SO<sub>2</sub>R. More preferred R<sup>5</sup> substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic). Examples of such preferred R<sup>5</sup> substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHMe, -NMe<sub>2</sub>, -OEt, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, and -CO<sub>2</sub>Et.

When G is Ring D, preferred formula VI Ring D monocyclic rings include substituted and unsubstituted phenyl, pyridinyl, piperidinyl, piperazinyl,



pyrrolidinyl, thienyl, azepanyl, and morpholinyl rings. When two adjacent substituents on Ring D are taken together to form a fused ring, the Ring D system is bicyclic. Preferred formula VI Ring D bicyclic rings include 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, and naphthyl. Examples of more preferred bicyclic Ring D systems include naphthyl and isoquinolinyl.

Preferred substituents on formula VI Ring D include one or more of the following: halo, oxo, CN,  $-\text{NO}_2$ ,  $-\text{N}(\text{R}^4)_2$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{CONH}(\text{R}^4)$ ,  $-\text{N}(\text{R}^4)\text{COR}$ ,  $-\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ ,  $-\text{SR}$ ,  $-\text{OR}$ ,  $-\text{C}(\text{O})\text{R}$ , or substituted or unsubstituted group selected from 5-6 membered heterocyclyl,  $\text{C}_{6-10}$  aryl, or  $\text{C}_{1-6}$  aliphatic. More preferred Ring D substituents include -halo, -CN, -oxo, -SR, -OR,  $-\text{N}(\text{R}^4)_2$ ,  $-\text{C}(\text{O})\text{R}$ , or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl,  $\text{C}_{6-10}$  aryl, or  $\text{C}_{1-6}$  aliphatic. Examples of Ring D substituents include -OH, phenyl, methyl,  $\text{CH}_2\text{OH}$ ,  $\text{CH}_2\text{CH}_2\text{OH}$ , pyrrolidinyl, OPh,  $\text{CF}_3$ ,  $\text{C}\equiv\text{CH}$ , Cl, Br, F, I,  $\text{NH}_2$ ,  $\text{C}(\text{O})\text{CH}_3$ , *i*-propyl, *tert*-butyl, SEt, OMe,  $\text{N}(\text{Me})_2$ , methylene dioxy, and ethylene dioxy.

Preferred formula VI compounds have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is selected from a phenyl or pyridinyl ring, optionally substituted by  $-\text{R}^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a naphthyl, quinolinyl or isoquinolinyl ring, and  $\text{R}^1$  is -halo, an optionally substituted  $\text{C}_{1-6}$  aliphatic group, phenyl,  $-\text{COR}^6$ ,  $-\text{OR}^6$ , -CN,  $-\text{SO}_2\text{R}^6$ ,  $-\text{SO}_2\text{NH}_2$ ,  $-\text{N}(\text{R}^6)_2$ ,  $-\text{CO}_2\text{R}^6$ ,  $-\text{CONH}_2$ ,  $-\text{NHCOR}^6$ ,  $-\text{OC}(\text{O})\text{NH}_2$ , or  $-\text{NHSO}_2\text{R}^6$ ; or Ring D is an optionally substituted ring selected from a

phenyl, pyridinyl, piperidinyl, piperazinyl,  
pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-  
tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl,  
2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl,  
5 isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^y$  is  $T-R^{3'}$ , wherein T is a valence bond or  
a methylene; and

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a  
substituted or unsubstituted group selected from aryl,  
10 heteroaryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are  
taken together with their intervening atoms to form a  
substituted or unsubstituted benzo, pyrido, pyrimido or  
partially unsaturated 6-membered carbocyclo ring.

More preferred compounds of formula VI have one  
15 or more, and more preferably all, of the features  
selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring,  
optionally substituted by  $-R^5$ , wherein when Ring C and two  
adjacent substituents thereon form a bicyclic ring  
20 system, the bicyclic ring system is a naphthyl ring, and  
 $R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic  
group, phenyl, or -CN; or Ring D is an optionally  
substituted ring selected from phenyl, pyridinyl,  
piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl,  
25 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-  
tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-  
dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or  
naphthyl;

(b)  $R^y$  is  $T-R^{3'}$ , wherein T is a valence bond or  
30 a methylene and  $R^{3'}$  is an optionally substituted group  
selected from  $C_{1-6}$  aliphatic,  $C_{3-6}$  carbocyclyl,  $C_{6-10}$  aryl, a  
heteroaryl ring having 5-10 ring atoms, or a heterocyclyl  
ring having 5-10 ring atoms;

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring; and

(d) Ring D is substituted by oxo or  $R^5$ , wherein each  $R^5$  is independently selected from -halo, -CN, -NO<sub>2</sub>, -N( $R^4$ )<sub>2</sub>, optionally substituted  $C_{1-6}$  aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH( $R^4$ ), -N( $R^4$ )COR, -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, or -N( $R^4$ )SO<sub>2</sub>R.

Even more preferred compounds of formula VI have one or more, and more preferably all, of the features selected from the group consisting of:

(a)  $R^Y$  is T- $R^{3'}$ , wherein T is a valence bond or a methylene and  $R^{3'}$  is an optionally substituted group selected from  $C_{1-4}$  aliphatic,  $C_{3-6}$  carbocyclyl, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring;


(b) Ring C is a phenyl or pyridinyl ring, optionally substituted by - $R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or naphthyl;

(c)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring optionally substituted with -halo, -N( $R^4$ )<sub>2</sub>, - $C_{1-4}$  alkyl, - $C_{1-4}$  haloalkyl, -NO<sub>2</sub>, -O( $C_{1-4}$  alkyl), -CO<sub>2</sub>( $C_{1-4}$  alkyl), -CN,

-SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -OC(O)NH<sub>2</sub>, -NH<sub>2</sub>SO<sub>2</sub>(C<sub>1-4</sub> alkyl), -NHC(O)(C<sub>1-4</sub> alkyl), -C(O)NH<sub>2</sub>, or -CO(C<sub>1-4</sub> alkyl), wherein the (C<sub>1-4</sub> alkyl) is a straight, branched, or cyclic alkyl group; and

- 5 (d) Ring D is substituted by oxo or R<sup>5</sup>, wherein each R<sup>5</sup> is independently selected from -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic).

Another embodiment of this invention relates to  
10 compounds of formula VIa:



VIa

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

G is Ring C or Ring D;

- 15 Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from -R<sup>1</sup>, any substitutable non-ortho carbon position on Ring C is independently  
20 substituted by -R<sup>5</sup>, and two adjacent substituents on Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen,  
25 said fused ring being optionally substituted by halo, oxo, or -R<sup>8</sup>;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or -R<sup>5</sup>, and at any substitutable ring nitrogen by -R<sup>4</sup>, provided that when Ring D is a six-membered aryl or heteroaryl ring, -R<sup>5</sup> is hydrogen at each ortho carbon position of Ring D;

R<sup>1</sup> is selected from -halo, -CN, -NO<sub>2</sub>, T-V-R<sup>6</sup>, phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or C<sub>1-6</sub> aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or -R<sup>8</sup>, said C<sub>1-6</sub> aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or R<sup>1</sup> and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

T is a valence bond or a C<sub>1-4</sub> alkylidene chain;

R<sup>2</sup> and R<sup>2'</sup> are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable carbon on said fused ring formed by R<sup>2</sup> and R<sup>2'</sup> is substituted by halo, oxo, -CN, -NO<sub>2</sub>, -R<sup>7</sup>, or -V-R<sup>6</sup>, and any substitutable nitrogen on said ring formed by R<sup>2</sup> and R<sup>2'</sup> is substituted by R<sup>4</sup>;

each R is independently selected from hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-\text{COR}^7$ ,  
 $-\text{CO}_2(\text{optionally substituted } \text{C}_{1-6} \text{ aliphatic})$ ,  $-\text{CON}(\text{R}^7)_2$ ,  
 or  $-\text{SO}_2\text{R}^7$ , or two  $R^4$  on the same nitrogen are taken  
 together to form a 5-8 membered heterocyclyl or  
 5 heteroaryl ring;

each  $R^5$  is independently selected from  $-R$ , halo,  $-\text{OR}$ ,  
 $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{SO}_2\text{R}$ ,  $-\text{SR}$ ,  
 $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^4)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^4)\text{COR}$ ,  
 $-\text{N}(\text{R}^4)\text{CO}_2(\text{optionally substituted } \text{C}_{1-6} \text{ aliphatic})$ ,  
 10  $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$ ,  
 $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$ , or  $R^5$  and  
 an adjacent substituent taken together with their  
 intervening atoms form said ring fused to Ring C;

V is  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2-$ ,  $-\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 15  $-\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{N}(\text{R}^6)\text{CO}-$ ,  $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  
 $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  
 20  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ , or  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ;

W is  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  
 $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$ ,  $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$ ,  
 25  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ , or  $-\text{CON}(\text{R}^6)-$ ;


each  $R^6$  is independently selected from hydrogen, an  
 optionally substituted  $\text{C}_{1-4}$  aliphatic group, or two  $R^6$   
 30 groups on the same nitrogen atom are taken together  
 with the nitrogen atom to form a 5-6 membered  
 heterocyclyl or heteroaryl ring;

each  $R^7$  is independently selected from hydrogen or an  
 optionally substituted  $\text{C}_{1-6}$  aliphatic group, or two  $R^7$

on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring; and

each  $R^8$  is independently selected from an optionally substituted  $C_{1-4}$  aliphatic group,  $-OR^6$ ,  $-SR^6$ ,  $-COR^6$ ,  $-SO_2R^6$ ,  $-N(R^6)_2$ ,  $-N(R^6)N(R^6)_2$ ,  $-CN$ ,  $-NO_2$ ,  $-CON(R^6)_2$ , or  $-CO_2R^6$ .

Preferred rings formed by the  $R^2$  and  $R^{2'}$  groups of formula VIa include benzo, pyrido, pyrimido, and a partially unsaturated 6-membered carbocyclo ring. These are exemplified in the following formula VIa compounds having a pyrazole-containing bicyclic ring system:



Preferred substituents on the  $R^2/R^{2'}$  fused ring include one or more of the following:  $-\text{halo}$ ,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl),  $-CN$ ,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , and  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the  $(C_{1-4}$  alkyl) group is methyl.

When G is Ring C, preferred formula VIa Ring C groups are phenyl and pyridinyl. When two adjacent substituents on Ring C are taken together to form a fused ring, Ring C is contained in a bicyclic ring system. Preferred fused rings include a benzo or pyrido ring. Such rings preferably are fused at ortho and meta

positions of Ring C. Examples of preferred bicyclic Ring C systems include naphthyl and isoquinolinyl. Preferred R<sup>1</sup> groups include -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>,  
5 -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>. When R<sup>1</sup> is an optionally substituted C<sub>1-6</sub> aliphatic group, the most preferred optional substituents are halogen. Examples of preferred R<sup>1</sup> groups include -CF<sub>3</sub>, -Cl, -F, -CN, -COCH<sub>3</sub>, -OCH<sub>3</sub>, -OH, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -CH<sub>3</sub>, -CF<sub>2</sub>CH<sub>3</sub>,  
10 cyclohexyl, t-butyl, isopropyl, cyclopropyl, -C≡CH, -C≡C-CH<sub>3</sub>, -SO<sub>2</sub>CH<sub>3</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, -CO<sub>2</sub>CH<sub>3</sub>, -CONH<sub>2</sub>, -NHCOCH<sub>3</sub>, -OC(O)NH<sub>2</sub>, -NHSO<sub>2</sub>CH<sub>3</sub>, and -OCF<sub>3</sub>.

On Ring C preferred R<sup>5</sup> substituents, when present, include -halo, -CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, optionally  
15 substituted C<sub>1-6</sub> aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, and -N(R<sup>4</sup>)SO<sub>2</sub>R. More preferred R<sup>5</sup> substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic).  
20 Examples of such preferred R<sup>5</sup> substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHMe, -NMe<sub>2</sub>, -OEt, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, and -CO<sub>2</sub>Et.

When G is Ring D, preferred formula VIa Ring D monocyclic rings include substituted and unsubstituted  
25 phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, and morpholinyl rings. When two adjacent substituents on Ring D are taken together to form a fused ring, the Ring D system is bicyclic. Preferred formula VIa Ring D bicyclic rings  
30 include 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, and naphthyl. Examples of more preferred bicyclic Ring D systems include naphthyl and isoquinolinyl.



Preferred substituents on the formula **Via** Ring D include one or more of the following: halo, oxo, CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -SR, -OR, -C(O)R, or substituted or  
5 unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. More preferred Ring D substituents include -halo, -CN, -oxo, -SR, -OR, -N(R<sup>4</sup>)<sub>2</sub>, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub>  
10 aliphatic. Examples of Ring D substituents include -OH, phenyl, methyl, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH, pyrrolidinyl, OPh, CF<sub>3</sub>, C≡CH, Cl, Br, F, I, NH<sub>2</sub>, C(O)CH<sub>3</sub>, *i*-propyl, *tert*-butyl, SEt, OMe, N(Me)<sub>2</sub>, methylene dioxy, and ethylene dioxy.

Preferred formula **Via** compounds have one or  
15 more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by -R<sup>5</sup>, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring  
20 system, the bicyclic ring system is selected from a naphthyl, quinolinyl or isoquinolinyl ring, and R<sup>1</sup> is -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>; or Ring D is an  
25 optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1*H*-isoindolyl, 2,3-dihydro-1*H*-indolyl,  
30 isoquinolinyl, quinolinyl, or naphthyl ring; and

(b) R<sup>2</sup> and R<sup>2'</sup> are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring.

More preferred compounds of formula VIa have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclic ring optionally substituted with -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl), -CN,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , and  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group; and

(c) Ring D is substituted by oxo or  $R^5$ , wherein each  $R^5$  is independently selected from -halo, -CN,  $-NO_2$ ,  $-N(R^4)_2$ , optionally substituted  $C_{1-6}$  aliphatic group, -OR,  $-C(O)R$ ,  $-CO_2R$ ,  $-CONH(R^4)$ ,  $-N(R^4)COR$ ,  $-SO_2N(R^4)_2$ , or  $-N(R^4)SO_2R$ .

Even more preferred compounds of formula VIa have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two


adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, or partially unsaturated 6-membered carbocyclo ring optionally substituted with -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl), -CN,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , or  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group; and


(d) Ring D is substituted by oxo or  $R^5$ , wherein each  $R^5$  is independently selected from -Cl, -F, -CN,  $-CF_3$ ,  $-NH_2$ ,  $-NH(C_{1-4}$  aliphatic),  $-N(C_{1-4}$  aliphatic) $_2$ ,  $-O(C_{1-4}$  aliphatic),  $C_{1-4}$  aliphatic, and  $-CO_2(C_{1-4}$  aliphatic).

Representative compounds of formula VI and IVa are set forth in Table 5 below.


Table 5.




VI-1




VI-2




VI-3




VI-4




VI-5




VI-6




VI-7




VI-8




VI-9




VI-10




VI-11




VI-12




VI-13




VI-14




VI-15




VI-16




VI-17




VI-18




VI-19




VI-20




VI-21




VI-22




VI-23




VI-24




VI-25




VI-26




VI-27




VI-28




VI-29




VI-30



VI-31




VI-32




VI-33


5




VI-34




VI-35




VI-36



VI-37




VI-38




VI-39


10




VI-40




VI-41




VI-42



VI-43




VI-44




VI-45


5




VIa-1




VIa-2




VIa-3



VIa-4




VIa-5




VIa-6


10




VIA-7




VIA-8



VIA-9



VIA-10



VIA-11



VIA-12

In another embodiment, this invention provides a composition comprising a compound of formula VI or VIA and a pharmaceutically acceptable carrier.

One aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VI or VIA.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VI or VIA.

Another aspect relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective



amount of a composition comprising a compound of formula VI or VIa. This method is especially useful for diabetic patients.

Another aspect relates to a method of  
5 inhibiting the production of hyperphosphorylated Tau protein in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula VI or VIa. This method is especially useful in halting  
10 or slowing the progression of Alzheimer's disease.

Another aspect relates to a method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition  
15 comprising a compound of formula VI or VIa. This method is especially useful for treating schizophrenia.

One aspect of this invention relates to a method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically  
20 effective amount of a composition comprising a compound of formula VI or VIa.

Another aspect relates to a method of treating a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of  
25 administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VI or VIa. This method is especially useful for treating cancer, such as colon, ovarian, and breast cancer.


30 One aspect of this invention relates to a method of inhibiting CDK-2 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VI or VIa.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a CDK-2 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VI or VIa. This method is especially useful for treating cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis.

Another method relates to inhibiting GSK-3, Aurora, or CDK-2 activity in a biological sample, which method comprises contacting the biological sample with the GSK-3 or Aurora inhibitor of formula VI or VIa, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3, Aurora or CDK-2.

Each of the aforementioned methods directed to the inhibition of GSK-3, Aurora or CDK-2, or the treatment of a disease alleviated thereby, is preferably carried out with a preferred compound of formula VI or VIa, as described above.

Another embodiment of this invention relates to compounds of formula VII:



VII

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

G is Ring C or Ring D;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from  $-R^1$ , any substitutable non-ortho carbon position on Ring C is independently substituted by  $-R^5$ , and two adjacent substituents on Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo, oxo, or  $-R^8$ ;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or  $-R^5$ , and at any substitutable ring nitrogen by  $-R^4$ , provided that when Ring D is a six-membered aryl or heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^1$  is selected from -halo, -CN,  $-\text{NO}_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $\text{C}_{1-6}$  aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or  $-R^8$ , said  $\text{C}_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

- $R^y$  is hydrogen or  $T-R^{3''}$ ;  
 $T$  is a valence bond, hydrogen, or a  $C_{1-4}$  alkylidene chain;  
 $R^2$  and  $R^{2'}$  are independently selected from  $-R$ ,  $-T-W-R^6$ , or  
 $R^2$  and  $R^{2'}$  are taken together with their intervening  
atoms to form a fused, 5-8 membered, unsaturated or  
partially unsaturated, ring having 0-3 ring heteroatoms  
5 selected from nitrogen, oxygen, or sulfur, wherein each  
substitutable carbon on said fused ring formed by  $R^2$   
and  $R^{2'}$  is substituted by halo, oxo,  $-CN$ ,  $-NO_2$ ,  $-R^7$ , or  
 $-V-R^6$ , and any substitutable nitrogen on said ring  
formed by  $R^2$  and  $R^{2'}$  is substituted by  $R^4$ ;  
 $R^{3''}$  is selected from an optionally substituted group  
selected from  $C_{3-10}$  carbocyclyl,  $C_{6-10}$  aryl, a heteroaryl  
ring having 5-10 ring atoms, or a heterocyclyl ring  
having 5-10 ring atoms;  
each  $R$  is independently selected from hydrogen or an  
optionally substituted group selected from  $C_{1-6}$   
aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10  
ring atoms, or a heterocyclyl ring having 5-10 ring  
atoms;  
10 each  $R^4$  is independently selected from  $-R^7$ ,  $-COR^7$ ,  
 $-CO_2$ (optionally substituted  $C_{1-6}$  aliphatic),  $-CON(R^7)_2$ ,  
or  $-SO_2R^7$ , or two  $R^4$  on the same nitrogen are taken  
together to form a 5-8 membered heterocyclyl or  
heteroaryl ring;  
15 each  $R^5$  is independently selected from  $-R$ , halo,  $-OR$ ,  
 $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  
 $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  
 $-N(R^4)CO_2$ (optionally substituted  $C_{1-6}$  aliphatic),  
 $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  
20  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ , or  $R^5$  and  
an adjacent substituent taken together with their  
intervening atoms form said ring fused to Ring C;

V is -O-, -S-, -SO-, -SO<sub>2</sub>-, -N(R<sup>6</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>6</sup>)-,  
 -N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -N(R<sup>6</sup>)CO-, -N(R<sup>6</sup>)C(O)O-,  
 -N(R<sup>6</sup>)CON(R<sup>6</sup>)-, -N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)N(R<sup>6</sup>)-,  
 -C(O)N(R<sup>6</sup>)-, -OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-,  
 5 -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-,  
 -C(R<sup>6</sup>)=N-O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, or  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-;

W is -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-,  
 10 -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-,  
 -C(R<sup>6</sup>)OC(O)-, -C(R<sup>6</sup>)OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CO-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-, -C(R<sup>6</sup>)=N-O-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-,  
 -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-, or -CON(R<sup>6</sup>)-;

15 each R<sup>6</sup> is independently selected from hydrogen, an  
 optionally substituted C<sub>1-4</sub> aliphatic group, or two R<sup>6</sup>  
 groups on the same nitrogen atom are taken together  
 with the nitrogen atom to form a 5-6 membered  
 heterocyclyl or heteroaryl ring;


20 each R<sup>7</sup> is independently selected from hydrogen or an  
 optionally substituted C<sub>1-6</sub> aliphatic group, or two R<sup>7</sup>  
 on the same nitrogen are taken together with the  
 nitrogen to form a 5-8 membered heterocyclyl or  
 heteroaryl ring;

25 each R<sup>8</sup> is independently selected from an optionally  
 substituted C<sub>1-4</sub> aliphatic group, -OR<sup>6</sup>, -SR<sup>6</sup>, -COR<sup>6</sup>,  
 -SO<sub>2</sub>R<sup>6</sup>, -N(R<sup>6</sup>)<sub>2</sub>, -N(R<sup>6</sup>)N(R<sup>6</sup>)<sub>2</sub>, -CN, -NO<sub>2</sub>, -CON(R<sup>6</sup>)<sub>2</sub>, or  
 -CO<sub>2</sub>R<sup>6</sup>; and

R<sup>9</sup> is selected from -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR,  
 30 -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>,  
 -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally  
 substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>,  
 -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or  
 -OC(=O)N(R<sup>4</sup>)<sub>2</sub>.

Preferred  $R^Y$  groups of formula VII include  $T-R^{3''}$  wherein T is a valence bond or a methylene. Preferred  $R^{3''}$  groups include an optionally substituted group selected from  $C_{3-6}$  carbocyclyl, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring. Examples of preferred  $R^Y$  include 2-pyridyl, 4-pyridyl, piperidinyl, cyclopropyl, and an optionally substituted phenyl such as phenyl or halo-substituted phenyl.

The  $R^2$  and  $R^{2'}$  groups of formula VII may be taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring. Preferred fused rings include benzo, pyrido, pyrimido, and a partially unsaturated 6-membered carbocyclo ring. These are exemplified in the following formula VII compounds having a pyrazole-containing bicyclic ring system:



Preferred substituents on the  $R^2/R^{2'}$  fused ring include one or more of the following: -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl),  $-CN$ ,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , and  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the  $(C_{1-4}$  alkyl) group is methyl.

When the pyrazole ring system of formula VII is monocyclic, preferred  $R^2$  groups include hydrogen,  $C_{1-4}$

aliphatic, alkoxycarbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred R<sup>2</sup> substituents include methyl, cyclopropyl, ethyl, isopropyl, propyl, t-butyl, cyclopentyl, phenyl, CO<sub>2</sub>H, CO<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>OH, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>Ph, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHCOOC(CH<sub>3</sub>)<sub>3</sub>, CONHCH(CH<sub>3</sub>)<sub>2</sub>, CONHCH<sub>2</sub>CH=CH<sub>2</sub>, CONHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CONHCH<sub>2</sub>Ph, CONH(cyclohexyl), CON(Et)<sub>2</sub>, CON(CH<sub>3</sub>)CH<sub>2</sub>Ph, CONH(n-C<sub>3</sub>H<sub>7</sub>), CON(Et)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, CONHCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, CON(n-C<sub>3</sub>H<sub>7</sub>)<sub>2</sub>, CO(3-methoxymethylpyrrolidin-1-yl), CONH(3-tolyl), CONH(4-tolyl), CONHCH<sub>3</sub>, CO(morpholin-1-yl), CO(4-methylpiperazin-1-yl), CONHCH<sub>2</sub>CH<sub>2</sub>OH, CONH<sub>2</sub>, and CO(piperidin-1-yl). A preferred R<sup>2'</sup> group is hydrogen.

When G is Ring C, preferred formula VII Ring C groups are phenyl and pyridinyl. When two adjacent substituents on Ring C are taken together to form a fused ring, Ring C is contained in a bicyclic ring system. Preferred fused rings include a benzo or pyrido ring. Such rings preferably are fused at ortho and meta positions of Ring C. Examples of preferred bicyclic Ring C systems include naphthyl and isoquinolinyl. Preferred R<sup>1</sup> groups include -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>. When R<sup>1</sup> is an optionally substituted C<sub>1-6</sub> aliphatic group, the most preferred optional substituents are halogen. Examples of preferred R<sup>1</sup> groups include -CF<sub>3</sub>, -Cl, -F, -CN, -COCH<sub>3</sub>, -OCH<sub>3</sub>, -OH, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -CH<sub>3</sub>, -CF<sub>2</sub>CH<sub>3</sub>, cyclohexyl, t-butyl, isopropyl, cyclopropyl, -C≡CH, -C≡C-CH<sub>3</sub>, -SO<sub>2</sub>CH<sub>3</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, -CO<sub>2</sub>CH<sub>3</sub>, -CONH<sub>2</sub>, -NHCOCH<sub>3</sub>, -OC(O)NH<sub>2</sub>, -NHSO<sub>2</sub>CH<sub>3</sub>, and -OCF<sub>3</sub>.

On Ring C preferred  $R^5$  substituents, when present, include -halo, -CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, optionally substituted C<sub>1-6</sub> aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, and -N(R<sup>4</sup>)SO<sub>2</sub>R. More preferred  $R^5$  substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic). Examples of such preferred  $R^5$  substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHMe, -NMe<sub>2</sub>, -OEt, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, and -CO<sub>2</sub>Et.

When G is Ring D, preferred formula VII Ring D monocyclic rings include substituted and unsubstituted phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, and morpholinyl rings. When two adjacent substituents on Ring D are taken together to form a fused ring, the Ring D system is bicyclic. Preferred formula VII Ring D bicyclic rings include 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, and naphthyl. Examples of more preferred bicyclic Ring D systems include naphthyl and isoquinolinyl.

Preferred substituents on Ring D include one or more of the following: halo, oxo, CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -SR, -OR, -C(O)R, or substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. More preferred Ring D substituents include -halo, -CN, -oxo, -SR, -OR, -N(R<sup>4</sup>)<sub>2</sub>, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. Examples of Ring D substituents include -OH, phenyl, methyl, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH, pyrrolidinyl, OPh, CF<sub>3</sub>, C≡CH, Cl,



Br, F, I, NH<sub>2</sub>, C(O)CH<sub>3</sub>, *i*-propyl, *tert*-butyl, SEt, OMe, N(Me)<sub>2</sub>, methylene dioxy, and ethylene dioxy.

Preferred formula VII compounds have one or more, and more preferably all, of the features selected  
5 from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by -R<sup>5</sup>, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a  
10 naphthyl, quinolinyl or isoquinolinyl ring, and R<sup>1</sup> is -halo, an optionally substituted C<sub>1-6</sub> aliphatic group; phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>; or Ring D is an  
15 optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1*H*-isoindolyl, 2,3-dihydro-1*H*-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

20 (b) R<sup>y</sup> is T-R<sup>3y</sup>, wherein T is a valence bond or a methylene; and

(c) R<sup>2'</sup> is hydrogen and R<sup>2</sup> is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a C<sub>1-6</sub> aliphatic group, or R<sup>2</sup> and R<sup>2'</sup> are  
25 taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring.

More preferred compounds of formula VII have one or more, and more preferably all, of the features  
30 selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by -R<sup>5</sup>, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and

$R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^Y$  is  $T-R^{3''}$ , wherein T is a valence bond or a methylene and  $R^{3''}$  is an optionally substituted group selected from  $C_{3-6}$  carbocyclyl, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring;

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring; and

(d) Ring D is substituted by oxo or  $R^5$ , wherein each  $R^5$  is independently selected from -halo, -CN, -NO<sub>2</sub>, -N( $R^4$ )<sub>2</sub>, optionally substituted  $C_{1-6}$  aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH( $R^4$ ), -N( $R^4$ )COR, -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, or -N( $R^4$ )SO<sub>2</sub>R.

Even more preferred compounds of formula VII have one or more, and more preferably all, of the features selected from the group consisting of:

(a)  $R^Y$  is  $T-R^{3''}$ , wherein T is a valence bond or a methylene and  $R^{3''}$  is an optionally substituted group selected from phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring;

(b) Ring C is a phenyl or pyridinyl ring, optionally substituted by - $R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and

$R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or naphthyl;


(c)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring optionally substituted with -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl), -CN,  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , or  $-CO(C_{1-4}$  alkyl), wherein the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group; and

(d) Ring D is substituted by oxo or  $R^5$ , wherein each  $R^5$  is independently selected from -Cl, -F, -CN,  $-CF_3$ ,  $-NH_2$ ,  $-NH(C_{1-4}$  aliphatic),  $-N(C_{1-4}$  aliphatic) $_2$ ,  $-O(C_{1-4}$  aliphatic),  $C_{1-4}$  aliphatic, and  $-CO_2(C_{1-4}$  aliphatic).


Representative compounds of formula VII are set forth in Table 6 below.

Table 6.


25




VII-1




VII-2




VII-3




VII-4




VII-5




VII-6




VII-7




VII-8




VII-9




VII-10




VII-11




VII-12




VII-13




VII-14




VII-15




VII-16




VII-17




VII-18




VII-19




VII-20




VII-21




VII-22




VII-23




VII-24




VII-25




VII-26




VII-27




VII-28




VII-29




VII-30




VII-31




VII-32




VII-33



VII-34



VII-35



VII-36

5

10

In another embodiment, this invention provides a composition comprising a compound of formula VII and a pharmaceutically acceptable carrier.

15

One aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VII.

20

Another aspect relates to a method of treating a disease that is alleviated by treatment with a GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a

therapeutically effective amount of a composition comprising a compound of formula VII.

Another aspect relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula VII. This method is especially useful for diabetic patients.

Another aspect relates to a method of inhibiting the production of hyperphosphorylated Tau protein in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula VII. This method is especially useful in halting or slowing the progression of Alzheimer's disease.

Another aspect relates to a method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula VII. This method is especially useful for treating schizophrenia.

One aspect of this invention relates to a method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VII.

Another aspect relates to a method of treating a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VII. This method is

especially useful for treating cancer, such as colon, ovarian, and breast cancer.

One aspect of this invention relates to a method of inhibiting CDK-2 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VII.


Another aspect relates to a method of treating a disease that is alleviated by treatment with a CDK-2 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VII. This method is especially useful for treating cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis.

Another method relates to inhibiting GSK-3, Aurora, or CDK-2 activity in a biological sample, which method comprises contacting the biological sample with the GSK-3 or Aurora inhibitor of formula VII, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3, Aurora or CDK-2.

Each of the aforementioned methods directed to the inhibition of GSK-3, Aurora or CDK-2, or the treatment of a disease alleviated thereby, is preferably carried out with a preferred compound of formula VII, as described above.

Another embodiment of this invention relates to compounds of formula VIII:





VIII

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

$Z^1$  is N or  $CR^9$ ,  $Z^2$  is N or CH, and  $Z^3$  is N or  $CR^x$ , provided that one of  $Z^1$  and  $Z^3$  is nitrogen;

G is Ring C or Ring D;

- 5 Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from  $-R^1$ , any substitutable non-ortho carbon position on Ring C is independently substituted by  $-R^5$ , and two adjacent substituents on
- 10 Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo,
- 15 oxo, or  $-R^8$ ;

- Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or
- 20 heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by halo, oxo, or  $-R^5$ , and at any substitutable ring nitrogen by  $-R^4$ , provided that when Ring D is a six-membered aryl

or heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^1$  is selected from -halo, -CN,  $-NO_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $C_{1-6}$  aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or  $-R^8$ , said  $C_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

$R^x$  is T- $R^3$ ;

T is a valence bond or a  $C_{1-4}$  alkylidene chain;

$R^2$  and  $R^{2'}$  are independently selected from -R, -T-W- $R^6$ , or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable carbon on said fused ring formed by  $R^2$  and  $R^{2'}$  is substituted by halo, oxo, -CN,  $-NO_2$ ,  $-R^7$ , or -V- $R^6$ , and any substitutable nitrogen on said ring formed by  $R^2$  and  $R^{2'}$  is substituted by  $R^4$ ;

$R^3$  is selected from -R, -halo, -OR,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-COCH_2COR$ ,  $-NO_2$ , -CN,  $-S(O)R$ ,  $-S(O)_2R$ , -SR,  $-N(R^4)_2$ ,  $-CON(R^7)_2$ ,  $-SO_2N(R^7)_2$ ,  $-OC(=O)R$ ,  $-N(R^7)COR$ ,  $-N(R^7)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^7)CON(R^7)_2$ ,  $-N(R^7)SO_2N(R^7)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^7)_2$ ;

each R is independently selected from hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-\text{COR}^7$ ,  
 $-\text{CO}_2$  (optionally substituted  $\text{C}_{1-6}$  aliphatic),  $-\text{CON}(\text{R}^7)_2$ ,  
 or  $-\text{SO}_2\text{R}^7$ , or two  $R^4$  on the same nitrogen are taken  
 together to form a 5-8 membered heterocyclyl or  
 5 heteroaryl ring;

each  $R^5$  is independently selected from  $-R$ , halo,  $-\text{OR}$ ,  
 $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{SO}_2\text{R}$ ,  $-\text{SR}$ ,  
 $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^4)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^4)\text{COR}$ ,  
 $-\text{N}(\text{R}^4)\text{CO}_2$  (optionally substituted  $\text{C}_{1-6}$  aliphatic),  
 10  $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$ ,  
 $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$ , or  $R^5$  and  
 an adjacent substituent taken together with their  
 intervening atoms form said ring fused to Ring C;

V is  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2-$ ,  $-\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 15  $-\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{N}(\text{R}^6)\text{CO}-$ ,  $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  
 $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  
 20  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ , or  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ;

W is  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  
 $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$ ,  $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$ ,  
 25  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ , or  $-\text{CON}(\text{R}^6)-$ ;

each  $R^6$  is independently selected from hydrogen, an  
 optionally substituted  $\text{C}_{1-4}$  aliphatic group, or two  $R^6$   
 30 groups on the same nitrogen atom are taken together  
 with the nitrogen atom to form a 5-6 membered  
 heterocyclyl or heteroaryl ring;


each  $R^7$  is independently selected from hydrogen or an  
 optionally substituted  $\text{C}_{1-6}$  aliphatic group, or two  $R^7$

on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring;

each  $R^8$  is independently selected from an optionally substituted  $C_{1-4}$  aliphatic group,  $-OR^6$ ,  $-SR^6$ ,  $-COR^6$ ,  $-SO_2R^6$ ,  $-N(R^6)_2$ ,  $-N(R^6)N(R^6)_2$ ,  $-CN$ ,  $-NO_2$ ,  $-CON(R^6)_2$ , or  $-CO_2R^6$ ; and

$R^9$  is selected from  $-R$ , halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ .

Accordingly, the present invention relates to compounds of formula VIIIa, VIIIb, VIIIc and VIId as shown below:




Preferred  $R^x$  groups of formula VIII include  $T-R^3$  wherein  $T$  is a valence bond or a methylene and  $R^3$  is  $CN$ ,  $-R$ , or  $-OR$ . When  $R^3$  is  $-R$ , preferred  $R^3$  groups include an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring. When  $R^3$  is  $-OR$ , preferred  $R$  groups include an optionally substituted group  $C_{1-6}$  aliphatic group such as alkyl- or dialkylaminoalkyl and aminoalkyl. Examples of

preferred  $R^x$  include acetamido, CN, piperidinyl, piperazinyl, phenyl, pyridinyl, imidazol-1-yl, imidazol-2-yl, cyclohexyl, cyclopropyl, methyl, ethyl, isopropyl, t-butyl,  $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}$ , and  $\text{NH}_2\text{CH}_2\text{CH}_2\text{O}$ .

5 Preferred  $R^9$  groups of formula VIII, when present, include R, OR, and  $\text{N}(\text{R}^4)_2$ . Examples of preferred  $R^9$  include methyl, ethyl,  $\text{NH}_2$ ,  $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}$ ,  $\text{N}(\text{CH}_3)_2\text{CH}_2\text{CH}_2\text{NH}$ ,  $\text{N}(\text{CH}_3)_2\text{CH}_2\text{CH}_2\text{O}$ , (piperidin-1-yl) $\text{CH}_2\text{CH}_2\text{O}$ , and  $\text{NH}_2\text{CH}_2\text{CH}_2\text{O}$ .

The  $R^2$  and  $R^{2'}$  groups of formula VIII may be  
 10 taken together to form a fused ring, thus providing a bicyclic ring system containing a pyrazole ring.  
 Preferred fused rings include benzo, pyrido, pyrimido, and a partially unsaturated 6-membered carbocyclo ring.  
 These are exemplified in the following formula VIII  
 15 compounds having a pyrazole-containing bicyclic ring system:



20 Preferred substituents on the formula VIII  $R^2/R^{2'}$  fused ring include one or more of the following:  
 -halo,  $-\text{N}(\text{R}^4)_2$ ,  $-\text{C}_{1-4}$  alkyl,  $-\text{C}_{1-4}$  haloalkyl,  $-\text{NO}_2$ ,  $-\text{O}(\text{C}_{1-4}$  alkyl),  $-\text{CO}_2(\text{C}_{1-4}$  alkyl),  $-\text{CN}$ ,  $-\text{SO}_2(\text{C}_{1-4}$  alkyl),  $-\text{SO}_2\text{NH}_2$ ,  
 $-\text{OC}(\text{O})\text{NH}_2$ ,  $-\text{NH}_2\text{SO}_2(\text{C}_{1-4}$  alkyl),  $-\text{NHC}(\text{O})(\text{C}_{1-4}$  alkyl),  
 25  $-\text{C}(\text{O})\text{NH}_2$ , and  $-\text{CO}(\text{C}_{1-4}$  alkyl), wherein the  $(\text{C}_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl group. Preferably, the  $(\text{C}_{1-4}$  alkyl) group is methyl.

When the pyrazole ring system of formula VIII is monocyclic, preferred  $R^2$  groups include hydrogen,  $\text{C}_{1-4}$

aliphatic, alkoxycarbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred R<sup>2</sup> substituents include methyl, cyclopropyl, ethyl, isopropyl, propyl, t-butyl, cyclopentyl, phenyl, CO<sub>2</sub>H, CO<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>OH, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>Ph, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHCOOC(CH<sub>3</sub>)<sub>3</sub>, CONHCH(CH<sub>3</sub>)<sub>2</sub>, CONHCH<sub>2</sub>CH=CH<sub>2</sub>, CONHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CONHCH<sub>2</sub>Ph, CONH(cyclohexyl), CON(Et)<sub>2</sub>, CON(CH<sub>3</sub>)CH<sub>2</sub>Ph, CONH(n-C<sub>3</sub>H<sub>7</sub>), CON(Et)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, CONHCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, CON(n-C<sub>3</sub>H<sub>7</sub>)<sub>2</sub>, CO(3-methoxymethylpyrrolidin-1-yl), CONH(3-tolyl), CONH(4-tolyl), CONHCH<sub>3</sub>, CO(morpholin-1-yl), CO(4-methylpiperazin-1-yl), CONHCH<sub>2</sub>CH<sub>2</sub>OH, CONH<sub>2</sub>, and CO(piperidin-1-yl). A preferred R<sup>2'</sup> group is hydrogen.

When G is Ring C, preferred formula VIII Ring C groups are phenyl and pyridinyl. When two adjacent substituents on Ring C are taken together to form a fused ring, Ring C is contained in a bicyclic ring system. Preferred fused rings include a benzo or pyrido ring. Such rings preferably are fused at ortho and meta positions of Ring C. Examples of preferred bicyclic Ring C systems include naphthyl and isoquinolinyl. Preferred R<sup>1</sup> groups include -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>. When R<sup>1</sup> is an optionally substituted C<sub>1-6</sub> aliphatic group, the most preferred optional substituents are halogen. Examples of preferred R<sup>1</sup> groups include -CF<sub>3</sub>, -Cl, -F, -CN, -COCH<sub>3</sub>, -OCH<sub>3</sub>, -OH, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>2</sub>CH<sub>3</sub>, -CH<sub>3</sub>, -CF<sub>2</sub>CH<sub>3</sub>, cyclohexyl, t-butyl, isopropyl, cyclopropyl, -C≡CH, -C≡C-CH<sub>3</sub>, -SO<sub>2</sub>CH<sub>3</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, -CO<sub>2</sub>CH<sub>3</sub>, -CONH<sub>2</sub>, -NHCOCH<sub>3</sub>, -OC(O)NH<sub>2</sub>, -NHSO<sub>2</sub>CH<sub>3</sub>, and -OCF<sub>3</sub>.

On Ring C preferred  $R^5$  substituents, when present, include -halo, -CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, optionally substituted C<sub>1-6</sub> aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, and -N(R<sup>4</sup>)SO<sub>2</sub>R. More preferred  $R^5$  substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic). Examples of such preferred  $R^5$  substituents include -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHMe, -NMe<sub>2</sub>, -OEt, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, and -CO<sub>2</sub>Et.

When G is Ring D, preferred formula VIII Ring D monocyclic rings include substituted and unsubstituted phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, and morpholinyl rings. When two adjacent substituents on Ring D are taken together to form a fused ring, the Ring D system is bicyclic. Preferred formula VIII Ring D bicyclic rings include 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, and naphthyl. Examples of more preferred bicyclic Ring D systems include naphthyl and isoquinolinyl.

Preferred  $R^5$  substituents on Ring D of formula VIII include halo, oxo, CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -SR, -OR, -C(O)R, or substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. More preferred  $R^5$  substituents include -halo, -CN, -oxo, -SR, -OR, -N(R<sup>4</sup>)<sub>2</sub>, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl, C<sub>6-10</sub> aryl, or C<sub>1-6</sub> aliphatic. Examples of Ring D substituents include -OH, phenyl, methyl, CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>OH, pyrrolidinyl, OPh, CF<sub>3</sub>, C≡CH, Cl, Br, F, I, NH<sub>2</sub>,

C(O)CH<sub>3</sub>, *i*-propyl, *tert*-butyl, SEt, OMe, N(Me)<sub>2</sub>, methylene dioxy, and ethylene dioxy.

Preferred formula VIII compounds have one or more, and more preferably all, of the features selected  
5 from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by -R<sup>5</sup>, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a  
10 naphthyl, quinolinyll or isoquinolinyll ring, and R<sup>1</sup> is -halo, an optionally substituted C<sub>1-6</sub> aliphatic group, phenyl, -COR<sup>6</sup>, -OR<sup>6</sup>, -CN, -SO<sub>2</sub>R<sup>6</sup>, -SO<sub>2</sub>NH<sub>2</sub>, -N(R<sup>6</sup>)<sub>2</sub>, -CO<sub>2</sub>R<sup>6</sup>, -CONH<sub>2</sub>, -NHCOR<sup>6</sup>, -OC(O)NH<sub>2</sub>, or -NHSO<sub>2</sub>R<sup>6</sup>; or Ring D is an  
15 pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyll, 1,2,3,4-tetrahydroquinolinyll, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyll, quinolinyll, or naphthyl ring;

20 (b) R<sup>x</sup> is T-R<sup>3</sup> wherein T is a valence bond or a methylene; and

(c) R<sup>2'</sup> is hydrogen and R<sup>2</sup> is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a C<sub>1-6</sub> aliphatic group, or R<sup>2</sup> and R<sup>2'</sup> are  
25 taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring.

More preferred compounds of formula VIII have one or more, and more preferably all, of the features  
30 selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by -R<sup>5</sup>, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and



$R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^x$  is T- $R^3$  wherein T is a valence bond or a methylene and  $R^3$  is CN, -R or -OR;

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring; and

(d) each  $R^5$  is independently selected from -halo, -CN, -NO<sub>2</sub>, -N( $R^4$ )<sub>2</sub>, optionally substituted  $C_{1-6}$  aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH( $R^4$ ), -N( $R^4$ )COR, -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, or -N( $R^4$ )SO<sub>2</sub>R.

Even more preferred compounds of formula VIII have one or more, and more preferably all, of the features selected from the group consisting of:

(a)  $R^x$  is T- $R^3$  wherein T is a valence bond or a methylene and  $R^3$  is -R or -OR wherein R is an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring;

(b) Ring C is a phenyl or pyridinyl ring, optionally substituted by - $R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl,

piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl,  
1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-  
tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or  
naphthyl;


5 (c)  $R^2$  and  $R^{2'}$  are taken together with their  
intervening atoms to form a benzo, pyrido, pyrimido or  
partially unsaturated 6-membered carbocyclic ring  
optionally substituted with -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  
 $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4}$  alkyl),  $-CO_2(C_{1-4}$  alkyl),  $-CN$ ,  
10  $-SO_2(C_{1-4}$  alkyl),  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4}$  alkyl),  
 $-NHC(O)(C_{1-4}$  alkyl),  $-C(O)NH_2$ , or  $-CO(C_{1-4}$  alkyl), wherein  
the  $(C_{1-4}$  alkyl) is a straight, branched, or cyclic alkyl  
group;

(d) each  $R^5$  is independently selected from  $-Cl$ ,  
15  $-F$ ,  $-CN$ ,  $-CF_3$ ,  $-NH_2$ ,  $-NH(C_{1-4}$  aliphatic),  $-N(C_{1-4}$   
aliphatic) $_2$ ,  $-O(C_{1-4}$  aliphatic),  $C_{1-4}$  aliphatic, and  
 $-CO_2(C_{1-4}$  aliphatic); and


(e)  $R^9$  is  $R$ ,  $OR$ , or  $N(R^4)_2$ .

Representative compounds of formula VIII are  
20 set forth in Table 7 below.


Table 7.



VIII-1




VIII-2




VIII-3




VIII-4




VIII-5




VIII-6




VIII-7




VIII-8




VIII-9




VIII-10




VIII-11




VIII-12




VIII-13




VIII-14




VIII-15




VIII-16




VIII-17




VIII-18



VIII-19




VIII-20




VIII-21


5




VIII-22




VIII-23




VIII-24



VIII-25




VIII-26




VIII-27


10




VIII-28




VIII-29




VIII-30




VIII-31




VIII-32




VIII-33




VIII-34




VIII-35




VIII-36




VIII-37




VIII-38




VIII-39




VIII-40




VIII-41




VIII-42



VIII-43




VIII-44




VIII-45


5




VIII-46




VIII-47




VIII-48



VIII-49




VIII-50




VIII-51


10




VIII-52




VIII-53




VIII-54



VIII-55



VIII-56



VIII-57

In another embodiment, this invention provides a composition comprising a compound of formula VIII and a pharmaceutically acceptable carrier.

One aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VIII.

Another aspect relates to a method of treating a disease that is alleviated by treatment with a GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VIII.

Another aspect relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula

VIII. This method is especially useful for diabetic patients.

Another aspect relates to a method of inhibiting the production of hyperphosphorylated Tau protein in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula VIII. This method is especially useful in halting or slowing the progression of Alzheimer's disease.

Another aspect relates to a method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula VIII. This method is especially useful for treating schizophrenia.

One aspect of this invention relates to a method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VIII.

Another aspect relates to a method of treating a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VIII. This method is especially useful for treating cancer, such as colon, ovarian, and breast cancer.

One aspect of this invention relates to a method of inhibiting CDK-2 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula VIII.




Another aspect relates to a method of treating a disease that is alleviated by treatment with a CDK-2 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula VIII. This method is especially useful for treating cancer, Alzheimer's disease, restenosis, angiogenesis, glomerulonephritis, cytomegalovirus, HIV, herpes, psoriasis, atherosclerosis, alopecia, and autoimmune diseases such as rheumatoid arthritis.

Another method relates to inhibiting GSK-3, Aurora, or CDK-2 activity in a biological sample, which method comprises contacting the biological sample with the GSK-3 or Aurora inhibitor of formula VIII, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3, Aurora or CDK-2.

Each of the aforementioned methods directed to the inhibition of GSK-3, Aurora or CDK-2, or the treatment of a disease alleviated thereby, is preferably carried out with a preferred compound of formula VIII, as described above.

The above formula I compounds contain a pyrazole ring bearing the  $R^2$  and  $R^{2'}$  substituents. In their search for further inhibitors of the protein kinases GSK and Aurora, applicants sought to replace the pyrazole moiety of formula I with other heteroaromatic rings. One of the more effective pyrazole ring replacements was found to be a triazole ring. Inhibitors having this triazole ring are otherwise structurally similar to the formula I compounds and are represented by the general formula IX:



IX

or a pharmaceutically acceptable derivative or prodrug  
 5 thereof, wherein:

$Z^1$  is nitrogen or  $CR^9$  and  $Z^2$  is nitrogen or CH, provided  
 that at least one of  $Z^1$  and  $Z^2$  is nitrogen;

G is Ring C or Ring D;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl,  
 10 pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring,  
 wherein said Ring C has one or two ortho substituents  
 independently selected from  $-R^1$ , any substitutable non-  
 ortho carbon position on Ring C is independently  
 substituted by  $-R^5$ , and two adjacent substituents on  
 15 Ring C are optionally taken together with their  
 intervening atoms to form a fused, unsaturated or  
 partially unsaturated, 5-6 membered ring having 0-3  
 heteroatoms selected from oxygen, sulfur or nitrogen,  
 said fused ring being optionally substituted by halo,  
 20 oxo, or  $-R^8$ ;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered  
 bicyclic ring selected from aryl, heteroaryl,  
 heterocyclyl or carbocyclyl, said heteroaryl or  
 heterocyclyl ring having 1-4 ring heteroatoms selected  
 25 from nitrogen, oxygen or sulfur, wherein Ring D is  
 substituted at any substitutable ring carbon by oxo or  
 $-R^5$ , and at any substitutable ring nitrogen by  $-R^4$ ,  
 provided that when Ring D is a six-membered aryl or

heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^1$  is selected from -halo, -CN,  $-NO_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $C_{1-6}$  aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or  $-R^8$ , said  $C_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

$R^x$  and  $R^y$  are independently selected from T- $R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 0-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring formed by  $R^x$  and  $R^y$  is substituted by oxo or T- $R^3$ , and any substitutable nitrogen on said ring formed by  $R^x$  and  $R^y$  is substituted by  $R^4$ ;

T is a valence bond or a  $C_{1-4}$  alkylidene chain;

$R^2$  is -R or -T-W- $R^6$ ;

$R^3$  is selected from -R, -halo, -OR,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-COCH_2COR$ ,  $-NO_2$ , -CN,  $-S(O)R$ ,  $-S(O)_2R$ , -SR,  $-N(R^4)_2$ ,  $-CON(R^7)_2$ ,  $-SO_2N(R^7)_2$ ,  $-OC(=O)R$ ,  $-N(R^7)COR$ ,  $-N(R^7)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^7)CON(R^7)_2$ ,  $-N(R^7)SO_2N(R^7)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^7)_2$ ;

each R is independently selected from hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-\text{COR}^7$ ,  
 $-\text{CO}_2$ (optionally substituted  $\text{C}_{1-6}$  aliphatic),  $-\text{CON}(\text{R}^7)_2$ ,  
 or  $-\text{SO}_2\text{R}^7$ , or two  $R^4$  on the same nitrogen are taken  
 together to form a 5-8 membered heterocyclyl or  
 5 heteroaryl ring;

each  $R^5$  is independently selected from  $-\text{R}$ , halo,  $-\text{OR}$ ,  
 $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{SO}_2\text{R}$ ,  $-\text{SR}$ ,  
 $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^4)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^4)\text{COR}$ ,  
 $-\text{N}(\text{R}^4)\text{CO}_2$ (optionally substituted  $\text{C}_{1-6}$  aliphatic),  
 10  $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$ ,  
 $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$ , or  $R^5$  and  
 an adjacent substituent taken together with their  
 intervening atoms form said ring fused to Ring C;  
 V is  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2-$ ,  $-\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 15  $-\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{N}(\text{R}^6)\text{CO}-$ ,  $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  
 $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  
 20  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ , or  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ;

W is  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  
 $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$ ,  $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$ ,  
 25  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ , or  $-\text{CON}(\text{R}^6)-$ ;

each  $R^6$  is independently selected from hydrogen, an  
 optionally substituted  $\text{C}_{1-4}$  aliphatic group, or two  $R^6$   
 30 groups on the same nitrogen atom are taken together  
 with the nitrogen atom to form a 5-6 membered  
 heterocyclyl or heteroaryl ring;

each  $R^7$  is independently selected from hydrogen or an  
 optionally substituted  $\text{C}_{1-6}$  aliphatic group, or two  $R^7$

on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring;

each R<sup>8</sup> is independently selected from an optionally


5 substituted C<sub>1-4</sub> aliphatic group, -OR<sup>6</sup>, -SR<sup>6</sup>, -COR<sup>6</sup>, -SO<sub>2</sub>R<sup>6</sup>, -N(R<sup>6</sup>)<sub>2</sub>, -N(R<sup>6</sup>)N(R<sup>6</sup>)<sub>2</sub>, -CN, -NO<sub>2</sub>, -CON(R<sup>6</sup>)<sub>2</sub>, or -CO<sub>2</sub>R<sup>6</sup>; and

R<sup>9</sup> is selected from -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR,


10 -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub> (optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>.

Compounds of formula IX may exist in


15 alternative tautomeric forms, as in tautomers 1-3 shown below. Unless otherwise indicated, the representation of any of these tautomers is meant to include the other two.




The R<sup>x</sup> and R<sup>y</sup> groups of formula IX may be taken together to form a fused ring, providing a bicyclic ring system containing Ring A. Preferred R<sup>x</sup>/R<sup>y</sup> rings include a  
 25 5-, 6-, 7-, or 8-membered unsaturated or partially unsaturated ring having 0-2 heteroatoms, wherein said R<sup>x</sup>/R<sup>y</sup> ring is optionally substituted. Examples of Ring A systems are shown below by compounds IX-A through IX-DD, wherein Z<sup>1</sup> is nitrogen or C(R<sup>9</sup>) and Z<sup>2</sup> is nitrogen or  
 30 C(H).




IX-A




IX-B




IX-C




IX-D




IX-E




IX-F




IX-G




IX-H




IX-I




IX-J




IX-K




IX-L




IX-M




IX-N




IX-O




IX-P




IX-Q




IX-R




IX-S




IX-T




IX-U




IX-V




IX-W




IX-X




IX-Y




IX-Z




IX-AA



IX-BB



IX-CC



IX-DD

Preferred bicyclic Ring A systems of formula IX include IX-A, IX-B, IX-C, IX-D, IX-E, IX-F, IX-G, IX-H, IX-I, IX-J, IX-K, IX-L, and IX-M, more preferably IX-A,

IX-B, IX-C, IX-F, and IX-H, and most preferably IX-A, IX-B, and IX-H.

In the monocyclic Ring A system of formula IX, preferred  $R^x$  groups include hydrogen, alkyl- or  
5 dialkylamino, acetamido, or a  $C_{1-4}$  aliphatic group such as methyl, ethyl, cyclopropyl, isopropyl or t-butyl.  
Preferred  $R^y$  groups, when present, include  $T-R^3$  wherein T is a valence bond or a methylene, and  $R^3$  is  $-R$ ,  $-N(R^4)_2$ , or  $-OR$ . Examples of preferred  $R^y$  include 2-pyridyl, 4-  
10 pyridyl, piperidinyl, methyl, ethyl, cyclopropyl, isopropyl, t-butyl, alkyl- or dialkylamino, acetamido, optionally substituted phenyl such as phenyl or halo-substituted phenyl, and methoxymethyl.


In the bicyclic Ring A system of formula IX,  
15 the ring formed by  $R^x$  and  $R^y$  taken together may be substituted or unsubstituted. Suitable substituents include  $-R$ , halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$   
20 aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ , wherein R and  $R^4$  are as defined above.  
Preferred  $R^x/R^y$  ring substituents include -halo,  $-R$ ,  $-OR$ ,  $-COR$ ,  $-CO_2R$ ,  $-CON(R^4)_2$ ,  $-CN$ , or  $-N(R^4)_2$  wherein R is an  
25 optionally substituted  $C_{1-6}$  aliphatic group.

Preferred  $R^2$  groups of formula IX include hydrogen,  $C_{1-4}$  aliphatic, alkoxycarbonyl, (un)substituted phenyl, hydroxyalkyl, alkoxyalkyl, aminocarbonyl, mono- or dialkylaminocarbonyl, aminoalkyl, alkylaminoalkyl,  
30 dialkylaminoalkyl, phenylaminocarbonyl, and (N-heterocyclyl)carbonyl. Examples of such preferred  $R^2$  substituents include methyl, cyclopropyl, ethyl, isopropyl, propyl, t-butyl, cyclopentyl, phenyl,  $CO_2H$ ,  $CO_2CH_3$ ,  $CH_2OH$ ,  $CH_2OCH_3$ ,  $CH_2CH_2CH_2OH$ ,  $CH_2CH_2CH_2OCH_3$ ,



CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>Ph, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NHCOOC(CH<sub>3</sub>)<sub>3</sub>,  
 CONHCH(CH<sub>3</sub>)<sub>2</sub>, CONHCH<sub>2</sub>CH=CH<sub>2</sub>, CONHCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, CONHCH<sub>2</sub>Ph,  
 CONH(cyclohexyl), CON(Et)<sub>2</sub>, CON(CH<sub>3</sub>)CH<sub>2</sub>Ph, CONH(n-C<sub>3</sub>H<sub>7</sub>),  
 CON(Et)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, CONHCH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, CON(n-C<sub>3</sub>H<sub>7</sub>)<sub>2</sub>, CO(3-  
 5 methoxymethylpyrrolidin-1-yl), CONH(3-tolyl), CONH(4-  
 tolyl), CONHCH<sub>3</sub>, CO(morpholin-1-yl), CO(4-methylpiperazin-  
 1-yl), CONHCH<sub>2</sub>CH<sub>2</sub>OH, CONH<sub>2</sub>, and CO(piperidin-1-yl). A  
 more preferred R<sup>2</sup> group for formula IX compounds is  
 hydrogen.

10 An embodiment that is particularly useful for  
 treating GSK3-mediated diseases relates to compounds of  
 formula X wherein ring A is a pyrimidine ring:



15

X

or a pharmaceutically acceptable derivative or prodrug  
 thereof, wherein;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl,  
 pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring,  
 wherein said Ring C has one or two ortho substituents  
 20 independently selected from -R<sup>1</sup>, any substitutable non-  
 ortho carbon position on Ring C is independently  
 substituted by -R<sup>5</sup>, and two adjacent substituents on  
 Ring C are optionally taken together with their  
 intervening atoms to form a fused, unsaturated or  
 25 partially unsaturated, 5-6 membered ring having 0-3  
 heteroatoms selected from oxygen, sulfur or nitrogen,

said fused ring being optionally substituted by halo, oxo, or  $-R^8$ ;

$R^1$  is selected from -halo, -CN,  $-\text{NO}_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $\text{C}_{1-6}$  aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or  $-R^8$ , said  $\text{C}_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

$R^x$  and  $R^y$  are independently selected from T- $R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 0-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring formed by  $R^x$  and  $R^y$  is substituted by oxo or T- $R^3$ , and any substitutable nitrogen on said ring formed by  $R^x$  and  $R^y$  is substituted by  $R^4$ ;

T is a valence bond or a  $\text{C}_{1-4}$  alkylidene chain;

$R^2$  is -R or -T-W- $R^6$ ;

$R^3$  is selected from -R, -halo, -OR,  $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{COCH}_2\text{COR}$ ,  $-\text{NO}_2$ , -CN,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{S}(\text{O})_2\text{R}$ , -SR,  $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^7)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^7)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^7)\text{COR}$ ,  $-\text{N}(\text{R}^7)\text{CO}_2$  (optionally substituted  $\text{C}_{1-6}$  aliphatic),  $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^7)\text{CON}(\text{R}^7)_2$ ,  $-\text{N}(\text{R}^7)\text{SO}_2\text{N}(\text{R}^7)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^7)_2$ ;

each R is independently selected from hydrogen or an optionally substituted group selected from  $\text{C}_{1-6}$  aliphatic,  $\text{C}_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-\text{COR}^7$ ,  
 $-\text{CO}_2$ (optionally substituted  $\text{C}_{1-6}$  aliphatic),  $-\text{CON}(\text{R}^7)_2$ ,  
 or  $-\text{SO}_2\text{R}^7$ , or two  $\text{R}^4$  on the same nitrogen are taken  
 together to form a 5-8 membered heterocyclyl or  
 5 heteroaryl ring;

each  $\text{R}^5$  is independently selected from  $-\text{R}$ , halo,  $-\text{OR}$ ,  
 $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{SO}_2\text{R}$ ,  $-\text{SR}$ ,  
 $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^4)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^4)\text{COR}$ ,  
 $-\text{N}(\text{R}^4)\text{CO}_2$ (optionally substituted  $\text{C}_{1-6}$  aliphatic),  
 10  $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$ ,  
 $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$ , or  $\text{R}^5$  and  
 an adjacent substituent taken together with their  
 intervening atoms form said ring fused to Ring C;

V is  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2-$ ,  $-\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 15  $-\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{N}(\text{R}^6)\text{CO}-$ ,  $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  
 $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  
 20  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ , or  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ;

W is  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  
 $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$ ,  $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$ ,  
 25  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ , or  $-\text{CON}(\text{R}^6)-$ ;

each  $\text{R}^6$  is independently selected from hydrogen, an  
 optionally substituted  $\text{C}_{1-4}$  aliphatic group, or two  $\text{R}^6$   
 30 groups on the same nitrogen atom are taken together  
 with the nitrogen atom to form a 5-6 membered  
 heterocyclyl or heteroaryl ring;

each  $\text{R}^7$  is independently selected from hydrogen or an  
 optionally substituted  $\text{C}_{1-6}$  aliphatic group, or two  $\text{R}^7$

on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring; and

each  $R^8$  is independently selected from an optionally substituted  $C_{1-4}$  aliphatic group,  $-OR^6$ ,  $-SR^6$ ,  $-COR^6$ ,  $-SO_2R^6$ ,  $-N(R^6)_2$ ,  $-N(R^6)N(R^6)_2$ ,  $-CN$ ,  $-NO_2$ ,  $-CON(R^6)_2$ , or  $-CO_2R^6$ .

Compounds of formula X are structurally similar to compounds of formula II except for the replacement of the pyrazole ring moiety by the triazole ring moiety. Preferred  $R^2$ ,  $R^x$ ,  $R^y$  and Ring C groups of formula X are as described above for the formula II compounds. Preferred formula X compounds have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a naphthyl, quinolinyl or isoquinolinyl ring;

(b)  $R^x$  is hydrogen or  $C_{1-4}$  aliphatic and  $R^y$  is  $T-R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted 5-7 membered unsaturated or partially unsaturated ring having 0-2 ring nitrogens;

(c)  $R^1$  is -halo, an optionally substituted  $C_{1-6}$  aliphatic group, phenyl,  $-COR^6$ ,  $-OR^6$ ,  $-CN$ ,  $-SO_2R^6$ ,  $-SO_2NH_2$ ,  $-N(R^6)_2$ ,  $-CO_2R^6$ ,  $-CONH_2$ ,  $-NHCOR^6$ ,  $-OC(O)NH_2$ , or  $-NHCO_2R^6$ ; and

(d)  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group.

More preferred compounds of formula **X** have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring;

(b)  $R^x$  is hydrogen or methyl and  $R^y$  is  $-R$ ,  $N(R^4)_2$ , or  $-OR$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a benzo ring or a 5-7 membered carbocyclo ring, wherein said ring formed by  $R^x$  and  $R^y$  is optionally substituted with  $-R$ , halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;

(c)  $R^1$  is  $-halo$ , a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or  $-CN$ ;

(d)  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl or a  $C_{1-6}$  aliphatic group; and

(e) each  $R^5$  is independently selected from  $-halo$ ,  $-CN$ ,  $-NO_2$ ,  $-N(R^4)_2$ , optionally substituted  $C_{1-6}$  aliphatic group,  $-OR$ ,  $-C(O)R$ ,  $-CO_2R$ ,  $-CONH(R^4)$ ,  $-N(R^4)COR$ ,  $-SO_2N(R^4)_2$ , or  $-N(R^4)SO_2R$ .

Even more preferred compounds of formula **X** have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring;


(b)  $R^x$  is hydrogen or methyl and  $R^y$  is methyl, methoxymethyl, ethyl, cyclopropyl, isopropyl, t-butyl, alkyl- or an optionally substituted group selected from 2-pyridyl, 4-pyridyl, piperidinyl, or phenyl, or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted benzo ring or a 6-membered carbocyclo ring;

(c)  $R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN;

(d)  $R^2$  is hydrogen or a  $C_{1-6}$  aliphatic group; and

(e) each  $R^5$  is independently selected from -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH( $C_{1-4}$  aliphatic), -N( $C_{1-4}$  aliphatic)<sub>2</sub>, -O( $C_{1-4}$  aliphatic),  $C_{1-4}$  aliphatic, and -CO<sub>2</sub>( $C_{1-4}$  aliphatic).

Another embodiment of this invention relates to compounds of formula XI:



XI

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or -R<sup>5</sup>, and at any substitutable ring nitrogen by -R<sup>4</sup>,

provided that when Ring D is a six-membered aryl or heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^x$  and  $R^y$  are taken together with their intervening atoms to form a fused benzo ring or 5-8 membered carbocyclic ring, wherein any substitutable carbon on said fused ring formed by  $R^x$  and  $R^y$  is substituted by oxo or  $T-R^3$ ;

T is a valence bond or a  $C_{1-4}$  alkylidene chain;

$R^2$  is  $-R$  or  $-T-W-R^6$ ;

- 5  $R^3$  is selected from  $-R$ ,  $-\text{halo}$ ,  $=O$ ,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-COCH_2COR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-S(O)_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  
10  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;

each R is independently selected from hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each  $R^4$  is independently selected from  $-R^7$ ,  $-COR^7$ ,  $-CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-CON(R^7)_2$ , or  $-SO_2R^7$ , or two  $R^4$  on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or  
15 heteroaryl ring;

- each  $R^5$  is independently selected from  $-R$ ,  $\text{halo}$ ,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  
20  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;

V is  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-N(R^6)SO_2-$ ,  $-SO_2N(R^6)-$ ,  $-N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  $-N(R^6)CO-$ ,  $-N(R^6)C(O)O-$ ,  $-N(R^6)CON(R^6)-$ ,  $-N(R^6)SO_2N(R^6)-$ ,  $-N(R^6)N(R^6)-$ ,

$-C(O)N(R^6)-$ ,  $-OC(O)N(R^6)-$ ,  $-C(R^6)_2O-$ ,  $-C(R^6)_2S-$ ,  
 $-C(R^6)_2SO-$ ,  $-C(R^6)_2SO_2-$ ,  $-C(R^6)_2SO_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)-$ ,  
 $-C(R^6)_2N(R^6)C(O)-$ ,  $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  
 $-C(R^6)=N-O-$ ,  $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ , or  
 5  $-C(R^6)_2N(R^6)CON(R^6)-$ ;

W is  $-C(R^6)_2O-$ ,  $-C(R^6)_2S-$ ,  $-C(R^6)_2SO-$ ,  $-C(R^6)_2SO_2-$ ,  
 $-C(R^6)_2SO_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  
 $-C(R^6)OC(O)-$ ,  $-C(R^6)OC(O)N(R^6)-$ ,  $-C(R^6)_2N(R^6)CO-$ ,  
 $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  $-C(R^6)=N-O-$ ,  
 10  $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ ,  
 $-C(R^6)_2N(R^6)CON(R^6)-$ , or  $-CON(R^6)-$ ;

each  $R^6$  is independently selected from hydrogen or an  
 optionally substituted  $C_{1-4}$  aliphatic group, or two  $R^6$   
 groups on the same nitrogen atom are taken together  
 15 with the nitrogen atom to form a 5-6 membered  
 heterocyclyl or heteroaryl ring; and

each  $R^7$  is independently selected from hydrogen or an  
 optionally substituted  $C_{1-6}$  aliphatic group, or two  $R^7$   
 on the same nitrogen are taken together with the  
 20 nitrogen to form a 5-8 membered heterocyclyl or  
 heteroaryl ring.

Compounds of formula XI are structurally  
 similar to compounds of formula III except for the  
 replacement of the pyrazole ring moiety by the triazole  
 25 ring moiety. Preferred  $R^2$ ,  $R^x$ ,  $R^y$ , and Ring D groups of  
 formula XI are as described above for the formula III  
 compounds. Preferred formula XI compounds have one or  
 more, and more preferably all, of the features selected  
 from the group consisting of:

30 (a) Ring D is an optionally substituted ring  
 selected from a phenyl, pyridinyl, piperidinyl,  
 piperazinyl, pyrrolidinyl, thienyl, azepanyl,  
 morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-  
 tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-



dihydro-1*H*-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted benzo ring or 5-7 membered carbocyclo ring; and

(c)  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group.

More preferred compounds of formula XI have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1*H*-isoindolyl, 2,3-dihydro-1*H*-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a benzo ring or 5-7 membered carbocyclo ring, wherein said ring formed by  $R^x$  and  $R^y$  is optionally substituted with -R, oxo, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted  $C_{1-6}$  aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>;


(c)  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl or a  $C_{1-6}$  aliphatic group; and

(d) each  $R^5$  is independently selected from halo, oxo, CN, NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, -SR, -OR, -C(O)R, or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl,  $C_{6-10}$  aryl, or  $C_{1-6}$  aliphatic.

Even more preferred compounds of formula XI have one or more, and more preferably all, of the features selected from the group consisting of:

- (a)  $R^x$  and  $R^y$  are taken together with their  
 5 intervening atoms to form a benzo ring or 6-membered carbocyclo ring, wherein said ring formed by  $R^x$  and  $R^y$  is optionally substituted with halo, CN, oxo,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, ( $C_{1-6}$  alkyl)carbonyl, ( $C_{1-6}$  alkyl)sulfonyl, mono- or dialkylamino, mono- or dialkylaminocarbonyl, mono- or  
 10 dialkylaminocarbonyloxy, or 5-6 membered heteroaryl;  
 (b) each  $R^5$  is independently selected from -halo, -CN, -oxo, -SR, -OR,  $-N(R^4)_2$ ,  $-C(O)R$ , or a substituted or unsubstituted group selected from 5-6 membered heterocyclyl,  $C_{6-10}$  aryl, or  $C_{1-6}$  aliphatic; and  
 15 (c)  $R^2$  is hydrogen or a  $C_{1-6}$  aliphatic group.

Another embodiment of this invention relates to compounds of formula XII:



XII

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

- Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or  
 25 heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or

-R<sup>5</sup>, and at any substitutable ring nitrogen by -R<sup>4</sup>, provided that when Ring D is a six-membered aryl or heteroaryl ring, -R<sup>5</sup> is hydrogen at each ortho carbon position of Ring D;

R<sup>x</sup> and R<sup>y</sup> are independently selected from T-R<sup>3</sup>, or R<sup>x</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 1-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring is optionally and independently substituted by T-R<sup>3</sup>, and any substitutable nitrogen on said ring is substituted by R<sup>4</sup>;

T is a valence bond or a C<sub>1-4</sub> alkylidene chain;

5 R<sup>2</sup> is -R or -T-W-R<sup>6</sup>;

R<sup>3</sup> is selected from -R, -halo, =O, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -COCH<sub>2</sub>COR, -NO<sub>2</sub>, -CN, -S(O)R, -S(O)<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>;

each R is independently selected from hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each R<sup>4</sup> is independently selected from -R<sup>7</sup>, -COR<sup>7</sup>, -CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -CON(R<sup>7</sup>)<sub>2</sub>, or -SO<sub>2</sub>R<sup>7</sup>, or two R<sup>4</sup> on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or  
10 heteroaryl ring;

each R<sup>5</sup> is independently selected from -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR,

$-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  
 $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  
 $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;  
V is  $-O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-N(R^6)SO_2-$ ,  $-SO_2N(R^6)-$ ,  
5  $-N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  $-N(R^6)CO-$ ,  $-N(R^6)C(O)O-$ ,  
 $-N(R^6)CON(R^6)-$ ,  $-N(R^6)SO_2N(R^6)-$ ,  $-N(R^6)N(R^6)-$ ,  
 $-C(O)N(R^6)-$ ,  $-OC(O)N(R^6)-$ ,  $-C(R^6)_2O-$ ,  $-C(R^6)_2S-$ ,  
 $-C(R^6)_2SO-$ ,  $-C(R^6)_2SO_2-$ ,  $-C(R^6)_2SO_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)-$ ,  
 $-C(R^6)_2N(R^6)C(O)-$ ,  $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  
10  $-C(R^6)=N-O-$ ,  $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ , or  
 $-C(R^6)_2N(R^6)CON(R^6)-$ ;  
W is  $-C(R^6)_2O-$ ,  $-C(R^6)_2S-$ ,  $-C(R^6)_2SO-$ ,  $-C(R^6)_2SO_2-$ ,  
 $-C(R^6)_2SO_2N(R^6)-$ ,  $-C(R^6)_2N(R^6)-$ ,  $-CO-$ ,  $-CO_2-$ ,  
 $-C(R^6)OC(O)-$ ,  $-C(R^6)OC(O)N(R^6)-$ ,  $-C(R^6)_2N(R^6)CO-$ ,  
15  $-C(R^6)_2N(R^6)C(O)O-$ ,  $-C(R^6)=NN(R^6)-$ ,  $-C(R^6)=N-O-$ ,  
 $-C(R^6)_2N(R^6)N(R^6)-$ ,  $-C(R^6)_2N(R^6)SO_2N(R^6)-$ ,  
 $-C(R^6)_2N(R^6)CON(R^6)-$ , or  $-CON(R^6)-$ ;

each  $R^6$  is independently selected from hydrogen or an  
optionally substituted  $C_{1-4}$  aliphatic group, or two  $R^6$   
20 groups on the same nitrogen atom are taken together  
with the nitrogen atom to form a 5-6 membered  
heterocyclyl or heteroaryl ring; and

each  $R^7$  is independently selected from hydrogen or an  
optionally substituted  $C_{1-6}$  aliphatic group, or two  $R^7$   
25 on the same nitrogen are taken together with the  
nitrogen to form a 5-8 membered heterocyclyl ring or  
heteroaryl.

Compounds of formula XII are structurally  
similar to compounds of formula IV except for the  
30 replacement of the pyrazole ring moiety by the triazole  
ring moiety. Preferred  $R^2$ ,  $R^x$ ,  $R^y$ , and Ring D groups of  
formula XII are as described above for the formula IV  
compounds. Preferred formula XII compounds have one or

more, and more preferably all, of the features selected from the group consisting of:

(a) Ring D is an optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^x$  is hydrogen or  $C_{1-4}$  aliphatic and  $R^y$  is T- $R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted 5-7 membered unsaturated or partially unsaturated ring having 1-2 ring heteroatoms; and

(c)  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group.

More preferred compounds of formula XII have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^x$  is hydrogen or methyl and  $R^y$  is -R,  $N(R^4)_2$ , or -OR, or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a 5-7 membered unsaturated or partially unsaturated ring having 1-2 ring nitrogens, wherein said ring is optionally substituted with -R, halo, oxo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted  $C_{1-6}$  aliphatic),

$-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  
 $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ ;

(c)  $R^2$  is hydrogen or a substituted or  
unsubstituted group selected from aryl or a  $C_{1-6}$  aliphatic  
5 group; and

(d) each  $R^5$  is independently selected from halo,  
oxo, CN,  $NO_2$ ,  $-N(R^4)_2$ ,  $-CO_2R$ ,  $-CONH(R^4)$ ,  $-N(R^4)COR$ ,  
 $-SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ ,  $-SR$ ,  $-OR$ ,  $-C(O)R$ , or a substituted  
or unsubstituted group selected from 5-6 membered  
10 heterocyclyl,  $C_{6-10}$  aryl, or  $C_{1-6}$  aliphatic.


Even more preferred compounds of formula XII  
have one or more, and more preferably all, of the  
features selected from the group consisting of:

(a)  $R^x$  and  $R^y$  are taken together with their  
15 intervening atoms to form a 6-membered unsaturated or  
partially unsaturated ring having 1-2 ring nitrogens,  
optionally substituted with halo, CN, oxo,  $C_{1-6}$  alkyl,  $C_{1-6}$   
alkoxy, ( $C_{1-6}$  alkyl)carbonyl, ( $C_{1-6}$  alkyl)sulfonyl, mono- or  
dialkylamino, mono- or dialkylaminocarbonyl, mono- or  
20 dialkylaminocarbonyloxy, or 5-6 membered heteroaryl;

(b) each  $R^5$  is independently selected from  
-halo, -CN, -oxo, -SR, -OR,  $-N(R^4)_2$ ,  $-C(O)R$ , or a  
substituted or unsubstituted group selected from 5-6  
membered heterocyclyl,  $C_{6-10}$  aryl, or  $C_{1-6}$  aliphatic; and

(c)  $R^2$  is hydrogen or a  $C_{1-6}$  aliphatic group.

Another embodiment of this invention relates to  
compounds of formula XIII:



XIII

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

$Z^1$  is nitrogen,  $CR^a$ , or CH, and  $Z^2$  is nitrogen or CH; provided that one of  $Z^1$  and  $Z^2$  is nitrogen;

G is Ring C or Ring D;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from  $-R^1$ , any substitutable non-ortho carbon position on Ring C is independently substituted by  $-R^5$ , and two adjacent substituents on Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo, oxo, or  $-R^8$ ;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is substituted at any substitutable ring carbon by oxo or  $-R^5$ , and at any substitutable ring nitrogen by  $-R^4$ , provided that when Ring D is a six-membered aryl or

heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^1$  is selected from -halo, -CN,  $-NO_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $C_{1-6}$  aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or  $-R^8$ , said  $C_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

$R^x$  and  $R^y$  are independently selected from T- $R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-8 membered ring having 0-3 ring heteroatoms selected from oxygen, sulfur, or nitrogen, wherein any substitutable carbon on said fused ring formed by  $R^x$  and  $R^y$  is substituted by oxo or T- $R^3$ , and any substitutable nitrogen on said ring formed by  $R^x$  and  $R^y$  is substituted by  $R^4$ ;

T is a valence bond or a  $C_{1-4}$  alkylidene chain;

$R^2$  is -R or -T-W- $R^6$ ;

$R^3$  is selected from -R, -halo, -OR,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-COCH_2COR$ ,  $-NO_2$ , -CN,  $-S(O)R$ ,  $-S(O)_2R$ , -SR,  $-N(R^4)_2$ ,  $-CON(R^7)_2$ ,  $-SO_2N(R^7)_2$ ,  $-OC(=O)R$ ,  $-N(R^7)COR$ ,  $-N(R^7)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^7)CON(R^7)_2$ ,  $-N(R^7)SO_2N(R^7)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^7)_2$ ;

each R is independently selected from hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;



each  $R^4$  is independently selected from  $-R^7$ ,  $-\text{COR}^7$ ,  
 $-\text{CO}_2$ (optionally substituted  $\text{C}_{1-6}$  aliphatic),  $-\text{CON}(\text{R}^7)_2$ ,  
 or  $-\text{SO}_2\text{R}^7$ , or two  $R^4$  on the same nitrogen are taken  
 together to form a 5-8 membered heterocyclyl or  
 heteroaryl ring;

each  $R^5$  is independently selected from  $-\text{R}$ , halo,  $-\text{OR}$ ,  
 $-\text{C}(=\text{O})\text{R}$ ,  $-\text{CO}_2\text{R}$ ,  $-\text{COCOR}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{S}(\text{O})\text{R}$ ,  $-\text{SO}_2\text{R}$ ,  $-\text{SR}$ ,  
 $-\text{N}(\text{R}^4)_2$ ,  $-\text{CON}(\text{R}^4)_2$ ,  $-\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{OC}(=\text{O})\text{R}$ ,  $-\text{N}(\text{R}^4)\text{COR}$ ,  
 $-\text{N}(\text{R}^4)\text{CO}_2$ (optionally substituted  $\text{C}_{1-6}$  aliphatic),  
 $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$ ,  $-\text{C}=\text{NN}(\text{R}^4)_2$ ,  $-\text{C}=\text{N}-\text{OR}$ ,  $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$ ,  
 $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$ ,  $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$ , or  $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$ , or  $R^5$  and  
 an adjacent substituent taken together with their  
 intervening atoms form said ring fused to Ring C;

V is  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{SO}-$ ,  $-\text{SO}_2-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2-$ ,  $-\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 $-\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{N}(\text{R}^6)\text{CO}-$ ,  $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  
 $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ , or  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ ;

W is  $-\text{C}(\text{R}^6)_2\text{O}-$ ,  $-\text{C}(\text{R}^6)_2\text{S}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}-$ ,  $-\text{C}(\text{R}^6)_2\text{SO}_2-$ ,  
 $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$ ,  $-\text{CO}-$ ,  $-\text{CO}_2-$ ,  
 $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$ ,  $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$ ,  $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$ ,  $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$ ,  
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$ , or  $-\text{CON}(\text{R}^6)-$ ;

each  $R^6$  is independently selected from hydrogen, an  
 optionally substituted  $\text{C}_{1-4}$  aliphatic group, or two  $R^6$   
 groups on the same nitrogen atom are taken together  
 with the nitrogen atom to form a 5-6 membered  
 heterocyclyl or heteroaryl ring;


each  $R^7$  is independently selected from hydrogen or an  
 optionally substituted  $\text{C}_{1-6}$  aliphatic group, or two  $R^7$

on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring;

each  $R^6$  is independently selected from an optionally substituted  $C_{1-4}$  aliphatic group,  $-OR^6$ ,  $-SR^6$ ,  $-COR^6$ ,  $-SO_2R^6$ ,  $-N(R^6)_2$ ,  $-N(R^6)N(R^6)_2$ ,  $-CN$ ,  $-NO_2$ ,  $-CON(R^6)_2$ , or  $-CO_2R^6$ ; and

$R^a$  is selected from halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ ,  $-OC(=O)N(R^4)_2$ , or an optionally substituted group selected from  $C_{1-6}$  aliphatic,  $C_{6-10}$  aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms.

Compounds of formula XIII may be represented by specifying  $Z^1$  and  $Z^2$  as shown below:



Compounds of formula XIII are structurally similar to compounds of formula V except for the replacement of the pyrazole ring moiety by the triazole ring moiety. Preferred  $R^2$ ,  $R^x$ ,  $R^y$ ,  $R^a$ , and Ring G groups of formula XIII are as described above for the formula V compounds. Preferred formula XIII compounds have one or

more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a naphthyl, quinolinyl or isoquinolinyl ring, and  $R^1$  is -halo, an optionally substituted  $C_{1-6}$  aliphatic group, phenyl,  $-COR^6$ ,  $-OR^6$ ,  $-CN$ ,  $-SO_2R^6$ ,  $-SO_2NH_2$ ,  $-N(R^6)_2$ ,  $-CO_2R^6$ ,  $-CONH_2$ ,  $-NHCOR^6$ ,  $-OC(O)NH_2$ , or  $-NHSO_2R^6$ ; or Ring D is an optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^x$  is hydrogen or  $C_{1-4}$  aliphatic and  $R^y$  is  $T-R^3$ , or  $R^x$  and  $R^y$  are taken together with their intervening atoms to form an optionally substituted 5-7 membered unsaturated or partially unsaturated ring having 0-2 ring nitrogens; and

(c)  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group.

More preferred compounds of formula XIII have one or more, and more preferably all, of the features selected from the group consisting of:

(a) Ring C is a phenyl or pyridinyl ring, optionally substituted by  $-R^5$ , wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or  $-CN$ ; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl,

piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl,  
1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-  
tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-  
dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or  
5 naphthyl;

(b)  $R^x$  is hydrogen or methyl and  $R^y$  is -R,  
 $N(R^4)_2$ , or -OR, or  $R^x$  and  $R^y$  are taken together with their  
intervening atoms to form a benzo ring or a 5-7 membered  
carbocyclo ring, wherein said ring formed by  $R^x$  and  $R^y$  is  
10 optionally substituted with -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R,  
-COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>,  
-SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally  
substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>,  
-C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or  
15 -OC(=O)N(R<sup>4</sup>)<sub>2</sub>;

(c)  $R^2$  is hydrogen or a substituted or  
unsubstituted group selected from aryl, or a C<sub>1-6</sub>  
aliphatic group; and

(d) each  $R^5$  is independently selected from  
20 -halo, -CN, -NO<sub>2</sub>, -N(R<sup>4</sup>)<sub>2</sub>, optionally substituted C<sub>1-6</sub>  
aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH(R<sup>4</sup>), -N(R<sup>4</sup>)COR,  
-SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, or -N(R<sup>4</sup>)SO<sub>2</sub>R, and, when Ring G is Ring D, Ring  
D is substituted by oxo or  $R^5$ .

Even more preferred compounds of formula XIII  
25 have one or more, and more preferably all, of the  
features selected from the group consisting of:

(a)  $R^x$  is hydrogen or methyl and  $R^y$  is methyl,  
methoxymethyl, ethyl, cyclopropyl, isopropyl, t-butyl,  
alkyl- or an optionally substituted group selected from  
30 2-pyridyl, 4-pyridyl, piperidinyl, or phenyl, or  $R^x$  and  $R^y$   
are taken together with their intervening atoms to form a  
benzo ring or a 6-membered carbocyclo ring wherein said  
ring formed by  $R^x$  and  $R^y$  is optionally substituted with  
halo, CN, oxo, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, (C<sub>1-6</sub> alkyl)carbonyl,

(C<sub>1-6</sub> alkyl)sulfonyl, mono- or dialkylamino, mono- or dialkylaminocarbonyl, mono- or dialkylaminocarbonyloxy, or 5-6 membered heteroaryl;


(b) Ring C is a phenyl or pyridinyl ring, optionally substituted by -R<sup>5</sup>, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and R<sup>1</sup> is -halo, a C<sub>1-4</sub> aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or naphthyl;

(c) R<sup>2</sup> is hydrogen or a C<sub>1-6</sub> aliphatic group; and


(d) each R<sup>5</sup> is independently selected from -Cl, -F, -CN, -CF<sub>3</sub>, -NH<sub>2</sub>, -NH(C<sub>1-4</sub> aliphatic), -N(C<sub>1-4</sub> aliphatic)<sub>2</sub>, -O(C<sub>1-4</sub> aliphatic), C<sub>1-4</sub> aliphatic, and -CO<sub>2</sub>(C<sub>1-4</sub> aliphatic), and when Ring G is Ring D, Ring D is substituted by oxo or R<sup>5</sup>.

Representative compounds of formula IX are shown below in Table 8.

Table 8.




IX-1



IX-2




IX-3




IX-4




IX-5




IX-6




IX-7




IX-8




IX-9




IX-10




IX-11




IX-12




IX-13




IX-14




IX-15




IX-16




IX-17




IX-18



IX-19




IX-20




IX-21



IX-22




IX-23




IX-24



IX-25




IX-26




IX-27




IX-28




IX-29




IX-30



IX-31




IX-32




IX-33




IX-34




IX-35




IX-36




IX-37




IX-38




IX-39



IX-40




IX-41




IX-42







IX-43




IX-44




IX-45



IX-46




IX-47




IX-48


5




IX-49




IX-50




IX-51



IX-52




IX-53




IX-54


10




IX-55




IX-56




IX-57



IX-58




IX-59




IX-60




IX-61




IX-62




IX-63




IX-64




IX-65




IX-66




IX-67




IX-68




IX-69




IX-70




IX-71




IX-72




IX-73




IX-74




IX-75




IX-76




IX-77




IX-78




IX-79




IX-80




IX-81




IX-82




IX-83




IX-84




IX-85




IX-86




IX-87




IX-88




IX-89




IX-90




IX-91




IX-92




IX-93




IX-94




IX-95




IX-96




IX-97




IX-98




IX-99




IX-100




IX-101




IX-102




IX-103




IX-104




IX-105




IX-106




IX-107




IX-108




IX-109




IX-110



IX-111




IX-112




IX-113




IX-114




IX-115




IX-116




IX-117




IX-118




IX-119




IX-120




IX-121




IX-122




IX-123



IX-124




IX-125




IX-126


5




IX-127




IX-128



IX-129



IX-130




IX-131




IX-132


10




IX-133




IX-134




IX-135




IX-136



IX-137




IX-138




IX-139




IX-140




IX-141




IX-142




IX-143




IX-144



IX-145




IX-146




IX-147







IX-148




IX-149




IX-150




IX-151



IX-152




IX-153




IX-154




IX-155




IX-156




IX-157




IX-158




IX-159




IX-160




IX-161




IX-162




IX-163




IX-164




IX-165




IX-166




IX-167




IX-168




IX-169




IX-170




IX-171




IX-172




IX-173




IX-174



IX-175



IX-176



IX-177

In another embodiment, this invention provides a composition comprising a compound of formula IX and a pharmaceutically acceptable carrier.

5           One aspect of this invention relates to a method of inhibiting GSK-3 activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula IX.

10           Another aspect relates to a method of treating a disease that is alleviated by treatment with a GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition  
15 comprising a compound of formula IX.

          Another aspect relates to a method of enhancing glycogen synthesis and/or lowering blood levels of glucose in a patient in need thereof, comprising administering to said patient a therapeutically effective  
20 amount of a composition comprising a compound of formula IX. This method is especially useful for diabetic patients.

          Another aspect relates to a method of inhibiting the production of hyperphosphorylated Tau  
25 protein in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition comprising a compound of formula IX. This method is especially useful in halting or slowing the progression of Alzheimer's disease.

30           Another aspect relates to a method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, comprising administering to said patient a therapeutically effective amount of a composition

comprising a compound of formula IX. This method is especially useful for treating schizophrenia.

One aspect of this invention relates to a method of inhibiting Aurora activity in a patient, comprising administering to the patient a therapeutically effective amount of a composition comprising a compound of formula IX.

Another aspect relates to a method of treating a disease that is alleviated by treatment with an Aurora inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of a composition comprising a compound of formula IX. This method is especially useful for treating cancer, such as colon, ovarian, and breast cancer.

Another method relates to inhibiting GSK-3 or Aurora activity in a biological sample, which method comprises contacting the biological sample with the GSK-3 or Aurora inhibitor of formula IX, or a pharmaceutical composition thereof, in an amount effective to inhibit GSK-3 or Aurora.

Each of the aforementioned compositions and methods directed to the inhibition of GSK-3 or Aurora, or the treatment of a disease alleviated thereby, is preferably carried out with a preferred compound of formula IX, as described above.


The compounds of this invention may be prepared as illustrated by the Synthetic Methods below, by the Synthetic Examples described herein and by general methods known to those skilled in the art.

#### General Synthetic Methods

The general synthetic methods below provide a series of general reaction routes that were used to

prepare compounds of this invention. Methods A-F below are particularly useful for preparing formula II compounds. In most cases, Ring C is drawn as a phenyl ring bearing an ortho R<sup>1</sup> substituent. However, it will be apparent to one skilled in the art that compounds having other Ring C groups may be obtained in a similar manner. Methods analogous to methods A-F are also useful for preparing other compounds of this invention. Methods F-I below are particularly useful for preparing compounds of formula III or IV.

#### Method A



Method A is a general route for the preparation of compounds wherein ring C is an aryl or heteroaryl ring. Preparation of the starting dichloropyrimidine 1 may be achieved in a manner similar to that described in *Chem. Pharm. Bull.*, 30, 9, 1982, 3121-3124. The chlorine in position 4 of intermediate 1 may be replaced by an aminopyrazole or aminoindazole to provide intermediate 2 in a manner similar to that described in *J. Med. Chem.*, 38, 3547-3557 (1995). Ring C is then introduced using a boronic ester under palladium catalysis (see *Tetrahedron*, 48, 37, 1992, 8117-8126). This method is illustrated by the following procedure.


A suspension of 1H-quinazoline-2,4-dione (10.0 g, 61.7 mmol) in POCl<sub>3</sub> (60 mL, 644 mmol) and N,N-dimethylaniline (8mL, 63.1 mmol) is heated under reflux


for 2 h. Excess POCl<sub>3</sub> is evaporated under vacuum, the residue is poured into ice, and the precipitate is collected by filtration. The crude solid 2,4-dichloroquinazoline product may be used without further purification.

To a solution of 2,4-dichloro-quinazoline (3.3 g, 16.6 mmol) in anhydrous ethanol (150 mL) is added 5-methyl-1H-pyrazol-3-yl amine (3.2 g, 32.9 mmol). The mixture is stirred at room temperature for 4 h, and the resulting precipitate is collected by filtration, washed with ethanol, and dried under vacuum to afford (2-chloro-quinazolin-4-yl)-(5-methyl-1H-pyrazol-3-yl)-amine.

To a solution of (2-chloro-quinazolin-4-yl)-(5-methyl-1H-pyrazol-3-yl)-amine (50 mg, 0.19 mmol) in DMF (1.0 mL) is added the desired arylboronic acid (0.38 mmol), 2M Na<sub>2</sub>CO<sub>3</sub> (0.96 mmol), and tri-*t*-butylphosphine (0.19 mmol). Under nitrogen, PdCl<sub>2</sub>(dppf) (0.011 mmol) is added in one portion. The reaction mixture is then heated at 80°C for 5 to 10 hours, cooled to room temperature, and poured into water (2 mL). The resulting precipitate is collected by filtration, washed with water, and purified by HPLC.

#### Method B





Methods B through F describe routes where the  
 5 pyrazole ring system is introduced after Ring C and the  
 pyrimidine ring portion are first constructed. A  
 versatile intermediate is the 4-chloropyrimidine 4, which  
 is readily obtained from pyrimidinone 3 as shown in  
 Method B(i). This reaction sequence is generally  
 10 applicable for a variety of Ring C groups including  
 aliphatic, aryl, heteroaryl, or heterocyclyl. See *J.*  
*Med. Chem.*, 38, 3547-3557 (1995).

For quinazoline ring systems (where R<sup>x</sup> and R<sup>y</sup>  
 are taken together to form a benzo ring), the useful  
 15 intermediate 6 may be obtained by condensing an  
 anthranilic acid or its derivative with a benzamidine as  
 shown in Method B(ii) or by condensing a benzoylchloride  
 with an anthranilamide as shown in Method B(iii). Many  
 substituted anthranilic acid, anthranilamide, benzamidine  
 20 and benzoylchloride starting materials may be obtained by  
 known methods. See *Aust. J. Chem.*, 38, 467-474 and *J.*  
*Med. Chem.*, 38, 3547-3557 (1995). Method B(iii) is  
 illustrated by the following procedure.


To a solution of anthranilamide (33 mmol) in  
 25 THF and CH<sub>2</sub>Cl<sub>2</sub> (1:1, 70 mL) is added the desired  
 benzoylchloride (33 mmol), and triethylamine (99 mmol) at  
 room temperature. The mixture is stirred for about 14  
 hours. The resulting precipitate is collected by  
 filtration, washed with CH<sub>2</sub>Cl<sub>2</sub> and water, and dried under  
 30 vacuum. The crude 2-benzoylaminoanthranilamide may be used  
 directly for the next step without further purification.

To a solution of the above crude product (13 mmol) in ethanol (50 mL) is added NaOEt (26 mmol) at room temperature. The mixture is heated under reflux for 48 to 96 h. The solvent is evaporated and the residue is  
5 neutralized using concentrated HCl to pH 7. The product is then collected by filtration and dried under vacuum to provide 2-phenyl-3H-quinazolin-4-one that may be used without further purification.


To a suspension of the above product (12 mmol)  
10 in POCl<sub>3</sub> (120 mmol) is added tri-*n*-propylamine (24 mmol). The mixture is heated under reflux for 1h. After removal of the excess POCl<sub>3</sub> by evaporation, the residue is dissolved in ethyl acetate, and washed with 1N NaOH (twice) and water (twice). The organic layer is dried  
15 over MgSO<sub>4</sub>, the solvent is evaporated under vacuum, and the crude product is purified by flash chromatography (eluting with 10% of ethyl acetate in hexanes) to give 4-chloro-2-aryl quinazoline.

To a solution of 4-chloro-2-aryl quinazoline  
20 (0.16 mmol) in DMF (or THF, ethanol) (1 mL) is added the desired aminopyrazole or aminoindazole (0.32 mmol). The mixture is heated in DMF (or THF under reflux) at 100 to 110°C for 16 h (or in ethanol at 130-160°C for 16 hours) and then poured into water (2 mL). The precipitate is  
25 collected by filtration and purified by HPLC.

#### Method C






Method D(i)

Methods C and D(i) above employ  $\beta$ -ketoesters 8 and 10, respectively, as pyrimidinone precursors. The substitution pattern of the  $R^x$  and  $R^y$  groups on the pyrimidinone ring will be reversed if a chlorocrotonate 11 (Synth. Comm, (1986), 997-1002), instead of the corresponding  $\beta$ -ketoester 10, is condensed with the desired benzamidine. These methods are illustrated by the following general procedure.

To a solution of a  $\beta$ -ketoester (5.2 mmol) and amidinium chloride (5.7 mmol) in ethanol (5 mL) is added sodium ethoxide (7.8 mmol). The mixture is heated under reflux for 7-14 hours. After evaporation the resulting residue is dissolved in water, acidified with concentrated HCl to pH 6, and then filtered to obtain a solid product 2-aryl-3H-pyrimidin-4-one (yield 75-87%), which may be purified by flash column chromatography if needed. To this pyrimidinone (3.7 mmol) is added  $POCl_3$  (4 mL) and  $n-Pr_3N$  (1.4 mL). The mixture is heated under reflux for 1 hour. After evaporation of the excess  $POCl_3$ , the residue is dissolved in ethyl acetate, washed with 1N NaOH solution (three times) and  $NaHCO_3$  (once), and dried over  $MgSO_4$ . The solvent is removed under vacuum and the residue is purified by flash column chromatography eluting with 10% of ethyl acetate in hexanes to give 2-aryl-4-chloro-pyrimidine as a pale yellow syrup. This crude product may be treated with a 3-aminopyrazole or 3-aminoindazole as described above.

Method D(ii)

5                    Method D(ii) above shows a general route for the preparation of the present compounds, such as compound 40, wherein R<sup>y</sup> is N(R<sup>4</sup>)<sub>2</sub>. See *Il Farmaco*, 52(1) 61-65 (1997). Displacement of the 6-chloro group is exemplified here using morpholine. This method is  
10 illustrated by the following procedure.


To a solution of 2-methylmalonic acid diethyl ester (5 mmol) and sodium ethoxide (15 mmol) is added the appropriate amidine salt (5 mmol) in ethanol (10 mL) and the reaction heated at reflux for 2-24 hours. The  
15 residue is dissolved in water and acidified with 2N HCl. The resulting precipitate is filtered off and further purified by flash chromatography (yield 5-35%) to afford the pyrimidinedione 37. To 37 (1.6 mmol) is added POCl<sub>3</sub> (32 mmol) and tri-n-propylamine (6.4 mmol) and the  
20 reaction refluxed is for 1h. After evaporation of excess POCl<sub>3</sub>, the residue is dissolved in ethyl acetate, basified with 1N NaOH, separated and the aqueous phase twice more extracted with ethyl acetate. The combined organics are dried (sodium sulfate) and evaporated. Purification by

flash chromatography provides the dichloropyrimidine (38) as a yellow oil in 23% yield.

A solution of 38 (0.33 mmol) in methanol (5 mL) is treated with an amine, exemplified here using morpholine (0.64 mmol) and refluxed 1 hour. After evaporation of solvent, the residue is purified by flash chromatography to provide the mono-chloropyrimidine 39 as a colorless oil in 75% yield.

The mono-chloropyrimidine, 39, (0.19 mmol) may be treated with a 3-aminopyrazole or 3-aminoindazole compound in a manner substantially similar those described above in Methods A and B.

#### Method E




As shown by Method E, an acyl isocyanate 12 may be condensed with an enamine to provide pyrimidinone 9 (J. Org. Chem (1993), 58, 414-418; J. Med. Chem., (1992), 35, 1515-1520; J. Org. Chem., 1997, 32, 313-214). This method is illustrated by the following general procedure.

The enamine is prepared according to W. White, et al, J. Org Chem. (1967), 32, 213-214. The acyl isocyanate is prepared according to G Bradley, et al, J Med. Chem. (1992), 35, 1515-1520. The coupling reaction then follows the procedure of S Kawamura, et al, J. Org. Chem, (1993), 58, 414-418. To the enamine (10 mmol) in tetrahydrofuran (30 mL) at 0°C under nitrogen is added dropwise over 5 min a solution of acyl isocyanate (10 mmol) in tetrahydrofuran (5 mL). After stirring for 0.5 h, acetic acid (30 mL) is added, followed by ammonium

acetate (50 mmol). The mixture is refluxed for 2 h with continuous removal of tetrahydrofuran. The reaction is cooled to room temperature and is poured into water (100 mL). The precipitate is filtered, washed with water and ether and dried to provide the 2-aryl-3H-pyrimidin-4-one.

#### Method F



Method F shows a general route for the preparation of the present compounds wherein  $\text{R}^x$  and  $\text{R}^y$  are taken together to form a 5-8 membered partially unsaturated saturated or unsaturated ring having 1-3 heteroatoms. The condensation of a 2-amino-carboxylic acid, such as 2-amino-nicotinic acid 13, and an acid chloride 7 provides an oxazinone 14. Treatment of 14 with ammonium hydroxide will furnish the benzamide 15 which may be cyclized to a 2-(substituted)-pyrido[2,3-d][1,3]pyrimidin-4-one 16. This method is illustrated by the following procedure.

2-(Trifluoromethyl)benzoyl chloride (4.2 ml, 29.2 mmol) is added dropwise to a solution of 2-aminonicotinic acid (2.04g, 14.76 mmol) in 20 ml of pyridine. The reaction mixture is heated at 158 C for 30 min then cooled to room temperature. The reaction is poured into 200 ml of water and an oil forms which solidifies upon stirring. The solid is collected by vacuum filtration and washed with water and diethyl ether. The product is dried to give 2-(2-trifluoromethyl-phenyl)-pyrido[2,3-d][1,3]oxazin-4-one

(2.56 g, 60% yield) which may be used in the next step without further purification.

2-(2-Trifluoromethyl-phenyl)-pyrido[2,3-d][1,3]oxazin-4-one (2.51g) is stirred in 30% ammonium hydroxide (25 ml) at room temperature overnight. The resulting precipitate is filtered and rinsed with water and diethyl ether. The precipitate is dried under vacuum at 50 C overnight to give 2-(2-trifluoromethyl-benzoylamino)-nicotinamide (850 mg, 33% yield)

2-(2-Trifluoromethyl-benzoylamino)-nicotinamide (800mg, 2.6mmol) is dissolved in 10ml of ethanol. Potassium ethoxide (435mg, 5.2mmol) is added to the solution which is heated to reflux for 16 h. The reaction mixture is evaporated in vacuo to afford a gummy residue that is dissolved in water and acidified with 10% sodium hydrogen sulfate to pH 7. The resulting precipitate is filtered and dried under vacuum at 50 C to give 2-(2-trifluoromethyl-phenyl)-3H-pyrido[2,3-d]pyrimidin-4-one.

#### Method G

Method G is analogous to Method B(i) above. This method is illustrated by the following general procedure.


2-(3,4-Dichloro-phenyl)-3H-quinazolin-4-one (1g, 3.43 mmol) is suspended in phosphorus oxychloride (4 mL) and the reaction mixture was stirred at 110°C for 3 hours. The solvents are then evaporated and the residue is treated carefully with an ice cold aqueous saturated solution of NaHCO<sub>3</sub>. The solid is collected by filtration and washed with ether to give 4-chloro-2-(3,5-dichloro-phenyl)-quinazoline as a white solid (993 mg, 93%).

To 4-chloro-2-(3,5-dichloro-phenyl)-quinazoline (400mg, 1.29 mmol) in THF (30 mL) is added 3-amino-5-

methyl pyrazole (396 mg, 2.58 mmol) and the reaction mixture is heated at 65°C overnight. The solvents are then evaporated and the residue triturated with ethyl acetate, filtered and washed with a minimum amount of ethanol to give [2-(3,4-dichlorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine as a white solid (311 mg 65%): mp 274°C; <sup>1</sup>H NMR (DMSO) δ 2.34 (3H, s), 6.69 (1H, s), 7.60 (1H, m), 7.84 (1H, d), 7.96 (2H, d), 8.39 (1H, dd), 8.60 (1H, d), 8.65 (1H, d), 10.51 (1H, s), 12.30 (1H, s); IR (solid) 1619, 1600, 1559, 1528, 1476, 1449, 1376, 1352, 797, 764, 738; MS 370.5 (M+H)<sup>+</sup>.

The THF solvent used in the previous step may be replaced by other organic solvents such as ethanol, N,N-dimethylformamide, or dioxane.

#### Method H



Method H shows routes in which a Ring D aryl group bearing a halogen (X is Br or I) may be converted to other formula III compounds. Method H(i) shows a phenylboronic acid coupling to Ring D to provide compound 18 and Method H(ii) shows an acetylene coupling to provide compound 19. Substituent X in compound 17 may be

bromine or iodine. These methods are illustrated by the following procedures.


Method H(i). To a mixture of [2-(4-bromophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (196 mg, 0.51 mmol) and phenylboronic acid (75 mg, 0.62 mmol) in THF/water (1/1, 4 mL) is added Na<sub>2</sub>CO<sub>3</sub> (219 mg, 2.06 mmol), triphenylphosphine (9mg, 1/15 mol%) and palladium acetate (1 mg, 1/135 mol%). The mixture is heated at 80°C overnight, the solvents are evaporated and the residue is purified by flash chromatography (gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH) to give (2-biphenyl-4-yl-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine as a yellow solid (99 mg, 51%): <sup>1</sup>H NMR (DMSO) δ 2.37 (3H, s), 6.82 (1H, s), 7.39-7.57 (4H, m), 7.73-7.87 (6H, m), 8.57 (2H, d), 8.67 (1H, d), 10.42 (1H, s), 12.27 (1H, s); MS 378.2 (M+H)<sup>+</sup>

Method H(ii). To a mixture of [2-(4-bromophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (114 mg, 0.3 mmol), and trimethylsilylacetylene (147 mg, 1.5 mmol) in DMF (2 mL) is added CuI (1.1 mg, 1/50 mol%), Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (4.2 mg, 1/50 mol%) and triethylamine (121 mg, 0.36 mmol). The mixture is heated at 120°C overnight and the solvent is evaporated. The residue is triturated in ethyl acetate and the precipitate is collected by filtration.

To the above precipitate suspended in THF (3 mL) is added tetrabutylammonium fluoride (1M in THF, 1.1eq). The reaction mixture is stirred at room temperature for two hours and the solvent is evaporated. The residue is purified by flash chromatography (gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH) to give [2-(4-ethynylphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine as a white solid (68 mg, 70%): <sup>1</sup>H NMR (DMSO) δ 2.34 (3H, s), 4.36 (1H, s), 6.74 (1H, s), 7.55 (1H, m), 7.65 (2H, d), 7.84 (2H, m), 8.47

(2H, d), 8.65 (1H, d), 10.43 (1H, s), 12.24 (1H, s); MS  
326.1 (M+H)<sup>+</sup>


### Method I



Method I above shows a general route for the preparation of the present compounds wherein ring D is a heteroaryl or heterocyclyl ring directly attached to the pyrimidine 2-position via a nitrogen atom. Displacement of the 2-chloro group, exemplified here using piperidine, may be carried out in a manner similar to that described in *J. Med. Chem.*, 38, 2763-2773 (1995) and *J. Chem. Soc.*, 1766-1771 (1948). This method is illustrated by the following procedure.


To a solution of (2-chloro-quinazolin-4-yl)-(1H-indazol-3-yl)-amine (1 equivalent, 0.1-0.2 mmol) in N, N-dimethylacetamide (1 ml) is added the desired amine (3 equivalents). The resulting mixture is maintained at 100°C for 6 h and then purified by reverse-phase HPLC.

### Method J



25






Method J above shows the preparation of  
 5 compounds of formula V via the displacement of a chloro group from an appropriately substituted pyridyl ring. Method J(i) is a route for preparing compounds of formula Va (see *Indian J. Chem. Sect. B*, 35, 8, 1996, 871-873). Method J(ii) is a route for preparing compounds of  
 10 formula Vb (see *Bioorg. Med. Chem.*, 6, 12, 1998, 2449-2458). For convenience, the chloropyridines 21 and 23 are shown with a phenyl substituent corresponding to Ring D of formula V. It would be apparent to one skilled in the art that Method J is also useful for preparing  
 15 compounds of formula V wherein Ring D is heteroaryl, heterocyclyl, carbocyclyl or other aryl rings. Method J is illustrated by the following procedures.

Method J(i). (5-Methyl-2H-pyrazol-3-yl)-(2-phenyl-quinolin-4-yl)-amine. To 4-chloro-2-  
 20 phenylquinoline (*J. Het. Chem.*, 20, 1983, 121-128) (0.53g, 2.21 mmol) in diphenylether (5 mL) was added 3-amino-5-methylpyrazole (0.43g, 4.42 mmol) and the mixture was heated at 200°C overnight with stirring. To the cooled mixture was added petroleum ether (20 mL) and the  
 25 resulting crude precipitate was filtered and further washed with petroleum ether. The crude solid was purified by flash chromatography (SiO<sub>2</sub>, gradient DCM-MeOH) to give the title compound as a white solid: mp 242-244°C; <sup>1</sup>H NMR (DMSO) δ 2.27(3H, s), 6.02(1H, s), 7.47(2H, d), 7.53-  
 30 7.40(2H, br m), 7.67(1H, m), 7.92(1H, m), 8.09(2H, d),

8.48 (2H, m), 9.20 (1H, s), 12.17 (1H, br s); IR (solid) 1584, 1559, 1554, 1483, 1447, 1430, 1389; MS 301.2 (M+H)<sup>+</sup>

Method J(ii). (5-Methyl-2H-pyrazol-3-yl)-(3-phenyl-isoquinolin-1-yl)-amine. To 1-chloro-3-phenylisoquinoline (J. Het. Chem., 20, 1983, 121-128) (0.33g, 1.37 mmol) in dry DMF (5 mL) was added 3-amino-5-methylpyrazole (0.27g, 2.74 mmol) and potassium carbonate (0.57g, 4.13 mmol) and the mixture was heated under reflux for 6 hours. The mixture was cooled and the bulk of DMF was evaporated. The residue was extracted twice with ethyl acetate and the combined organic layers were washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The crude was purified by flash chromatography (SiO<sub>2</sub>, gradient DCM-MeOH) to give the title compound as a colourless oil; <sup>1</sup>H NMR (MeOD) δ 2.23 (3H, s), 5.61 (1H, s), 7.41 (1H, m), 7.52 (2H, m), 7.62 (1H, m), 7.81 (1H, m), 8.07 (1H, d), 8.19 (2H, m), 8.29 (1H, s), 8.54 (1H, d); MS 301.2 (M+H)<sup>+</sup>

## 20 Method K




Method K shows a route for the preparation of compounds of formula VI. A versatile starting material is 2,4,6-trichloro-[1,3,5]triazine 25 in which the chlorine substituents may be sequentially displaced. The displacement of one of the chlorines by an aryl Grignard reagent or an aryl boronic acid is described in PCT patent application WO 01/25220 and *Helv. Chim. Acta*, 33, 1365 (1950). The displacement of one of the chlorines by a heteroaryl ring is described in WO 01/25220; *J. Het.*

*Chem.*, 11, 417 (1974); and *Tetrahedron* 31, 1879 (1975).

These reactions provide a 2,4-dichloro-(6-substituted) [1,3,5]triazine 26 that is a useful intermediate for the preparation of compounds of formula VI. Alternatively, intermediate 26 may be obtained by constructing the triazine ring by known methods. See US patent 2,832,779; and US patent 2,691,020 together with *J. Am. Chem. Soc.* 60, 1656 (1938). In turn, one of the chlorines of 26 may be displaced as described above to provide 2-chloro-(4,6-disubstituted) [1,3,5]triazine 27. The treatment of 27 with an appropriate aminopyrazole provides the desired compound of formula VI.

#### Method L







Method L shows a route for preparing compounds of formula VII. For illustration purposes the trifluoromethylchalcone 28 is used as a starting material; however, it would be apparent to one skilled in the art that other rings may be used in place of the

trifluoromethylphenyl and phenyl rings of compound 28. Substituted chalcones may be prepared by known methods, for example as described in the *Indian J. Chemistry*, 32B, 449 (1993). Condensation of a chalcone with urea provides the pyrimidinone 29, which may be treated with POCl<sub>3</sub> to give the chloropyrimidine 30. See *J. Chem. Eng. Data*, 30(4) 512 (1985) and *Egypt. J. Chem.*, 37(3), 283 (1994). In an alternative approach to compound 30, one of the aryl rings attached to the pyrimidine is introduced by displacement of the 4-chloro group of 2,4-dichloro-(6-aryl)-pyrimidine by an aryl boronic acid using a palladium catalyst such as (Ph<sub>3</sub>P)<sub>4</sub>Pd in the presence of a base such as sodium carbonate as described in *Bioorg. Med. Lett.*, 9(7), 1057 (1999). Displacement of the chlorine of compound 30 by an appropriate aminopyrazole provides compounds of this invention, such as 31. The last step of this method is illustrated by the following procedure.

[4-(4-Methylpiperidin-1-yl)-pyrimidin-2-yl]-(5-methyl-2H-pyrazol-3-yl)-amine. To a solution of 2-chloro-4-(4-methylpiperidin-1-yl)-pyrimidine (prepared using a procedure similar to the one reported in *Eur. J. Med. Chem.*, 26(7) 729(1991)) (222 mg, 1.05 mmol) in BuOH (5 mL) was added 3-amino-5-methyl-2H-pyrazole (305mg, 3.15 mmol) and the reaction mixture was then heated under reflux overnight. The solvent was evaporated and the residue dissolved in a mixture ethanol/water (1/3, 4 mL). Potassium carbonate (57mg, 0.41 mmol) was added and the mixture was stirred at room temperature for 2 hours. The resulting suspension was filtered, washed with water twice and rinsed with ether twice to give the title compound as a white solid (143mg, 50%): mp 193-195°C; <sup>1</sup>H NMR (DMSO) δ 0.91 (3H, d), 1.04 (2H, m), 1.67 (3H, m), 2.16 (3H, s), 2.83 (2H, t), 4.31 (2H, m), 6.19 (2H, m),

7.87 (1H, d), 8.80 (1H, br s), 11.71 (1H, s); IR (solid) 1627, 1579, 1541, 1498, 1417, 1388, 1322, 1246; MS 273.3 (M+H)<sup>+</sup>.

## 5 Method M




Method M provides routes for obtaining compounds of formula VIII. A general procedure for displacing the chlorine of a 4-chloro-6-substituted-pyridazine, 32, with an appropriately substituted

pyrazole to provide VIIIA is described in *J. Het. Chem.*, 20, 1473 (1983). Analogous reactions may be carried out as follows: (a) with 3-chloro-5-substituted-pyridazine, 33, to provide VIIIB is described in *J. Med. Chem.*, 41(3), 311 (1998); (b) with 5-chloro-3-substituted-[1,2,4]triazine, 34, to provide VIIIC is described in *Heterocycles*, 26(12), 3259 (1987); and (c) with 3-chloro-5-substituted-[1,2,4]triazine, 35, to provide IIID is described in *Pol. J. Chem.*, 57, 7, (1983); *Indian J. Chem. Sect. B*, 26, 496 (1987); and *Agric. Biol. Chem.*, 54(12), 3367 (1990). An alternative procedure to compounds of formula VIIIC is described in *Indian J. Chem. Sect. B*, 29(5), 435 (1990).

Compounds of formula IX are prepared by methods substantially similar to those described above for the pyrazole-containing compounds of formula I. Methods A-J may be used to prepare the triazole-containing compounds of formula IX by replacing the amino-pyrazole compound with an amino-triazole compound. Such methods are specifically exemplified by Synthetic Examples 415-422 set forth below. The amino-triazole intermediate may be obtained by methods described in *J. Org. Chem. USSR*, 27, 952-957 (1991).


Certain synthetic intermediates that are useful for preparing the protein kinase inhibitors of this invention are new. Accordingly, another aspect of this invention relates to a 3-aminoindazole compound of formula A:




A

where R<sup>10</sup> is one to three substituents that are each independently selected from fluoro, bromo, C<sub>1-6</sub> haloalkyl,


nitro, or 1-pyrrolyl. Examples of such compounds include the following:




A1



A2




A3




A4


5




A5




A6




A7



A8




A9



A10

10


Another aspect of this invention relates to a 4-chloropyrimidine compound of formula B:




B

15 wherein  $R^x$  and  $R^y$  are as defined above;  $R^1$  is selected from Cl, F,  $CF_3$ , CN, or  $NO_2$ ; and is one to three substituents that are each independently selected from H, Cl, F,  $CF_3$ ,  $NO_2$ , or CN; provided that  $R^1$  and  $R^5$  are not simultaneously Cl. Examples of compounds of formula B


20 are shown below:




B1




B2




B3



B4




B5




B6


5




B7




B8




B9



B10




B11




B12


10




B13



B14




B15



B16




B17




B18

15




B19



B20




Another aspect of this invention relates to compounds of formula C:



C


5

wherein  $R^x$ ,  $R^y$ ,  $R^2$ , and  $R^{2'}$  are as defined above. Examples of compounds of formula C are shown below:




C1


10




C2



C3




C4



C5




C6




C7


15




C8



C9




C10



C11




C12



C13




C14



C15

5


Yet another aspect of this invention relates to compounds of formula D:




D

10


where  $R^5$ ,  $R^x$  and  $R^y$  are as defined above. Examples of formula D compounds and other useful pyrimidinone intermediates are shown below:




D1




D2




D3



D4




D5




D6


20




D7




D8




D9




D10




D11




D12




D13




D14




D15




D16



D17



D18



D20

In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

SYNTHETIC EXAMPLES

The following HPLC methods were used in the analysis of the compounds as specified in the Synthetic Examples set forth below. As used herein, the term "R<sub>t</sub>" refers to the retention time observed for the compound using the HPLC method specified.

## HPLC-Method A:

10 Column: C18, 3 um, 2.1 X 50 mm, "Lighting" by Jones Chromatography.  
Gradient: 100% water (containing 1% acetonitrile, 0.1% TFA) to 100% acetonitrile (containing 0.1% TFA) over 4.0 min, hold at 100% acetonitrile for 1.4 min  
15 and return to initial conditions. Total run time 7.0 min. Flow rate: 0.8 mL/min.

## HPLC-Method B:

20 Column: C18, 5 um, 4.6 X 150 mm "Dynamax" by Rainin  
Gradient: 100% water (containing 1% acetonitrile, 0.1% TFA) to 100% acetonitrile (containing 0.1% TFA) over 20 min, hold at 100% acetonitrile for 7.0 min and return to initial conditions. Total run time 31.5 min. Flow rate: 1.0 mL/min.

25

## HPLC-Method C:

Column: Cyano, 5 um, 4.6 X 150 mm "Microsorb" by Varian.  
Gradient: 99% water (0.1% TFA), 1% acetonitrile  
30 (containing 0.1% TFA) to 50% water (0.1% TFA), 50% acetonitrile (containing 0.1% TFA) over 20 min, hold for 8.0 min and return to initial conditions. Total run time 30 min. Flow rate: 1.0 mL/min.

## HPLC-Method D:

Column: Waters (YMC) ODS-AQ 2.0x50mm, S5, 120A.

Gradient: 90% water (0.2% Formic acid), 10%  
acetonitrile (containing 0.1% Formic acid) to 10%  
5 water (0.1% formic acid), 90% acetonitrile  
(containing 0.1% formic acid) over 5.0 min, hold for  
0.8 min and return to initial conditions. Total run  
time 7.0 min.

Flow rate: 1.0 mL/min.

10

## HPLC-Method E:

Column: 50x2.0mm Hypersil C18 BDS; 5  $\mu$ m

Gradient: elution 100% water (0.1% TFA), to 5% water  
(0.1% TFA), 95% acetonitrile (containing 0.1% TFA)  
15 over 2.1 min, returning to initial conditions after  
2.3 min.

Flow rate: 1 mL/min.

Example 1 [2-(2-Clorophenyl)-5,6-dimethylpyrimidin-4-yl]-

20 (5-Methyl-2H-pyrazol-3-yl)-amine (II-1):  $^1\text{H}$ NMR (500 MHz,  
DMSO-d<sub>6</sub>)  $\delta$  10.4 (s, br, 1H), 7.74 (m, 2H), 7.68 (m, 1H),  
7.60 (m, 1H), 6.39 (s, 1H), 2.52 (s, 3H), 2.30 (s, 3H),  
2.22 (s, 3H); MS 314.1 (M+H).

25 Example 2 [2-(2-Chloro-phenyl)-6,7,8,9-tetrahydro-5H-  
cycloheptapyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-2):

Prepared in 30% yield.  $^1\text{H}$ NMR (500MHz, DMSO-d<sub>6</sub>)  $\delta$  1.72 (m,  
4H), 1.91 (m, 2H), 3.02 (m, 4H), 7.05 (t, 1H), 7.33 (t,  
1H), 7.39 (m, 1H), 7.47 (d, 1H), 7.55 (m, 3H), 7.59 (d,  
30 1H), 10.4 (m, 1H), 13.11 (br. s, 1H); EI-MS 390.2 (M+H);  
HPLC-Method A,  $R_t$  2.99 min.

Example 3 (5-Fluoro-1H-indazol-3-yl)-[2-(2-  
trifluoromethyl-phenyl)-5,6,7,8-tetrahydro-pyrido[3,4-

d]pyrimidin-4-yl]-amine (II-3): Compound II-18 (90 mg, 0.17 mmol) was treated with an equal weight of Pd/C (10%) in 4.4% formic acid in MeOH at room temperature for 14 h. The mixture was filtered through celite, the filtrate was evaporated, and crude product was purified by HPLC to provide 18 mg (24%) of the desired product as pale yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1H), 9.51 (s, 1H), 9.26 (s, 2H), 7.72 (d, 1H), 7.63 (t, 1H), 7.58 (t, 1H), 7.49 (m, 2H), 7.21 (td, 1H), 7.15 (dd, 1H), 4.24 (s, 2H), 3.56 (m, 2H), 2.95 (m, 2H) ppm. MS (ES<sup>+</sup>): m/e= 429.22 (M+H); HPLC-Method A, R<sub>t</sub> 2.88 min.

Example 4 [2-(2-Chloro-phenyl)-6,7,8,9-tetrahydro-5H-cycloheptapyrimidin-4-yl]-(7-fluoro-1H-indazol-3-yl)-amine (II-4): Prepared in 52% yield to afford a white solid. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 1.72 (m, 4H), 1.92 (m, 2H), 3.00 (m, 4H), 7.02 (td, 1H), 7.20 (dd, 1H), 7.40 (m, 1H), 7.42 (d, 1H), 7.52 (m, 3H), 10.5 (m, 1H), 13.50 (br. s, 1H); EI-MS 408.2 (M+H); HPLC-Method A, R<sub>t</sub> 3.00 min.

Example 5 [2-(2-Chloro-phenyl)-6,7,8,9-tetrahydro-5H-cycloheptapyrimidin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-5): Prepared in 51% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 1.71 (m, 4H), 1.91 (m, 2H), 3.01 (m, 4H), 7.24 (td, 1H), 7.41 (m, 2H), 7.54 (m, 4H), 10.5 (m, 1H), 13.1 (br. s, 1H); EI-MS 408.2 (M+H); HPLC-Method A, R<sub>t</sub> 3.05 min.

Example 6 [2-(2-Chloro-phenyl)-6,7,8,9-tetrahydro-5H-cycloheptapyrimidin-4-yl]-(5,7-difluoro-1H-indazol-3-yl)-amine (II-6): Prepared according to Method C in 72% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 1.72 (m, 4H), 1.91 (m, 2H), 3.01 (m, 4H), 7.31 (m, 2H), 7.41 (m, 1H), 7.54 (m, 3H), 10.5 (m, 1H), 13.6 (br. s, 1H); EI-MS 426.2 (M+H); HPLC-Method A, R<sub>t</sub> 3.21 min.

Example 7 (7-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydroquinazolin-4-yl]-amine (II-7): Prepared in 62% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.5 (s, br, 1H), 10.1 (s, br, 1H), 7.75 (m, 4H), 7.33 (d, 1H), 7.17 (dd, 1H), 7.00 (td, 1H), 2.80 (m, 2H), 2.71 (m, 2H), 1.89 (br, 4H) ppm; LC-MS (ES+) 428.44 (M+H), (ES-) 426.43 (M-H); HPLC-Method A, R<sub>t</sub> 3.02 min.

10 Example 8 (5-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydroquinazolin-4-yl]-amine (II-8): Prepared in 53% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.1 (s, 1H), 10.2 (s, br, 1H), 7.75 (m, 4H), 7.50 (dd, 1H), 7.27 (dd, 1H), 7.21 (td, 1H), 2.80 (m, 2H), 2.72 (m, 2H), 1.88 (m, 4H) ppm; MS (ES+) 428.43 (M+H), (ES-) 426.43 (M-H); HPLC-Method A, R<sub>t</sub> 3.01 min.

Example 9 (5,7-Difluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydroquinazolin-4-yl]-amine (II-9): Prepared in 37% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.7 (s, 1H), 10.2 (s, br, 1H), 7.80 (d, 1H), 7.76 (t, 1H), 7.69 (m, 2H), 7.31 (t, 1H), 7.18 (d, 1H), 2.81 (t, br, 2H), 2.72 (t, br, 2H), 1.90 (m, 4H) ppm; MS (ES+) 446.42 (M+H), (ES-) 444.37 (M-H); HPLC-Method A, R<sub>t</sub> 3.09 min.

Example 10 (5-Trifluoromethyl-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydroquinazolin-4-yl]-amine (II-10): Prepared by Method C in ethanol in 35% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.2 (s, 1H), 10.1 (s, br, 1H), 8.01 (s, 1H), 7.76 (d, 1H), 7.66 (m, 4H), 7.57 (d, 1H), 2.79 (m, 2H), 2.73 (m, 2H), 1.89 (m, 4H) ppm. MS (ES+) 478.45 (M+H), (ES-) 476.42 (M-H); HPLC-Method A, R<sub>t</sub> 3.21 min.

Example 11 (5,7-difluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-6,7,8,9-tetrahydro-5H-cycloheptapyrimidin-4-yl]-amine (II-11): Prepared in 60%  
5 yield. White solid. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 1.72 (m, 4H), 1.91 (m, 2H), 3.01 (m, 4H), 7.15 (dd, 1H), 7.30 (td, 1H), 7.66 (m, 2H), 7.72 (t, 1H), 7.78 (d, 1H), 10.2 (m, 1H), 13.5 (br. s, 1H); EI-MS 460.2 (M+H); HPLC-Method A, R<sub>t</sub> 3.13 min.

10

Example 12 (6-Benzyl-2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydro-pyrido[4,3-d]pyrimidin-4-yl)-(5-fluoro-1H-indazol-3-yl)-amine (II-12): Prepared in 49%  
15 yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.8 (s, 1H), 9.11 (s, 1H), 7.68 (d, 1H), 7.58 (t, 1H), 7.53 (t, 1H), 7.44 (m, 4H), 7.37 (t, 2H), 7.29 (t, 1H), 7.19 (m, 2H), 3.78 (s, 2H), 3.61 (s, 2H), 2.81 (s, br, 4H) ppm; LC-MS (ES+) 519.24 (M+H); HPLC-Method A, R<sub>t</sub> 3.11 min.

20 Example 13 (1H-Indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-6,7,8,9-tetrahydro-5H-cycloheptapyrimidin-4-yl]-amine (II-13): Prepared in 40% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 1.70 (m, 4H), 1.90 (m, 2H), 3.00 (m, 4H), 7.01 (t, 1H), 7.30 (td, 1H), 7.44 (d, 1H), 7.49 (d, 1H), 7.68  
25 (m, 3H), 7.77 (d, 1H), 10.01 (m, 1H), 12.83 (s, 1H); EI-MS 424.2 (M+H); HPLC-Method A, R<sub>t</sub> 3.17 min.

Example 14 (7-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-6,7,8,9-tetrahydro-5H-cycloheptapyrimidin-4-yl]-amine (II-14): Prepared in 78%  
30 yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 1.71 (m, 4H), 1.91 (m, 2H), 3.00 (m, 4H), 6.98 (td, 1H), 7.16 (dd, 1H), 7.31 (d, 1H), 7.68 (m, 3H), 7.77 (d, 1H), 10.25 (m, 1H), 13.40 (br. s, 1H); EI-MS 442.2 (M+H); HPLC-Method A, R<sub>t</sub> 3.12



min.

Example 15 (5-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-6,7,8,9-tetrahydro-5H-

5 cycloheptapyrimidin-4-yl]-amine (II-15): Prepared in 63% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 1.71 (m, 4H), 1.91 (m, 2H), 3.00 (m, 4H), 7.20 (td, 1H), 7.25 (dd, 1H), 7.49 (dd, 1H), 7.69 (br. t, 2H), 7.74 (m, 1H), 7.79 (d, 1H), 10.35 (m, 1H), 13.00 (br. s, 1H); EI-MS 442.2 (M+H);  
10 HPLC-Method A, R<sub>t</sub> 3.21 min.

Example 16 (5-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydro-pyrido[4,3-d]pyrimidin-4-yl]-amine (II-16): A solution of compound

15 II-12 (45mg, 0.087 mmol) in methanol (4.4% HCOOH) was treated with an equal weight of Pd/C (10%) at room temperature for 14 h. The mixture was filtered through celite, the filtrate evaporated, and the crude product was purified by preparative HPLC to provide 15 mg (41%)  
20 of the desired product as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1H), 9.52 (s, 1H), 9.32 (s, 2H, TFA-OH), 7.72 (d, 1H), 7.59 (m, 2H), 7.49 (m, 2H), 7.21 (m, 1H), 7.15 (m, 1H), 4.31 (s, 2H), 3.55 (s, 2H), 3.00 (m, 2H) ppm; LC-MS (ES+) 429.20 (M+H); HPLC-Method A, R<sub>t</sub> 2.79  
25 min.

Example 17 (1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydroquinazolin-4-yl]-amine (II-17):

Prepared in 58% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.0  
30 (s, 1H), 10.3 (s, br, 1H), 7.74 (m, 4H), 7.51 (d, 1H), 7.47 (d, 1H), 7.32 (t, 1H), 7.03 (t, 1H), 2.82 (m, 2H), 2.73 (m, 2H), 1.90 (m, 4H) ppm; LC-MS (ES+) 410.21 (M+H); HPLC-Method A, R<sub>t</sub> 2.99 min.

Example 18 (7-Benzyl-2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydro-pyrido[4,3-d]pyrimidin-4-yl)-(5-fluoro-1H-indazol-3-yl)-amine (II-18): Prepared from compound B11 in 92% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1H), 10.5 (s, br, 1H), 9.58 (s, 1H, TFA-OH), 7.71 (d, 1H), 7.52 (m, 9H), 7.19 (m, 2H), 4.57 (s, 2H), 4.20 (m, 2H), 3.70 (m, 2H), 3.00 (m, 2H) ppm; LC-MS (ES+) 519.23 (M+H); HPLC-Method A, R<sub>t</sub> 3.23 min.

10

Example 19 (1H-Indazol-3-yl)-[6-methyl-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-amine (II-19): Prepared in 42% yield. Melting point 235-237°C; <sup>1</sup>HNMR (500 MHz, DMSO) δ 2.44 (3H, s), 7.09 (1H, J=7.5 Hz, t), 7.40 (1H, J=7.1 Hz, t), 7.49 (1H, J=8.3 Hz, d), 7.70 (3H, m), 7.79 (1H, J=7.3 Hz, t), 7.87 (1H, J=8.3 Hz, d), 8.03 (1H, J=7.7 Hz, d), 10.3 (1H, s), 12.6 (1H, s) ppm; HPLC-Method A, R<sub>t</sub> 2.958 min; MS (FIA) 370.2 (M+H)<sup>+</sup>.

15

Example 20 (1H-Indazol-3-yl)-[6-phenyl-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-amine (II-20): Prepared in 32% yield. <sup>1</sup>HNMR (500 MHz, DMSO) δ 6.94 (1H, J=7.4 Hz, t), 7.24 (1H, J=7.4 Hz, t), 7.33 (1H, J=8.4 Hz, d), 7.42 (3H, m), 7.57 (1H, J=7.3 Hz, t), 7.68 (2H, m), 7.75 (1H, J=7.9 Hz, d), 7.93 (3H, m), 8.18 (1H, br s), 10.45 (1H, br s), 12.5 (1H, br s) ppm; HPLC-Method A, R<sub>t</sub> 4.0 min; MS (FIA) 432.2 (M+H)<sup>+</sup>.

25

Example 21 (1H-Indazol-3-yl)-[6-(pyridin-4-yl)-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-amine (II-21): Prepared in 12% yield. <sup>1</sup>HNMR (500 MHz, DMSO) δ 7.16 (1H, J=7.4 Hz, t), 7.46 (1H, J=7.6 Hz, t), 7.56 (1H, J=8.3 Hz, d), 7.80 (1H, J=7.2 Hz, t), 7.90 (2H, m), 7.97 (1H, J=7.8

30

Hz, d), 8.09 (1H, br), 8.22 (2H, J=4.9 Hz, d), 8.45 (1H, br s), 8.93 (2H, J=4.8 Hz, d), 10.9 (1H, br s), 12.8 (1H, br s) ppm; HPLC-Method A,  $R_t$  3.307 min; MS (FIA) 433.2 (M+H)<sup>+</sup>

5

Example 22 (1H-Indazol-3-yl)-[6-(pyridin-2-yl)-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-amine (II-22):

Prepared in 42% yield. <sup>1</sup>HNMR (500 MHz, DMSO)  $\delta$  7.07 (1H, J=7.4 Hz, t), 7.36 (1H, J=7.4 Hz, t), 7.46 (1H, J=7.4 Hz, d), 7.53 (1H, J=5.0 Hz, t), 7.70 (1H, J=7.4 Hz, t), 7.79 (1H, J=7.1 Hz, t), 7.83 (1H, J=7.4 Hz, d), 7.88 (1H, J=7.8 Hz, d), 7.97 (1H, J=7.7 Hz, t), 8.02 (1H, J=5.5 Hz, br d), 8.36 (1H, J=7.8 Hz, d), 8.75 (2H, J=4.1 Hz, d), 10.5 (1H, br s), 12.7 (1H, br s) ppm; HPLC-Method A,  $R_t$  3.677 min; MS (FIA) 433.2 (M+H)<sup>+</sup>.

15

Example 23 [6-(2-Chlorophenyl)-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-23):

Prepared in 44% yield; <sup>1</sup>HNMR (500 MHz, DMSO)  $\delta$  7.08 (1H, J=7.5 Hz, t), 7.37 (1H, J=7.5 Hz, t), 7.45 (1H, J=8.4 Hz, d), 7.51 (2H, m), 7.61 (1H, J=7.4, 1.9 Hz, dd), 7.69 (2H, m), 7.79 (2H, J=4.0 Hz, d), 7.86 (3H, J=7.8 Hz, d), 8.04 (2H, J=6.2 Hz, br d), 10.7 (1H, br s), 12.6 (1H, br s) ppm; HPLC-Method A,  $R_t$  3.552 min; MS (FIA) 466.2 (M+H)<sup>+</sup>.

25

Example 24 [5,6-Dimethyl-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-24): Prepared

in 35% yield; mp 183-186°C; <sup>1</sup>HNMR (500 MHz, DMSO)  $\delta$  2.14 (3H, s), 2.27 (3H, s), 6.85 (1H, J=7.5 Hz, t), 7.15 (1H, J=7.6 Hz, t), 7.32 (3H, m), 7.38 (1H, J=7.5 Hz, t), 7.42 (1H, J=7.4 Hz, t), 7.53 (1H, J=7.6 Hz, d), 8.88 (1H, s), 12.5 (1H, s) ppm; HPLC-Method A,  $R_t$  2.889 min.; MS (FIA) 384.2 (M+H)<sup>+</sup>.

30

Example 25 [5,6-Dimethyl-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-25):

Prepared in 44% yield. Melting point 160-163°C; <sup>1</sup>HNMR

5 (500 MHz, DMSO) δ 2.27 (3H, s), 2.40 (3H, s), 7.16 (2H, m), 7.44 (2H, m), 7.52 (1H, J=7.4 Hz, t), 7.57 (1H, J=7.4 Hz, t), 7.67 (1H, J=7.8 Hz, d), 9.03 (1H, s), 12.75 (1H, s) ppm; HPLC-Method A, R<sub>t</sub> 2.790 min; MS (FIA) 402.2 (M+H)<sup>+</sup>.

10

Example 26 [2-(2-Chlorophenyl)-5,6-dimethyl-pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-26): Prepared in 30%

yield. <sup>1</sup>HNMR (500 MHz, DMSO) δ 2.14 (3H, s), 2.33 (3H, s), 6.84 (1H, J=7.4 Hz, t), 7.13 (1H, J=7.4 Hz, t), 7.19  
15 (1H, J=6.9 Hz, br t), 7.27 (1H, J=7.4 Hz, d), 7.32 (3H, br m), 7.37 (1H, J=7.1 Hz, d), 10.0 (1H, br), 12.8 (1H, br s) ppm; δ 2.919 min; MS (FIA) 350.1 (M+H)<sup>+</sup>.

Example 27 [5,6-Dimethyl-2-(2-trifluoromethyl-phenyl)-

20 pyrimidin-4-yl]-(7-fluoro-1H-indazol-3-yl)-amine (II-27):

Prepared in 92% yield. <sup>1</sup>HNMR (500 MHz, DMSO) δ 2.33

(3H, s), 2.50 (3H, s), 6.97 (1H, m), 7.15 (1H, m), 7.30 (1H, J=8.1 Hz, d), 7.65 (3H, m), 7.76 (1H, J=7.5 Hz, d), 10.0 (1H, s), 13.4 (1H, s) ppm; HPLC-Method A, R<sub>t</sub> 3.053  
25 min; MS (FIA) 402.2 (M+H)<sup>+</sup>.

Example 28 (5,7-Difluoro-1H-indazol-3-yl)-[5,6-Dimethyl-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-amine (II-

28): Prepared in 50% yield. <sup>1</sup>HNMR (500 MHz, DMSO) δ 2.42

30 (3H, s), 2.63 (3H, s), 7.22 (1H, J=7.6 Hz, d), 7.38 (1H, J=9.3, 1.7 Hz, dt), 7.71 (1H, m), 7.75 (1H, J=7.0 Hz, d), 7.79 (1H, J=6.7 Hz, d), 7.86 (1H, J=8.0 Hz, d), 10.0 (1H,

s), 13.2 (1H, s) ppm; HPLC-Method A,  $R_t$  3.111 min; MS (FIA) 420.2 (M+H)<sup>+</sup>.

5 Example 29 [2-(2-Chlorophenyl)-5,6-dimethyl-pyrimidin-4-yl]-(5,7-difluoro-1H-indazol-3-yl)-amine (II-29):

Prepared in 58% yield. <sup>1</sup>HNMR (500 MHz, DMSO)  $\delta$  2.47 (3H, s), 2.66 (3H, s), 7.44 (2H, m), 7.53 (1H, m), 7.64 (3H, m), 10.4 (1H, br), 13.8 (1H, br s) ppm; HPLC-Method A,  $R_t$  2.921 min; MS (FIA) 386.1 (M+H)<sup>+</sup>.

10

Example 30 [2-(2-Chlorophenyl)-5,6-dimethyl-pyrimidin-4-yl]-(7-fluoro-1H-indazol-3-yl)-amine (II-30): Prepared in 70% yield. <sup>1</sup>HNMR (500 MHz, DMSO)  $\delta$  2.35 (3H, s), 2.51 (3H, s), 7.03 (1H, J=7.8, 4.4 Hz, dt), 7.22 (1H, m), 7.33 (1H, J=7.4 Hz, t), 7.42 (1H, m), 9.19 (1H, s), 13.3 (1H, s) ppm; HPLC-Method A,  $R_t$  2.859 min; MS (FIA) 368.2 (M+H)<sup>+</sup>.

15

Example 31 [2-(2-Chlorophenyl)-5,6-dimethyl-pyrimidin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-31): Prepared in 86% yield. <sup>1</sup>HNMR (500 MHz, DMSO)  $\delta$  2.49 (3H, s), 2.68 (3H, s), 7.38 (1H, J=9.0 Hz, t), 7.54 (2H, m), 7.67 (4H, m), 10.5 (1H, br), 13.2 (1H, br s) ppm; HPLC-Method A,  $R_t$  2.850 min; MS (FIA) 368.1 (M+H)<sup>+</sup>.

20

Example 32 [2-(2,4-Dichlorophenyl)-5,6-dimethyl-pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-32): Prepared in 52% yield. <sup>1</sup>HNMR (500 MHz, DMSO)  $\delta$  2.46 (3H, s), 2.64 (3H, s), 7.16 (1H, J=7.5 Hz, t), 7.46 (1H, J=7.6 Hz, t), 7.61 (2H, m), 7.68 (2H, J=8.2 Hz, d), 7.82 (1H, m), 10.2 (1H, br), 13.0 (1H, br s) ppm; HPLC-Method A,  $R_t$  2.983 min; MS (FIA) 384.1 (M+H).

25

30

Example 33 (5-Methyl-2H-pyrazol-3-yl)-[2-(2-methylphenyl)-quinazolin-4-yl]-amine (II-33): <sup>1</sup>HNMR (DMSO) δ 1.21 (3H, s), 2.25 (3H, s), 6.53 (1H, s), 7.38 (4H, m), 7.62 (1H, d), 7.73 (1H, d), 7.81 (1H, d), 7.89 (1H, t), 8.70 (1H, s), 12.20 (1H, s); MS 316.3 (M+H)<sup>+</sup>.

Example 34 [2-(2,4-Difluorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (II-34): <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.4 (br s, 1H), 10.8 (br s, 1H), 8.58 (d, 1H), 7.97 (m, 1H), 8.36 (m, 1H), 7.85 (m, 1H), 7.60 (m, 1H), 6.62 (s, 1H), 2.30 (s, 3H); MS 338.07 (M+H).

Example 35 [2-(2,5-Dimethoxyphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (II-35): <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.5 (br s, 1H), 8.68 (br, 1H), 7.92 (t, J = 7.5 Hz, 1H), 7.86 (d, J = 8.2 Hz, 1H), 7.65 (t, J = 7.5 Hz, 1H), 7.45 (s, 1H), 7.14 (m, 2H), 6.51 (s, 1H), 3.79 (s, 3H), 3.67 (s, 3H), 2.14 (s, 3H); MS 362.2 (M+H).

Example 36 [2-(2-Chlorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (II-36): <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.8 (br, 1H), 8.80 (d, J = 8.3 Hz, 1H), 8.00 (t, J = 7.6 Hz, 1H), 7.82 (d, J = 8.3 Hz, 1H), 7.78 (m, 2H), 7.67 (d, J = 7.8 Hz, 1H), 7.61 (t, J = 7.0 Hz, 1H), 7.55 (t, J = 7.4 Hz, 1H), 6.56 (s, 1H), 2.18 (s, 3H); MS 336.1 (M+H).

Example 37 [2-(2-Methoxyphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (II-37): <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.78 (s, br, 1H), 8.00 (t, J = 7.4 Hz, 1H), 7.90 (m, 2H), 7.74 (t, J = 7.5 Hz, 1H), 7.63 (t, J = 7.3 Hz, 1H), 7.30 (d, J = 8.4 Hz, 1H), 7.18 (t, J = 7.5 Hz,

1H), 6.58 (s, br, 1H), 3.90 (s, 3H), 2.21 (s, 3H); MS  
332.1 (M+H).

Example 38 [2-(2,6-Dimethylphenyl)-quinazolin-4-yl]-(5-  
5 methyl-2H-pyrazol-3-yl)-amine (II-38): <sup>1</sup>HNMR (500 MHz,  
DMSO-d<sub>6</sub>) δ 12.2 (s, br, 2H), 8.88 (d, J = 7.7 Hz, 1H),  
8.05 (t, J = 7.7 Hz, 1H), 7.80 (m, 2H), 7.37 (t, J = 7.6  
Hz, 1H), 7.21 (d, J = 7.7 Hz, 2H), 6.36 (s, 1H), 2.16 (s,  
3H), 2.15 (s, 6H); MS 330.1 (M+H).

10

Example 39 [2-(2-Acetylphenyl)-quinazolin-4-yl]-(5-  
methyl-2H-pyrazol-3-yl)-amine (II-39): <sup>1</sup>HNMR (500 MHz,  
DMSO-d<sub>6</sub>) δ 12.35 (s, br, 1H), 8.93 (d, J = 8.4 Hz, 1H),  
8.37 (d, J = 8.6 Hz, 1H), 8.20 (d, J = 7.6 Hz, 1H), 8.11  
15 (t, J = 8.0 Hz, 2H), 7.89 (m, 2H), 7.77 (m, 2H), 6.93 (s,  
1H), 2.33 (s, 3H), 2.04 (s, 3H) MS 344.1 (M+H).

Example 40 [2-(2,3-Dimethylphenyl)-quinazolin-4-yl]-(5-  
methyl-2H-pyrazol-3-yl)-amine (II-40): <sup>1</sup>HNMR (500 MHz,  
20 DMSO-d<sub>6</sub>) δ 12.6 (s, br, 1H), 12.1 (s, br, 1H), 8.91 (d, J  
= 7.7 Hz, 1H), 8.14 (t, J = 7.2 Hz, 1H), 7.95 (d, J = 8.4  
Hz, 1H), 7.89 (t, J = 7.7 Hz, 1H), 7.58 (d, J = 7.6 Hz,  
1H), 7.53 (d, J = 7.0 Hz, 1H), 7.42 (t, J = 7.6 Hz, 1H),  
6.60 (s, 1H), 2.43 (s, 3H), 2.35 (s, 3H), 2.32 (s, 3H);  
25 MS 330.1 (M+H).

Example 41 (5-Methyl-2H-pyrazol-3-yl)-[2-(2-  
trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-41):  
<sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.3 (s, 1H), 10.5 (s, 1H),  
30 8.77 (d, J = 8.2 Hz, 1H), 7.92 (m, 2H), 7.85 (m, 3H),  
7.56 (t, J = 8.1 Hz, 1H), 7.67 (t, J = 7.4 Hz, 1H), 6.63  
(s, 1H), 2.27 (s, 3H); MS 370.1 (M+H).

Example 42 [2-(2-Ethylphenyl)-quinazolin-4-yl]-(5-Methyl-2H-pyrazol-3-yl)-amine (II-42): <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>)

5    δ 8.80 (m, 1H), 8.02 (s, br, 1H), 7.82 (d, J = 8.4 Hz, 1H), 7.77 (m, 1H), 7.62 (d, J = 7.6 Hz, 1H), 7.54 (m, 1H), 7.41 (m, 2H), 6.40 (s, 1H), 2.75 (q, J = 7.1 Hz, 2H), 2.17 (s, 3H), 0.99 (t, J = 7.5 Hz, 3H); MS 330.1 (M+H).

10    Example 43 (2-Biphenyl-2-yl-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (II-43): <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>)

δ 8.76 (d, J = 7.6 Hz, 1H), 8.04 (m, 1H), 7.75 (m, 6H), 7.30 (m, 5H), 5.34 (s, 1H), 2.14 (s, 3H); MS 378.2 (M+H).

15    Example 44 [2-(2-Hydroxyphenyl)-quinazolin-4-yl]-(5-Methyl-2H-pyrazol-3-yl)-amine (II-44): <sup>1</sup>HNMR (500 MHz,

DMSO-d<sub>6</sub>) δ 10.9 (s, br, 1H), 8.62 (d, J = 8.2 Hz, 1H), 8.28 (d, J = 7.9 Hz, 1H), 7.87 (m, 2H), 7.60 (t, J = 7.9 Hz, 1H), 7.37 (t, J = 7.8 Hz, 1H), 6.92 (m, 2H), 6.45 (s, 20 1H), 2.27 (s, 3H); MS 318.1 (M+H).

Example 45 [2-(2-Ethoxyphenyl)-quinazolin-4-yl]-(5-Methyl-2H-pyrazol-3-yl)-amine (II-45): <sup>1</sup>HNMR (500 MHz,

DMSO-d<sub>6</sub>) δ 12.1 (s, br, 1H), 8.75 (d, J = 8.3 Hz, 1H), 7.97 (t, J = 7.8 Hz, 1H), 7.82 (d, J = 8.3 Hz, 1H), 7.78 (d, J = 7.5 Hz, 1H), 7.70 (t, J = 7.8 Hz, 1H), 7.56 (t, J = 7.8 Hz, 1H), 7.22 (d, J = 8.4 Hz, 1H), 7.12 (t, J = 7.6 Hz, 1H), 6.55 (s, 1H), 4.11 (q, J = 6.9 Hz, 2H), 2.16 (s, 25 3H), 1.22 (t, J = 6.9 Hz, 3H); MS 346.1 (M+H).

30

Example 46 [5-(Thiophen-2-yl)-2H-pyrazol-3-yl]-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-46):

<sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.04 (d, J = 8.3 Hz, 1H), 8.05



(dd,  $J = 7.3, 8.2$  Hz, 1H), 7.93 (d,  $J = 6.5$  Hz, 1H), 7.81 (m, 5H), 7.34 (d,  $J = 5.0$  Hz, 1H), 7.25 (m, 1H), 7.00 (m, 1H), 6.87 (s, 1H); MS 438.1 (M+H).

5 Example 47 [4-(Thiophen-2-yl)-2H-pyrazol-3-yl]-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-47):

Prepared according to Method B.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  6.97 (m, 1H), 7.08 (m, 1H), 7.27 (m, 1H), 7.36 (m, 1H), 7.66 (m, 2H), 7.77 (m, 3H), 7.83 (m, 1H), 8.00 (m, 1H), 8.18 (s, 1H), 8.62 (d,  $J = 8.2$  Hz, 1H), 10.7 (br. s, 1H); EI-MS 438.1 (M+H); HPLC-Method A,  $R_t$  2.97 min.

Example 48 (4-Phenyl-2H-pyrazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-48):

15 Prepared according to Method B.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  7.05 (br. s, 1H), 7.14 (t,  $J = 7.8$  Hz, 1H), 7.25 (m, 3H), 7.43 (m, 2H), 7.60 (m, 2H), 7.73 (m, 2H), 7.80 (d, 1H), 7.95 (m, 1H), 8.12 (br. s, 1H), 8.60 (m, 1H), 10.6 (br. s, 1H); EI-MS 432.2 (M+H); HPLC-Method A,  $R_t$  3.04 min.

20

Example 49 (5-tert-Butyl-2H-pyrazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-49):

$^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  8.76 (d,  $J = 8.3$  Hz, 1H), 7.94 (m, 2H), 7.79 (m, 4H), 7.70 (t,  $J = 7.6$  Hz, 1H), 6.51 (s, 1H), 1.16 (s, 9H); MS 412.2 (M+H).

25

Example 50 (5-Phenyl-2H-pyrazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-50):

$^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  7.09 (s, 1H), 7.36 (td,  $J = 7.8, 1.1$  Hz, 1H), 7.46 (t,  $J = 7.8$  Hz, 2H), 7.65 (br. d,  $J = 8.1$  Hz, 2H), 7.78 (m, 2H), 7.90 (m, 4H), 7.95 (d,  $J = 7.7$  Hz, 1H), 8.00 (t,  $J = 7.8$  Hz, 1H), 8.81 (d,  $J = 8.6$  Hz, 1H), 11.29 (br. s, 1H); EI-MS 432.1 (M+H); HPLC-Method A,  $R_t$  3.24 min.

30

Example 51 (4,5-Diphenyl-2H-pyrazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-51):

<sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 7.13 (m, 1H), 7.18 (m, 5H), 7.36 (m, 5H), 7.62 (m, 3H), 7.73 (m, 2H), 7.85 (m, 1H), 8.48 (d, J = 8.7 Hz, 1H), 10.02 (s, 1H), 13.19 (s, 1H); EI-MS 508.2 (M+H); HPLC-Method A, R<sub>t</sub> 3.39 min.

Example 52 (4-Carbamoyl-2H-pyrazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-52):

Prepared in 40% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>): δ 12.85 (s, 1H), 12.77 (s, 1H), 11.80 (s, 1H), 10.80 (s, 1H), 8.35-7.42 (m, 9H); MS 399.13 (M+H) HPLC-Method A, R<sub>t</sub> 2.782 min.

Example 53 (2H-Pyrazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-53):

Prepared in 38% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.52 (s, 1H), 10.65 (s, 1H), 8.75 (d, 1H), 7.91-7.68 (m, 8H), 6.87 (s, 1H). MS: (M+H) 356.17. HPLC-Method A, R<sub>t</sub> 2.798 min.

Example 54 (5-Hydroxy-2H-pyrazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (II-54):

Prepared in 36% yield; <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 10.61 (s, 1H), 8.75 (s, 1H), 8.03-7.75 (m, 9H), 5.97 (s, 1H); MS 372.18 (M+H); HPLC-Method A, R<sub>t</sub> 2.766 min.

Example 55 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-55):

Prepared in 30% yield. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.21 (s, 1H), 10.45 (s, 1H), 8.68 (s, 1H), 7.89-7.45 (m, 8H),

6.48 (s, 1H), 0.89 (m, 2H), 0.62 (s, 2H). MS 396.18 (M+H); HPLC-Method A,  $R_t$  3.069 min.

Example 56 (5-Methoxymethyl-2H-pyrazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-56):  
5 Prepared in 33% yield;  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.51 (s, 1H), 10.48 (s, 1H), 8.60 (s, 1H), 7.81-7.55 (m, 7H), 6.71 (s, 1H), 4.28 (s, 2H), 3.18 (s, 3H). MS 400.19 (M+H); HPLC-Method A,  $R_t$  2.881 min.

10

Example 57 (1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-57): Prepared to afford 51 mg (78% yield) as pale yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.7 (s, 1H), 10.4 (s, 1H), 8.55 (d, 1H),  
15 7.81 (t, 1H), 7.71 (d, 1H), 7.61 (d, 1H), 7.58 (t, 1H), 7.46 (m, 4H), 7.36 (d, 1H), 7.22 (t, 1H), 6.91 (t, 1H) ppm; LC-MS (ES+) 406.16 (M+H), (ES-) 404.19 (M-H); HPLC-Method A,  $R_t$  3.00 min.

20 Example 58 (4-Chloro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-58): Prepared in DMF (70% yield) as pale yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.3 (s, br, 1H), 10.9 (s, br, 1H), 8.60 (d, 1H), 7.97 (t, 1H), 7.81 (d, 1H), 7.75 (t, 1H),  
25 7.67 (d, 1H), 7.63 (dd, 1H), 7.57 (m, 2H), 7.43 (d, 1H), 7.28 (dd, 1H), 7.08 (d, 1H) ppm; LC-MS (ES+) 440.10 (M+H), (ES-) 438.12 (M-H); HPLC-Method A,  $R_t$  3.08 min.

Example 59 (5-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-59):  
30 Prepared in DMF (34% yield) as pale yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.0 (s, 1H), 10.6 (s, 1H), 8.72 (d, 1H), 7.99 (t, 1H), 7.89 (d, 1H), 7.79 (d, 1H), 7.75 (t, 1H), 7.68 (m, 3H), 7.56 (dd, 1H), 7.39 (d, 1H), 7.28 (t,

1H) ppm; LC-MS (ES+) 424.12 (M+H), (ES-) m/e= 422.13 (M-H); HPLC-Method A, R<sub>t</sub> 3.05 min.

Example 60 (7-Fluoro-1H-indazol-3-yl)-[2-(2-

5 trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-60):

Prepared in DMF (51% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.4 (s, 1H), 10.6 (s, 1H), 8.68 (d, 1H), 7.95 (t, 1H), 7.85 (d, 1H), 7.72 (m, 2H), 7.63 (m, 2H), 7.58 (m, 1H), 7.43 (d, 1H), 7.18 (dd, 1H), 7.00 (m, 1H)

10 ppm; LC-MS (ES+) 424.11 (M+H), (ES-) 422.15 (M-H); HPLC-Method A, R<sub>t</sub> 3.06 min.

Example 61 (5-Methyl-1H-indazol-3-yl)-[2-(2-

trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-61):

15 Prepared in DMF (81% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.0 (s, br, 1H), 8.79 (br, 1H), 8.11 (br, 1H), 7.96 (d, 1H), 7.82 (m, 5H), 7.46 (s, 1H), 7.41 (d, 1H), 7.20 (d, 1H), 2.33 (s, 3H) ppm; MS (ES+) 420.15 (M+H), (ES-) 418.17 (M-H); HPLC-Method A, R<sub>t</sub> 3.07 min.

20

Example 62 [2-(2,6-Dichloro-phenyl)-quinazolin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-62): Prepared in DMF (37% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>)

δ 13.0 (s, 1H), 10.8 (s, 1H), 8.72 (d, 1H), 7.97 (t, 1H), 7.90 (d, 1H), 7.75 (t, 1H), 7.53 (m, 3H), 7.43 (t, 1H), 7.35 (d, 1H), 7.23 (t, 1H) ppm; LCMS (ES+) 424.08 (M+H), (ES-) 422.10 (M-H); HPLC-Method A, R<sub>t</sub> 3.06 min.

25

Example 63 [2-(2-Chloro-phenyl)-quinazolin-4-yl]-(1H-

30 indazol-3-yl)-amine (II-63): Prepared in 91% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 7.06 (t, 1H), 7.36 (t, 1H), 7.39 (t, 1H), 7.52 (m, 3H), 7.62 (d, 1H), 7.72 (d, 1H), 7.82 (m, 1H), 7.90 (d, 1H), 8.05 (m, 1H), 8.76 (d, 1H), 11.5 (m, 1H), 13.02 (s, 1H); EI-MS 372.1 (M+1); HPLC-Method A, R<sub>t</sub>

2.93 min.

Example 64 (5-Trifluoromethyl-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-64):

5 Prepared in DMF (57% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.4 (s, br, 1H), 11.4 (br, 1H), 8.72 (d, 1H), 8.12 (s, 1H), 7.98 (t, 1H), 7.83 (d, 1H), 7.76 (d, 1H), 7.73 (dd, 1H), 7.60 (m, 4H), 7.52 (d, 1H) ppm; LC-MS (ES+) 474.12 (M+H), (ES-) 472.17 (M-H); HPLC-Method A, R<sub>t</sub>  
10 3.25 min.

Example 65 (4-Trifluoromethyl-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-65):

Prepared in DMF (8% yield) as yellow solid. <sup>1</sup>HNMR (500  
15 MHz, DMSO-d<sub>6</sub>) δ 13.7 (s, br, 1H), 11.2 (br, 1H), 8.70 (d, 1H), 8.05 (s, 1H), 7.85 (m, 3H), 7.65 (m, 4H), 7.51 (m, 2H) ppm; LC-MS (ES+) 474.13 (M+H), (ES-) 472.17 (M-H);  
HPLC-Method A, R<sub>t</sub> 3.15 min.

20 Example 66 [2-(2,6-Dichloro-phenyl)-quinazolin-4-yl]-(1H-indazol-3-yl)-amine (II-66): Prepared in DMF (30% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1H), 11.1 (s, 1H), 8.69 (d, 1H), 7.95 (t, 1H), 7.82 (d, 1H), 7.73 (t, 1H), 7.56 (d, 1H), 7.47 (s, 1H), 7.45 (s, 1H),  
25 7.39 (m, 2H), 7.26 (t, 1H), 6.92 (t, 1H) ppm; LC-MS (ES+) 406.11 (M+H), (ES-) 404.12 (M-H); HPLC-Method A, R<sub>t</sub> 3.00 min.

Example 67 (1H-indazol-3-yl)-[2-(2-methyl-phenyl)-

30 quinazolin-4-yl]-amine (II-67): Prepared in 55% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 2.15 (s, 3H), 7.09 (t, 1H), 7.26 (d, 1H), 7.31 (t, 1H), 7.39 (t, 1H), 7.42 (m, 1H), 7.55 (d 1H), 7.64 (d, 1H), 7.74 (d, 1H), 7.89 (m, 1H), 7.96

(d, 1H), 8.10 (m, 1H), 8.81 (d, 1H), 12.0 (m, 1H), 13.18 (s, 1H); EI-MS 352.2 (M+1); HPLC-Method A,  $R_t$  2.93 min.

Example 68 (7-Trifluoromethyl-1H-indazol-3-yl)-[2-(2-

5 trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-68):

Prepared in DMF (75% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.5 (s, br, 1H), 11.2 (s, br, 1H), 8.68 (d, 1H), 7.97 (t, 1H), 7.92 (d, 1H), 7.82 (d, 1H), 7.74 (t, 1H), 7.70 (d, 1H), 7.68 (d, 1H), 7.64 (m, 2H), 7.57 (m, 1H), 7.14 (t, 1H) ppm; LC-MS (ES+) 474.11 (M+H), (ES-) 472.14 (M-H); HPLC-Method A,  $R_t$  3.24 min.

Example 69 (6-Trifluoromethyl-1H-indazol-3-yl)-[2-(2-

15 trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-69):

Prepared by Method B in DMF (78% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.4 (s, br, 1H), 11.1 (s, br, 1H), 8.67 (d, 1H), 7.95 (t, 1H), 7.82 (m, 3H), 7.72 (m, 2H), 7.63 (m, 2H), 7.57 (t, 1H), 7.23 (d, 1H) ppm; LC-MS (ES+) 474.12 (M+H), (ES-) 472.15 (M-H); HPLC-Method A,  $R_t$  3.28 min.

Example 70 (5-Nitro-1H-indazol-3-yl)-[2-(2-

trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-70):

Prepared in DMF (82% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.6 (s, br, 1H), 11.4 (s, br, 1H), 8.75 (s, 1H), 8.72 (d, 1H), 8.09 (dd, 1H), 7.98 (t, 1H), 7.83 (d, 1H), 7.75 (t, 1H), 7.70 (m, 2H), 7.61 (m, 3H) ppm; LC-MS (ES+) 451.14 (M+H), (ES-) 449.12 (M-H); HPLC-Method A,  $R_t$  3.02 min.

Example 71 (5,7-Difluoro-1H-indazol-3-yl)-[2-(2-

trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-71):

Prepared in DMF (60% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.7 (s, br, 1H), 11.2 (s, br, 1H), 8.73

(d, 1H), 8.03 (t, 1H), 7.88 (d, 1H), 7.80 (m, 2H), 7.70 (m, 3H), 7.32 (m, 2H) ppm; LC-MS (ES+) 442.14 (M+H), (ES-) 440.14 (M-H); HPLC-Method A,  $R_t$  3.11 min.

- 5 Example 72 (4-Pyrrol-1-yl-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-72): Prepared in DMF (33% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.4 (s, br, 1H), 11.0 (s, br, 1H), 8.53 (d, 1H), 7.98 (t, 1H), 7.75 (m, 4H), 7.62 (m, 2H), 7.52 (d, 1H), 7.43 (t, 1H), 7.05 (d, 1H), 6.80 (s, 2H), 5.61 (s, 2H) ppm; LC-MS (ES+) 471.18 (M+H), (ES-) 469.18 (M-H); HPLC-Method A,  $R_t$  3.12 min.

- Example 73 (5-Amino-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-73): A solution of compound II-70 (70 mg, 0.16 mmol) in MeOH (2 mL) was treated with Raney Ni until solution was colorless (about 1.5 g Raney Ni was added). After stirring at room temperature for 40 min, the mixture was filtered through celite, the resulting celite was washed with MeOH (5 times), and the solvent was evaporated in vacuo to provide a crude product that was then purified by HPLC to give the title compound as a yellow solid (10 mg, 15%). m.p. 221-223°C;  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.2 (s, br, 1H), 10.7 (s, br, 1H), 9.80 (br, 2H), 8.68 (d, 1H), 7.97 (t, 1H), 7.87 (d, 1H), 7.75 (m, 2H), 7.65 (m, 5H), 7.30 (d, 1H) ppm; MS (ES+) 421.16 (M+H), (ES-) 419.17 (M-H); HPLC-Method A,  $R_t$  2.41 min.

- 30 Example 74 [2-(2-Chloro-phenyl)-quinazolin-4-yl]-(7-fluoro-1H-indazol-3-yl)-amine (II-74): Prepared in DMF (35% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.7 (s, 1H), 11.7 (s, br, 1H), 8.80 (d, 1H), 8.15 (t, 1H), 7.99 (d, 1H), 7.88 (t, 1H), 7.68 (d, 1H), 7.60 (m,

2H), 7.53 (t, 1H), 7.46 (t, 1H), 7.25 (dd, 1H), 7.04 (m, 1H) ppm; LC-MS (ES+) 390.16 (M+H); HPLC-Method A,  $R_t$  3.00 min.

- 5 Example 75 [2-(2-Chloro-phenyl)-quinazolin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-75): Prepared in DMF.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.2 (s, 1H), 11.7 (s, br, 1H), 8.80 (d, 1H), 8.10 (t, 1H), 7.91 (m, 2H), 7.70 (d, 1H), 7.58 (m, 4H), 7.50 (t, 1H), 7.29 (t, 1H) ppm; LC-MS (ES+) 390.17 (M+H); HPLC-Method A,  $R_t$  3.00 min.

- Example 76 [2-(2-Chloro-phenyl)-quinazolin-4-yl]-(5,7-difluoro-1H-indazol-3-yl)-amine (II-76): Prepared in DMF (55% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.8 (s, 1H), 11.5 (s, br, 1H), 8.76 (d, 1H), 8.08 (t, 1H), 7.93 (d, 1H), 7.84 (t, 1H), 7.64 (d, 1H), 7.55 (d, 1H), 7.50 (t, 1H), 7.44 (m, 2H), 7.36 (t, 1H) ppm; LC-MS (ES+) 408.15 (M+H), (ES-) 406.17 (M-H); HPLC-Method A,  $R_t$  3.08 min.

- 20 Example 77 [2-(2-Chloro-phenyl)-quinazolin-4-yl]-(5-trifluoromethyl-1H-indazol-3-yl)-amine (II-77): Prepared in DMF (66% yield) as yellow solid.  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.5 (s, 1H), 11.4 (s, br, 1H), 8.79 (d, 1H), 8.29 (s, 1H), 8.07 (t, 1H), 7.93 (d, 1H), 7.84 (t, 1H), 7.72 (d, 1H), 7.63 (d, 2H), 7.53 (d, 1H), 7.48 (t, 1H), 7.36 (t, 1H) ppm; LC-MS (ES+):  $m/e$  = 440.16 (M+H); (ES-):  $m/e$  = 438.18 (M-H); HPLC-Method A,  $R_t$  3.22 min.

- 30 Example 78 [2-(2-cyano-phenyl)-quinazolin-4-yl]-(1H-indazol-3-yl)-amine (II-78): Prepared in 13% yield.  $^1\text{H}$ -NMR (500 MHz, DMSO)  $\delta$  12.9 (br, 1H), 10.8 (br, 1H), 8.73 (br s, 1H), 7.97 (m, 4H), 7.74 (m, 1H), 7.5 (m, 4H), 7.42



(m, 1H), 7.08 (m, 1H) ppm; MS (FIA) 363.2 (M+H); HPLC-Method A, R<sub>t</sub> 2.971 min.

Example 79 (5-Bromo-1H-indazol-3-yl)-[2-(2-

5 trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-79):

Prepared in DMF (64% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.4 (s, 1H), 11.6 (s, br, 1H), 8.93 (d, 1H), 8.21 (t, 1H), 8.14 (s, 1H), 8.05 (d, 1H), 7.95 (m, 4H), 7.86 (t, 1H), 7.65 (d, 1H), 7.59 (d, 1H) ppm; MS

10 (ES+) 486.10 (M+H), (ES-) 484.09 (M-H); HPLC-Method A, R<sub>t</sub> 3.22 min.

Example 80 (6-Chloro-1H-indazol-3-yl)-[2-(2-

trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-80):

15 Prepared in DMF (94% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.1 (s, 1H), 11.2 (s, br, 1H), 8.73 (d, 1H), 8.03 (t, 1H), 7.87 (d, 1H), 7.79 (m, 2H), 7.73 (m, 2H), 7.67 (m, 2H), 7.58 (s, 1H), 7.04 (dd, 1H) ppm. LC-MS (ES+) 440.14 (M+H), (ES-) 438.16 (M-H); HPLC-Method A, R<sub>t</sub>

20 3.25 min.

Example 81 (7-Fluoro-6-trifluoromethyl-1H-indazol-3-yl)-

[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-81): Prepared in DMF (30% yield) as yellow solid. <sup>1</sup>HNMR

25 (500 MHz, DMSO-d<sub>6</sub>) δ 13.9 (s, 1H), 11.0 (s, br, 1H), 8.64 (d, 1H), 7.94 (t, 1H), 7.81 (d, 1H), 7.71 (m, 2H), 7.60 (m, 4H), 7.20 (dd, 1H) ppm. LC-MS (ES+) 492.18 (M+H), (ES-) 490.18 (M-H); HPLC-Method A, R<sub>t</sub> 3.44 min.

30 Example 82 (6-Bromo-1H-indazol-3-yl)-[2-(2-

trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-82):

Prepared in DMF (40% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.1 (s, 1H), 11.2 (s, br, 1H), 8.73 (d, 1H), 8.03 (t, 1H), 7.87 (d, 1H), 7.80 (m, 2H), 7.73 (m,

3H), 7.67 (m, 1H), 7.61 (d, 1H), 7.15 (dd, 1H) ppm; MS (ES+) 486.07 (M+H); HPLC-Method A,  $R_t$  3.28 min.

Example 83 [2-(2,4-Bis-trifluoromethyl-phenyl)-

5 quinazolin-4-yl]-(5,7-difluoro-1H-indazol-3-yl)-amine (II-83): Prepared in DMF in 28% yield.  $^1\text{HNMR}$  (500MHz, MeOH- $d_4$ )  $\delta$  8.81 (d,  $J=8.4\text{Hz}$ , 1H), 8.35-8.20 (m, 3H), 8.19-7.96 (m, 3H), 7.40-7.34 (m, 1H), 7.29-7.14 (m, 1H); LC-MS (ES+) 510.14 (M+H); HPLC-Method C,  $R_t$  8.29 min.

10

Example 84 (5,7-Difluoro-1H-indazol-3-yl)-[2-(4-fluoro-2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-84):

Prepared in 48% yield.  $^1\text{HNMR}$  (500MHz, MeOH- $d_4$ )  $\delta$  8.74-8.63 (m, 1H), 8.23-8.10 (m, 1H), 7.99-7.90 (m, 2H), 7.89-15 7.80 (m, 1H), 7.71-7.61 (m, 1H), 7.61-7.50 (m, 1H), 7.24-7.15 (m, 1H), 7.14-7.02 (m, 1H); LC-MS (ES+) 460.14 (M+H); HPLC-Method C,  $R_t$  7.59 min.

Example 85 [2-(2-Bromo-phenyl)-quinazolin-4-yl]-(5,7-

20 difluoro-1H-indazol-3-yl)-amine (II-85): Prepared in THF (21% yield).  $^1\text{HNMR}$  (500MHz, MeOH- $d_4$ )  $\delta$  8.81 (d,  $J=8.4\text{Hz}$ , 1H), 8.35-8.20 (m, 3H), 8.19-7.96 (m, 3H), 7.40-7.34 (m, 1H), 7.29-7.14 (m, 1H); LC-MS (ES+) 510.14 (M+H); HPLC-Method C,  $R_t$  8.29 min.

25

Example 86 (5,7-Difluoro-1H-indazol-3-yl)-[2-(5-fluoro-2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-86):

Prepared in THF (26% yield).  $^1\text{HNMR}$  (500MHz, MeOH- $d_4$ )  $\delta$  8.62 (d,  $J=8.4\text{Hz}$ , 1H), 8.16-8.02 (m, 1H), 7.96-7.73 (m, 30 3H), 7.59-7.48 (m, 1H), 7.48-7.35 (m, 1H), 7.21-7.09 (m, 1H), 7.09-6.89 (m, 1H); LC-MS (ES+) 460.16 (M+H); HPLC-Method C,  $R_t$  7.28 min.

**Example 87** [2-(2,4-Dichloro-phenyl)-quinazolin-4-yl]-(5,7-Difluoro-1H-indazol-3-yl)-amine (II-87): Prepared in THF (16% yield). <sup>1</sup>HNMR (500MHz, MeOH-d<sub>4</sub>) δ 8.81 (d, J=8.4Hz, 1H), 8.35-8.20 (m, 3H), 8.19-7.96 (m, 3H), 7.40-7.34 (m, 1H), 7.29-7.14 (m, 1H); LC-MS (ES+) 510.14 (M+H); HPLC-Method C, R<sub>t</sub> 8.29 min.

**Example 88** [2-(2-Chloro-5-trifluoromethyl-phenyl)-quinazolin-4-yl]-(5,7-Difluoro-1H-indazol-3-yl)-amine (II-88): Prepared in THF (33% yield). <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 10.76 (s, 1H), 8.66 (d, J=8.3Hz, 1H), 8.06-7.84 (m, 3H), 7.81-7.63 (m, 3H), 7.48-7.16 (m, 2H); LC-MS (ES+) 476.16 (M+H); HPLC-Method C, R<sub>t</sub> 19.28 min.

**Example 89** (4-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-89): Prepared in NMP (79% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.2 (s, 1H), 10.8 (s, br, 1H), 8.63 (d, 1H), 7.97 (t, 1H), 7.85 (d, 1H), 7.74 (m, 2H), 7.64 (t, 1H), 7.57 (m, 2H), 7.32 (m, 2H), 6.82 (m, 1H) ppm; LC-MS (ES+) 424.17 (M+H); HPLC-Method A, R<sub>t</sub> 3.14 min.

**Example 90** (1H-Indazol-3-yl)-[8-methoxy-2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-90): Prepared using THF as solvent to afford the title compound as a TFA salt (23% yield). HPLC-Method A, R<sub>t</sub> 2.97 min (95%); <sup>1</sup>HNMR (DMSO-d<sub>6</sub>, 500 MHz) δ 12.9 (1H, bs), 11.0 - 10.7 (1H, bs), 8.25 (1H, m), 7.75-7.50 (8H, s), 7.30 (1H, m), 6.90 (1H, m), 4.0 (3H, s); MS (m/z) 436.2 (M+H).

**Example 91** (5-Fluoro-1H-indazol-3-yl)-[8-methoxy-2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-91): Prepared using TFA as solvent to afford the title compound as a TFA salt (23% yield). HPLC-Method A, R<sub>t</sub>

3.10 min. (99%);  $^1\text{H}$ NMR (DMSO- $d_6$ , 500 MHz): 13.0 (1H, bs), 11.0 - 10.7 (1H, bs), 8.25 (1H, m), 7.75-7.50 (7H, m), 7.35 (1H, m), 7.25 (1H, m), 4.0 (3H, s); MS (m/z) 454.2 (M+H).

5

Example 92 (7-Fluoro-1H-indazol-3-yl)-[8-methoxy-2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-92):

Prepared using THF as solvent to afford the title compound as a TFA salt (98 mg, 58% yield). HPLC-Method

10 A,  $R_t$  3.20 min (92%);  $^1\text{H}$ NMR (DMSO- $d_6$ , 500 MHz)  $\delta$  13.45 (1H, bs), 11.0 - 10.7 (1H, bs), 8.25 (1H, m), 7.75-7.60 (5H, m), 7.50 (1H, m), 7.40 (1H, m), 7.15 (1H, m), 6.95 (1H, m) 4.0 (3H, s); MS (m/z) 454.2 (M+H).

15 Example 93 (5,7-Difluoro-1H-indazol-3-yl)-[8-methoxy-2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-93): Prepared using THF as solvent to afford the title compound as a TFA salt (36% yield). HPLC-Method A,  $R_t$  3.27 min. (95%);  $^1\text{H}$ NMR (DMSO- $d_6$ , 500 MHz): 13.65 (1H, bs),  
20 11.0 - 10.7 (1H, bs), 8.22 (1H, m), 7.75-7.60 (5H, m), 7.40 (1H, m), 7.35 (1H, m), 7.19 (1H, m), 4.0 (3H, s); MS (m/z) 472.2 (M+H).

Example 94 [2-(2-Chloro-pyridin-3-yl)-quinazolin-4-yl]-

25 (5,7-Difluoro-1H-indazol-3-yl)-amine (II-94): Prepared in DMF.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  13.62 (br s, 1H, 11.06-10.71 (m, 1H), 8.16-7.70 (m, 4H), 7.60-7.09 (m, 3H); LC-MS (ES+) 409.14 (M+H); HPLC-Method A,  $R_t$  2.89 min.

30 Example 95 [2-(2-Chloro-4-nitro-phenyl)-quinazolin-4-yl]- (5,7-difluoro-1H-indazol-3-yl)-amine (II-95): Prepared in THF.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  13.35 (s, 1H), 10.74 (s, 1H), 8.67 (d,  $J=8.4\text{Hz}$ , 1H), 8.29 (d,  $J=2.05\text{Hz}$ , 1H), 8.18-

8.08 (m, 1H), 8.07-7.60 (m, 4H), 7.53-7.10 (m, 2H). LC-MS (ES+) 453.15 (M+H); HPLC-Method D,  $R_t$  3.63 min.

Example 96 [2-(4-Amino-2-chloro-phenyl)-quinazolin-4-yl]-

5 (5,7-Difluoro-1H-indazol-3-yl)-amine (II-96):

A solution of compound II-95 (8mg, 0.018mmol) and tin chloride dihydrate (22mg, 0.1mmol) in ethanol (2mL) was heated at 100°C for 24h. The reaction was diluted with EtOAc (10mL), washed with 1N NaOH solution (2x10mL),  
10 brine, and dried over anhydrous sodium sulfate to afford the crude product. Purification was achieved by flash chromatography on silica gel (eluting with 1-3% MeOH in  $\text{CH}_2\text{Cl}_2$ .) The title compound was isolated as pale yellow solid (1.2mg, 16% yield). LC-MS (ES+) 423.12 (M+H),  
15 HPLC-Method C,  $R_t$  13.78 min.

Example 97 (4,5,6,7-Tetrahydro-1H-indazol-3-yl)

-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine

(II-97): Prepared in 34% yield.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$   
20 1.58 (m, 2H), 1.66 (m, 2H), 2.24 (m, 2H), 2.54 (m, 2H),  
7.63 (m, 3H), 7.71 (t, 1H), 7.75 (d, 1H), 7.78 (d, 1H),  
7.85 (t, 1H), 8.53 (d, 1H), 9.99 (s, 1H), 12.09 (s, 1H);  
EI-MS 410.2 (M+1); HPLC-Method A,  $R_t$  3.05 min.

25 Example 98 (1H-Pyrazolo[4,3-b]pyridin-3-yl)-[2-(2-

trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-98):

Prepared in DMF (37% yield) as yellow solid.  $^1\text{H}$ NMR (500  
MHz, DMSO- $d_6$ )  $\delta$  13.1 (s, br, 1H), 11.2 (s, br, 1H), 8.73  
(d, 1H), 8.54 (dd, 1H), 8.12 (d, 1H), 8.06 (t, 1H), 7.90  
30 (d, 1H), 7.84 (t, 1H), 7.75 (d, 1H), 7.69 (m, 2H), 7.65  
(t, 1H), 7.47 (dd, 1H) ppm; LC-MS (ES+) 407.18 (M+H);  
HPLC-Method A,  $R_t$  2.77 min.

Example 99 (1H-Pyrazolo[3,4-b]pyridin-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-99):

Prepared in DMF (45% yield). <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>)

5 δ 13.5 (s, br, 1H), 11.3 (s, br, 1H), 8.78 (d, 1H), 8.49 (d, 1H), 8.17 (d, 1H), 8.03 (t, 1H), 7.89 (d, 1H), 7.80 (m, 2H), 7.74 (m, 2H), 7.68 (m, 1H), 7.08 (dd, 1H) ppm. MS (ES+) 407.16 (M+H), (ES-) 405.16 (M-H); HPLC-Method A, R<sub>t</sub> 2.80 min.

10 Example 100 (6-Methyl-1H-pyrazolo[3,4-b]pyridin-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-100): Prepared in DMF (11% yield). <sup>1</sup>HNMR (500 MHz, DMSO-

d<sub>6</sub>) δ 13.2 (s, br, 1H), 10.8 (s, br, 1H), 8.57 (d, 1H), 7.95 (t, 1H), 7.82 (d, 1H), 7.72 (t, 1H), 7.65 (m, 2H), 15 7.58 (m, 2H), 2.44 (s, 3H, buried by DMSO), 2.20 (s, 3H) ppm. LC-MS (ES+) 435.22 (M+H), (ES-) 433.25 (M-H); HPLC-Method A, R<sub>t</sub> 2.94 min.

Example 101 (6-Oxo-5-phenyl-5,6-dihydro-1H-pyrazolo[4,3-c]pyridazin-3-yl)-[2-(2-trifluoromethyl-phenyl)-

20 quinazolin-4-yl]-amine II-101: Prepared in DMF (6% yield). <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.6 (s, 1H), 11.0 (s, br, 1H), 8.60 (d, 1H), 7.95 (t, 1H), 7.88 (d, 1H), 7.80 (d, 1H), 7.68 (m, 4H), 7.40 (s, 3H), 7.22 (s, 2H), 6.61 25 (s, 1H) ppm. LC-MS (ES+) 500.21 (M+H), (ES-) 498.16 (M-H); HPLC-Method A, R<sub>t</sub> 3.00 min.

Example 103 [6-Methyl-2-(2-trifluoromethoxy-phenyl)-pyrimidin-4-yl]-(5-phenyl-2H-pyrazol-3-yl)-amine (II-30 103): MS 412.13 (M+H); HPLC-Method E R<sub>t</sub> 1.248 min.

Example 104 (5-Furan-2-yl-2H-pyrazol-3-yl)-[6-methyl-2-(2-trifluoromethoxy-phenyl)-pyrimidin-4-yl]-amine (II-104); MS 402.12 (M+H); HPLC-Method E, R<sub>t</sub> 1.188 min.

Example 105 [6-Ethyl-2-(2-trifluoromethoxy-phenyl)-pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (II-105): MS 364.14 (M+H); HPLC-Method E,  $R_t$  1.112 min.

5

Example 106 [2-(2-Chloro-phenyl)-pyrido[2,3-d]pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (II-106):  $^1\text{H}$ NMR (500 MHz, DMSO)  $\delta$  12.23 (s, 1H), 10.78 (s, 1H), 7.73-7.47 (m, 7H), 6.72 (s, 1H), 2.21 (s, 3H). MS: (M+H) 337.02.

10 HPLC-Method A,  $R_t$  2.783 min.

Example 107 (5-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-6,7-dihydro-5H-cyclopentapyrimidin-4-yl]-amine (II-107): Prepared in 68% yield.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  2.16 (t, 2H), 2.88 (m, 2H), 2.98 (t, 2H), 7.21 (td, 1H), 7.29 (dd, 1H), 7.50 (dd, 1H), 7.65 (t, 1H), 7.67 (t, 1H), 7.73 (t, 1H), 7.79 (d, 1H), 10.22 (br. s, 1H), 12.99 (br. s, 1H); EI-MS 414.2 (M+H); HPLC-Method A,  $R_t$  2.92 min.

20

Example 108 (1H-Indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-pyrido[2,3-d]pyrimidin-4-yl]-amine (II-108): HPLC-Method A,  $R_t$  2.78 min. (95%);  $^1\text{H}$ NMR (DMSO- $d_6$ , 500 MHz): 12.95 (1H, bs), 11.45  $\delta$  11.15 (1H, bs), 9.20 (2H, m), 7.85-7.70 (2H, m), 7.70-7.55 (4H, m), 7.50 (1H, m), 7.35 (1H, m), 7.05 (1H, m); MS (m/z) 407.03 (M+H).

25

Example 109 (5,7-Difluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-pyrido[2,3-d]pyrimidin-4-yl]-amine (II-109): Yellow, di-TFA salt (25% yield). HPLC (Method A) 3.10 min. (95%);  $^1\text{H}$ NMR (DMSO- $d_6$ , 500 MHz): 13.8-13.6 (1H, bs), 11.4 - 11.2 (1H, bs), 9.15 (2H, m), 7.85-7.75 (2H, m), 7.75-7.62 (3H, m), 7.32 (2H, m); MS (m/z) 442.98 (M+H).

30

Example 110 [2-(2-Chloro-phenyl)-pyrido[2,3-d]pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-110): Prepared from 2-aminonicotinic acid and 2-chlorobenzoyl chloride afforded the title compound as a di-TFA salt (28% yield). HPLC-Method A,  $R_t$  2.85 min. (95%);  $^1\text{H}$ NMR (DMSO- $d_6$ , 500 MHz): 12.90 (1H, s), 11.10 - 10.90 (1H, bs), 9.05 (2H, m), 7.75-7.60 (2H, m), 7.51 (1H, m), 7.45-7.25 (5H, m), 6.95 (1H, m); MS (m/z) 372.99 (M+H).

10

Example 111 (5-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8,9,10-hexahydro-cyclooctapyrimidin-4-yl]-amine (II-111). Prepared in 43% yield.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  1.46 (m, 2H), 1.53 (m, 2H), 1.77 (m, 4H), 2.95 (m, 2H), 3.04 (m, 2H), 7.22 (m, 2H), 7.50 (dd, 1H), 7.72 (m, 3H), 7.80 (d, 1H), 10.5 (m, 1H), 13.05 (br s, 1H); EI-MS 456.2 (M+H); HPLC-Method C,  $R_t$  11.93 min.

Example 112 [2-(2-Chloro-phenyl)-6,7-dihydro-5H-cyclopentapyrimidin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-112): Prepared in 67% yield.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  2.18 (m, 2H), 2.89 (m, 2H), 3.02 (t, 2H), 7.24 (td, 1H), 7.42 (m, 2H), 7.49 (td, 1H), 7.52 (dd, 1H), 7.54 (d, 1H), 7.57 (dd, 1H), 10.50 (br. s, 1H), 13.06 (br. s, 1H); EI-MS 380.1 (M+1); HPLC-Method C,  $R_t$  9.68 min.

Example 113 (1H-Indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-6,7-dihydro-5H-cyclopentapyrimidin-4-yl]-amine (II-113): Prepared in 37% yield.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  2.65 (m, 2H), 2.85 (m, 2H), 2.99 (t, 2H), 7.02 (t, 1H), 7.32 (t, 1H), 7.47 (d, 1H), 7.55 (d, 1H), 7.68 (t, 1H),



7.74 (t, 1H), 7.80 (d, 1H), 10.37 (br. s, 1H), 12.91 (br. s, 1H); EI-MS 396.1 (M+H); HPLC-Method B, R<sub>t</sub> 9.88 min.

Example 114 (7-Fluoro-1H-indazol-3-yl)-[2-(2-

5 trifluoromethyl-phenyl)-6,7-dihydro-5H-

cyclopentapyrimidin-4-yl]-amine (II-114): Prepared in 40% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 2.15 (m, 2H), 2.87 (m, 2H), 2.97 (t, 2H), 6.99 (td, 1H), 7.17 (dd, 1H), 7.38 (d, 1H), 7.65 (m, 2H), 7.71 (t, 1H), 7.78 (d, 1H), 10.21 (br. s, 1H), 13.40 (br. s, 1H); EI-MS 414.1 (M+H); HPLC-Method C, R<sub>t</sub> 9.99 min.

Example 115 (5,7-Difluoro-1H-indazol-3-yl)-[2-(2-

15 trifluoromethyl-phenyl)-6,7-dihydro-5H-

cyclopentapyrimidin-4-yl]-amine (II-115): Prepared according to Method C in 52% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 2.16 (m, 2H), 2.89 (m, 2H), 2.97 (t, 2H), 7.19 (dd, 1H), 7.29 (td, 1H), 7.63 (t, 1H), 7.66 (d, 1H), 7.71 (t, 1H), 7.78 (d, 1H), 10.16 (br. s, 1H), 13.55 (br. s, 1H); EI-MS 432.1 (M+H); HPLC-Method C, R<sub>t</sub> 10.09 min.

Example 116 [2-(2-Chloro-phenyl)-6,7-dihydro-5H-

cyclopentapyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-116): Prepared in 56% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>)

25 δ 2.16 (m, 2H), 2.85 (m, 2H), 3.01 (t, 2H), 7.06 (t, 1H), 7.34 (t, 1H), 7.40 (t, 1H), 7.48 (m, 2H), 7.53 (d, 1H), 7.56 (d, 1H), 7.63 (d, 1H), 10.39 (br. s, 1H), 12.91 (s, 1H); EI-MS 362.1 (M+H); HPLC-Method A, R<sub>t</sub> 3.09 min.

30 Example 117 [2-(2-Chloro-phenyl)-6,7-dihydro-5H-

cyclopentapyrimidin-4-yl]-(7-fluoro-1H-indazol-3-yl)-amine (II-117): Prepared in 63% yield. <sup>1</sup>HNMR (500MHz, DMSO-d<sub>6</sub>) δ 2.15 (m, 2H), 2.87 (m, 2H), 3.00 (t, 2H), 7.01 (td, 1H), 7.19 (dd, 1H), 7.39 (t, 1H), 7.45 (m, 2H), 7.51

(d, 1H), 7.55 (d, 1H), 10.35 (br. s, 1H), 13.45 (br. s, 1H); EI-MS 380.1 (M+H); HPLC-Method A,  $R_t$   $R_t$  3.15 min.

Example 118 [2-(2-Chloro-phenyl)-6,7-dihydro-5H-

5 cyclopentapyrimidin-4-yl]-(5,7-difluoro-1H-indazol-3-yl)-amine (II-118): Prepared in 60% yield.  $^1\text{H}$ NMR (500MHz, DMSO-d<sub>6</sub>)  $\delta$  2.18 (m, 2H), 2.91 (m, 2H), 3.01 (t, 2H), 7.32 (t, 1H), 7.33 (td, 1H), 7.41 (t, 1H), 7.48 (t, 1H), 7.53 (d, 1H), 7.55 (dd, 1H), 10.35 (br. s, 1H), 13.45 (br. s, 10 1H); EI-MS 398.1 (M+H); HPLC-Method A,  $R_t$   $R_t$  3.24 min.

Example 119 (1H-Indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8,9,10-hexahydro-cyclooctapyrimidin-4-yl]-amine (II-119): Prepared in 36% yield.  $^1\text{H}$ NMR (500MHz,

15 DMSO-d<sub>6</sub>)  $\delta$  1.47 (m, 2H), 1.53 (m, 2H), 1.78 (m, 4H), 2.96 (m, 2H), 3.06 (t, 2H), 7.03 (t, 1H), 7.47 (t, 1H), 7.72 (d, 1H), 7.73 (d, 1H), 7.72 (m, 3H), 7.81 (d, 1H), 10.52 (m, 1H), 12.97 (br. s, 1H); EI-MS 438.2 (M+1); HPLC-Method A,  $R_t$  3.37 min.

20

Example 120 (7-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8,9,10-hexahydro-cyclooctapyrimidin-4-yl]-amine (II-120): Prepared in 40% yield.  $^1\text{H}$ NMR (500MHz, DMSO-d<sub>6</sub>)  $\delta$  1.46 (m, 2H), 1.52 (m, 25 2H), 1.77 (m, 4H), 2.94 (m, 2H), 3.04 (m, 2H), 7.00 (td, 1H), 7.17 (dd, 1H), 7.30 (d, 1H), 7.70 (m, 3H), 7.79 (d, 1H), 10.5 (m, 1H), 13.49 (br s, 1H); EI-MS 456.1 (M+H); HPLC-Method A,  $R_t$  3.43 min.

30 Example 121 (5,7-Difluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-5,6,7,8,9,10-hexahydro-cyclooctapyrimidin-4-yl]-amine (II-121): Prepared in 48% yield.  $^1\text{H}$ NMR (500MHz, DMSO-d<sub>6</sub>)  $\delta$  1.46 (m, 2H), 1.52 (m, 2H), 1.77 (m, 4H), 2.95 (m, 2H), 3.03 (m, 2H), 7.14 (d,

1H), 7.30 (t, 1H), 7.73 (m, 3H), 7.80 (d, 1H), 10.5 (m, 1H), 13.62 (br. s, 1H); EI-MS 475.1 (M+1); HPLC-Method A, R<sub>t</sub> 3.52 min.

5 Example 122 [6-Cyclohexyl-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-122):

Prepared in 45% yield. <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 1.30 (2H, m), 1.46 (2H, m), 1.65 (2H, m), 1.76 (2H, m), 1.91 (2H, m), 2.61 (1H, br m), 7.08 (1H, t, J=7.4 Hz), 7.27 (1H, d, J=8.0 Hz), 7.35 (1H, t, J= 7.1 Hz), 7.50 (1H, t, J=7.0 Hz), 7.58 (1H, t, J=7.4 Hz), 7.66 (3H, m), 7.72 (1H, d, J=7.8 Hz), 8.0 (1H, br), 9.87 (1H, br) ppm; HPLC-Method D, R<sub>t</sub> 3.57 min; LC-MS 438.17 (M+H)<sup>+</sup>

15 Example 123 [6-(2-Fluoro-phenyl)-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-123):

Prepared in 8% yield. <sup>1</sup>HNMR (500 MHz, CDCl<sub>3</sub>) δ 7.18 (3H, m), 7.37 (1H, m), 7.43 (1H, t, J=7.9 Hz), 7.51 (1H, d, J=7.9 Hz), 7.55 (1H, t, J=7.6 Hz), 7.65 (1H, t, J=7.4 Hz), 7.79 (1H, d, J=7.9 Hz), 7.85 (1H, d, J= 7.6 Hz), 8.19 (2H, m), 8.70 (1H, d, J= 8.5 Hz) ppm; HPLC-Method D, R<sub>t</sub> 4.93 min; LC-MS 450.13 (M+H)<sup>+</sup>

25 Example 124 (6-Fluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (II-124).

Prepared in DMF (87% yield) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.0 (s, 1H), 11.1 (s, br, 1H), 8.66 (d, 1H), 7.95 (t, 1H), 7.80 (d, 1H), 7.72 (m, 2H), 7.62 (m, 4H), 7.21 (dd, 1H), 6.84 (td, 1H) ppm. LC-MS (ES<sup>+</sup>) 424.15 (M+H); HPLC-Method A, R<sub>t</sub> 3.05 min.

Example 125 3-[2-(2-Trifluoromethyl-phenyl)-quinazolin-4-ylamino]-1H-indazole-5-carboxylic acid methyl ester (II-125): To a solution of compound II-79 (100 mg 0.21 mmol)

in DMF (2 mL) was added MeOH (1 mL), DIEA (54 uL, 0.31 mmol) and PdCl<sub>2</sub>(dppf) (4 mg, 0.005 mmol). The flask was flushed with CO three times and then charged with a CO balloon. The reaction mixture was heated at 80°C for 14 h then poured into water. The resulting precipitate was collected and washed with water. The crude product was then purified first by flash column (silica gel, 50% ethyl acetate in hexanes) then by preparative HPLC to afford II-125 (32%) as yellow solid. <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.3 (s, 1H), 11.3 (s, br, 1H), 8.70 (d, 1H), 8.36 (s, 1H), 7.97 (t, 1H), 7.82 (m, 2H), 7.71 (m, 3H), 7.58 (m, 2H), 7.51 (d, 1H), 3.75 (s, 3H) ppm; LC-MS (ES+) 464.13 (M+H); HPLC-Method A, R<sub>t</sub> 3.12 min.

Example 208 (5-Methyl-2H-pyrazol-3-yl)-[2-(2-naphthyl-1-yl)-quinazolin-4-yl]-amine (II-208): <sup>1</sup>HNMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.92 (s, 1H), 8.73 (m, 1H), 8.39 (m, 1H), 8.09 (m, 2H), 7.95 (m, 3H), 7.62 (m, 3H), 6.78 (s, 1H), 2.32 (s, 3H); MS 352.2 (M+H).

20

Example 209 [2-(2-Chloro-phenyl)-pyrido[2,3-d]pyrimidin-4-yl]-(7-fluoro-1H-indazol-3-yl)-amine (II-214): Prepared from 4-Chloro-2-(2-chloro-phenyl)-pyrido[2,3-d]pyrimidine (100 mg, 0.36mmol) and 7-Fluoro-1H-indazol-3-ylamine (108mg, 0.72mmol). Purification by preparative HPLC afforded the title compound as a yellow, di-TFA salt (93 mg, 46% yield). HPLC-Method A, R<sub>t</sub> 3.04 min; <sup>1</sup>H NMR (DMSO, 500 MHz): δ 13.67 (1H, s), 11.40-11.25 (1H, bs), 9.35-9.25 (2H, m), 7.95 (1H, m), 7.80-7.47 (5H, m), 7.35 (1H, m), 7.15 (1H, m); MS (m/z), MH<sup>+</sup> 391.1.

30

Example 210 [2-(2-Chloro-phenyl)-pyrido[2,3-d]pyrimidin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-215): Prepared from 4-Chloro-2-(2-chloro-phenyl)-pyrido[2,3-d]

pyrimidine (100 mg, 0.36mmol) and 5-Fluoro-1H-indazol-3-ylamine (108mg, 0.72mmol). Purification by preparative HPLC afforded the title compound as a yellow, di-TFA salt (45 mg, 22% yield). HPLC-Method A,  $R_t$  3.00 min;  $^1\text{H}$  NMR (DMSO, 500 MHz):  $\delta$  13.0 (1H, s), 10.90 (1H, bs), 9.15-9.05 (2H, m), 7.70 (1H, m), 7.60-7.30 (6H, m), 7.20 (1H, m); MS (m/z),  $\text{MH}^+$  391.1.

Example 211 [2-(2-Chloro-phenyl)-pyrido[2,3-d]pyrimidin-4-yl]-(5,7-difluoro-1H-indazol-3-yl)-amine (II-216): Prepared from 4-Chloro-2-(2-chloro-phenyl)-pyrido[2,3-d]pyrimidine (100 mg, 0.36mmol) and 7-Difluoro-1H-indazol-3-ylamine (112mg, 0.66mmol). Purification by preparative HPLC afforded the title compound as a yellow, di-TFA salt (130 mg, 62% yield). HPLC-Method A,  $R_t$  3.12 min;  $^1\text{H}$  NMR (DMSO, 500 MHz): 13.80-13.60 (1H, bs), 11.30-11.10 (1H, bs), 9.20-9.10 (2H, m), 7.80 (1H, m), 7.60-7.30 (6H, m); MS (m/z),  $\text{MH}^+$  409.1.

Example 212 [2-(2-Chloro-phenyl)-pyrido[3,4-d]pyrimidin-4-yl]-(1H-indazol-3-yl)-amine (II-217): Prepared from 4-Chloro-2-(2-chloro-phenyl)-pyrido[3,4-d]pyrimidine (100 mg, 0.36mmol) and 1H-indazol-3-ylamine (88mg, 0.66mmol). Purification by preparative HPLC afforded the title compound as a yellow, di-TFA salt (72 mg, 33% yield). HPLC-Method A,  $R_t$  3.21 min;  $^1\text{H}$  NMR (DMSO, 500 MHz):  $\delta$  12.95 (1H, s), 10.90 (1H, bs), 9.25 (1H, s), 8.75 (1H, m), 8.55 (1H, m), 7.65 (1H, m), 7.55 (1H, m), 7.50-7.30 (5H, m), 7.00 (1H, m); MS (m/z),  $\text{MH}^+$  373.1.

30

Example 213 [2-(2-Chloro-phenyl)-pyrido[3,4-d]pyrimidin-4-yl]-(7-fluoro-1H-indazol-3-yl)-amine (II-218): Prepared from 4-Chloro-2-(2-chloro-phenyl)-pyrido[3,4-d]pyrimidine (100 mg, 0.36mmol) and 7-Fluoro-1H-indazol-3-ylamine

(108mg, 0.72mmol). Purification by preparative HPLC afforded the title compound as a yellow, di-TFA salt (48.7 mg, 22% yield). HPLC-Method A,  $R_t$  3.35 min;  $^1\text{H}$  NMR (DMSO, 500 MHz):  $\delta$  12.95 (1H, s), 10.90 (1H, bs), 9.25 (1H, s), 8.75 (1H, m), 8.55 (1H, m), 7.70-7.35 (5H, m), 7.25 (1H, m), 6.95 (1H, m); MS (m/z),  $\text{MH}^+$  391.08.

Example 214 [2-(2-Chloro-phenyl)-pyrido[3,4-d]pyrimidin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (II-219): Prepared from 4-chloro-2-(2-chloro-5-fluoro-1H-indazol-3-yl)amine (108mg, 0.72mmol). Purification by preparative HPLC afforded the title compound as a yellow, di-TFA salt (57.2 mg, 26% yield). HPLC-Method A,  $R_t$  3.27 min;  $^1\text{H}$  NMR (DMSO, 500 MHz):  $\delta$  13.05 (1H, s), 10.95 (1H, s), 9.25 (1H, s), 8.75 (1H, m), 8.55 (1H, m), 7.60 (1H, m), 7.55 (1H, m), 7.50-7.30 (5H, m), 7.25 (1H, m); MS (m/z),  $\text{MH}^+$  391.1.

Example 215 [2-(2-Chloro-phenyl)-pyrido[3,4-d]pyrimidin-4-yl]-(5,7-difluoro-1H-indazol-3-yl)-amine (II-220): Prepared from 4-chloro-2-(2-chloro-7-difluoro-1H-indazol-3-yl)amine (112mg, 0.66mmol). Purification by preparative HPLC afforded the title compound as a yellow, di-TFA salt (57.2 mg, 26% yield). HPLC-Method A,  $R_t$  3.45 min;  $^1\text{H}$  NMR (DMSO, 500 MHz):  $\delta$  13.65 (1H, s), 11.0 (1H, s), 9.25 (1H, s), 8.80 (1H, m), 8.50 (1H, m), 7.60 (1H, m), 7.55 (1H, m), 7.50-7.30 (5H, m); MS (m/z),  $\text{MH}^+$  409.1.

Example 216 6-Fluoro-1H-indazol-3-ylamine (A1):  $^1\text{H}$ NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.4 (s, 1H), 7.68 (dd, 1H), 6.95 (dd, 1H), 6.75 (td, 1H), 5.45 (s, 2H) ppm; LC-MS (ES+) 152.03 (M+H); HPLC-Method A,  $R_t$  2.00 min.

Example 217 5-Fluoro-1H-indazol-3-ylamine (A2): <sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.3 (s, 1H), 7.43 (d, 1H), 7.22 (m, 1H), 7.08 (m, 1H), 5.29 (s, 2H) ppm; LC-MS (ES+) 152.01 (M+H); HPLC-Method A, R<sub>t</sub> 1.93 min.

5

Example 218 5,7-Difluoro-1H-indazol-3-yl-amine (A3): <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD) δ 7.22 (dd, J=2.0, 8.45Hz, 1H), 7.04-6.87 (m, 1H); LC-MS (ES+) 169.95 (M+H); HPLC-Method C, R<sub>t</sub> 2.94 min

10

Example 219 7-Fluoro-1H-indazol-3-ylamine (A4): <sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.8 (s, 1H), 7.42 (d, 1H), 6.97 (m, 1H), 6.78 (m, 1H), 5.40 (s, 2H) ppm; LCMS (ES+) 152.01 (M+H); HPLC-Method A, R<sub>t</sub> 2.00 min.

15

Example 220 7-Fluoro-6-trifluoromethyl-1H-indazol-3-ylamine (A5): <sup>1</sup>H-NMR (500 MHz, DMSO) δ 12.5 (s, 1H), 7.75 (d, 1H), 7.25 (m, 1H), 5.85 (m, 1H) ppm; MS (FIA) 220.0 (M+H); HPLC-Method A, R<sub>t</sub> 2.899 min.

20

Example 221 6-Bromo-1H-indazol-3-ylamine (A6): <sup>1</sup>H-NMR (500 MHz, DMSO) δ 11.5 (s, 1H), 7.65 (d, 1H), 7.40 (s, 1H), 7.00 (d, 1H), 5.45 (br s, 1H) ppm; MS (FIA) 213.8 (M+H); HPLC-Method A, R<sub>t</sub> 2.441 min.

25

Example 222 4-Fluoro-1H-indazol-3-ylamine (A7): <sup>1</sup>H-NMR (500 MHz, DMSO) δ 11.7 (s, 1H), 7.17 (m, 1H), 7.05 (d, 1H), 6.7 (br, 1H), 6.60 (dd, 1H), 5.20 (br s, 2H) ppm; MS (FIA) 152.0 (M+H); Method A, R<sub>t</sub> 2.256 min.

30

Example 223 5-Bromo-1H-indazol-3-ylamine (A8): <sup>1</sup>H-NMR (500 MHz, DMSO) δ 11.55 (br s, 1H), 7.95 (s, 1H), 7.30 (d,

1H), 7.20 (d, 1H), 5.45 (br s, 2H) ppm; MS (FIA) 213.8 (M+H); Method A,  $R_t$  2.451 min.

5 Example 224 5-Nitro-1H-indazol-3-ylamine (A9):  $^1\text{H}$ -NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.00 (s, 1H), 8.20 (d, 1H), 7.45 (d, 1H), 6.15 (br s, 1H) ppm; Method A,  $R_t$  2.184 min

10 Example 225 4-Pyrrol-1-yl-1H-indazol-3-ylamine (A10):  $^1\text{H}$ -NMR (500 MHz, DMSO)  $\delta$  7.20 (s, 2H), 7.00 (s, 2H), 6.75 (m, 1H), 6.25 (s, 2H), 4.30 (d, 1H) ppm; Method A,  $R_t$  2.625 min.

15 Example 226 4-Chloro-5,6-dimethyl-2-(2-trifluoromethyl-phenyl)-pyrimidine (B1): Prepared to afford a colorless oil in 75% yield.  $^1\text{H}$ -NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (d,  $J=7.8$  Hz, 1H), 7.64 (d,  $J=7.6$  Hz, 1H), 7.55 (t,  $J=7.6$  Hz, 1H), 7.48 (t,  $J=7.5$  Hz, 1H), 2.54 (s, 3H), 2.36 (s, 3H) ppm; MS (FIA) 287.0 (M+H); HPLC-Method A,  $R_t$  3.891 min.

20 Example 227 4-Chloro-2-(2-chloro-phenyl)-5,6-dimethyl-pyrimidine (B2): Prepared to afford a yellow-orange oil in 71% yield.  $^1\text{H}$ -NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.73 (m, 1H), 7.52 (m, 1H), 7.39 (m, 2H), 2.66 (s, 3H), 2.45 (s, 3H) ppm; MS (FIA) 253.0 (M+H); HPLC-Method A,  $R_t$  4.156 min.

25 Example 228 4-Chloro-6-methyl-2-(2-trifluoromethyl-phenyl)-pyrimidine (B3): Prepared to afford a pale yellow oil in 68% yield.  $^1\text{H}$ -NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.72 (d,  $J=7.8$  Hz, 1H), 7.65 (d,  $J=7.9$  Hz, 1H), 7.57 (t,  $J=7.5$  Hz, 1H), 7.52 (t,  $J=7.8$  Hz, 1H), 7.16 (s, 1H), 2.54 (s, 3H) ppm; MS (FIA) 273.0 (M+H); HPLC-Method A,  $R_t$  3.746 min.

30



Example 229 4-Chloro-6-cyclohexyl-2-(2-trifluoromethyl-phenyl)-pyrimidine (B4): Prepared to afford a yellow oil in 22% yield.  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.70 (m, 2H), 7.57 (t,  $J=7.5$  Hz, 1H), 7.50 (t,  $J=7.5$  Hz, 1H), 7.19 (s, 1H), 2.65 (m, 1H), 1.9 (m, 2H), 1.8 (m, 2H), 1.5 (m, 2H), 1.3 (m, 2H), 1.2 (m, 2H) ppm; MS (FIA) 341.0 (M+H).

Example 230 4-Chloro-6-phenyl-2-(2-trifluoromethyl-phenyl)-pyrimidine (B5): Prepared to afford a yellow oil in 53% yield.  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.08 (dd,  $J=7.9$ , 1.6 Hz, 2H), 7.80 (d,  $J=7.6$  Hz, 1H), 7.77 (d,  $J=7.8$  Hz, 1H), 7.67 (s, 1H), 7.61 (t,  $J=7.5$  Hz, 1H), 7.54 (t,  $J=7.6$  Hz, 1H), 7.47 (m, 3H) ppm; MS (FIA) 335.0 (M+H); HPLC-Method A,  $R_t$  4.393 min.

15

Example 231 4-Chloro-2-(2,4-dichloro-phenyl)-5,6-dimethyl-pyrimidine (B6): Prepared to afford a white solid in 91% yield.  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.62 (d,  $J=8.3$  Hz, 1H), 7.43 (d,  $J=7.0$  Hz, 1H), 7.27 (dd,  $J=8.3$ , 2.0 Hz, 1H), 2.55 (s, 3H), 2.35 (s, 3H) ppm; MS (FIA) 287, 289 (M+H); HPLC-Method A,  $R_t$  4.140 min.

20

Example 232 4-Chloro-6-(2-chloro-phenyl)-2-(2-trifluoromethyl-phenyl)-pyrimidine (B7): Prepared to afford a yellow oil in 52% yield.  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.75 (m, 3H), 7.65 (m, 2H), 7.53 (m, 1H), 7.44 (m, 1H), 7.36 (m, 2H) ppm; MS (FIA) 369.1 (M+H); HPLC-Method A,  $R_t$  4.426 min.

25

Example 233 4-Chloro-6-(2-fluoro-phenyl)-2-(2-trifluoromethyl-phenyl)-pyrimidine (B8): Prepared to afford a yellow oil in 95% yield.  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.24 (t,  $J=7.9$  Hz, 1H), 7.84 (s, 1H), 7.78 (d,  $J=7.7$

30

Hz, 1H), 7.76 (d, J=8.0 Hz, 1H), 7.60 (t, J=7.5 Hz, 1H), 7.53 (t, J=7.6 Hz, 1H), 7.43 (m, 1H), 7.23 (t, J=7.6 Hz, 1H), 7.13 (m, 1H) ppm; MS (FIA) 353.0 (M+H).

- 5 Example 234 4-Chloro-6-pyridin-2-yl-2-(2-trifluoromethyl-phenyl)-pyrimidine (B9): Prepared to afford a pale yellow solid in 50% yield. <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ 8.68 (m, 1H), 8.48 (dd, J=7.9, 0.8 Hz, 1H), 8.38 (d, J=2.3 Hz, 1H), 7.84 (m, 3H), 7.62 (t, J=7.6 Hz, 1H), 7.55 (t, J=7.6 Hz, 1H), 7.38 (m, 1H) ppm; MS (FIA) 336.0 (M+H); HPLC-Method A, R<sub>t</sub> 4.575 min.

- Example 235 6-Benzyl-4-chloro-2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydro-pyrido[4,3-d]pyrimidine (B10):  
15 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ 7.70 (d, 1H), 7.62 (d, 1H), 7.55 (t, 1H), 7.48 (t, 1H), 7.32 (m, 4H), 7.25 (m, 1H), 3.74 (s, 2H), 3.66 (s, 2H), 2.99 (t, 2H), 2.80 (t, 2H) ppm; LCMS (ES+) 404.17 (M+H); HPLC-Method A, R<sub>t</sub> 3.18 min.

- 20 Example 236 7-Benzyl-4-chloro-2-(2-trifluoromethyl-phenyl)-5,6,7,8-tetrahydro-pyrido[3,4-d]pyrimidine (B11):  
<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ 7.69 (d, 1H), 7.60 (d, 1H), 7.54 (t, 1H), 7.47 (t, 1H), 7.28 (m, 4H), 7.20 (m, 1H), 3.68 (s, 2H), 3.67 (s, 2H), 2.86 (t, 2H), 2.79 (t, 2H) ppm. MS  
25 (ES+) 404.18 (M+H); HPLC-Method A, R<sub>t</sub> 3.12 min.

- Example 237 4-Chloro-2-(4-fluoro-2-trifluoromethyl-phenyl)-quinazoline (B12): <sup>1</sup>H-NMR (500MHz, CD<sub>3</sub>OD) δ 8.43 (d, J=8.1Hz, 1H), 8.20-8.05 (m, 2H), 8.05-7.82 (m, 2H),  
30 7.71-7.51 (m, 2H). LC-MS (ES+) 327.09 (M+H). HPLC-Method D, R<sub>t</sub> 4.56 min.

Example 238 4-Chloro-2-(2-chloro-5-trifluoromethyl-phenyl)-quinazoline (B13): LC-MS (ES+) 342.97 (M+H).  
HPLC-Method D,  $R_t$  4.91 min.

5 Example 239 4-Chloro-2-(2-chloro-4-nitro-phenyl)-quinazoline (B14): LC-MS (ES+) 319.98 (M+H). HPLC-Method D,  $R_t$  4.45 min.

Example 240 4-Chloro-2-(2-trifluoromethyl-phenyl)-quinazoline (B15): Prepared in 57% yield. White solid.  
10  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  7.79 (t, 1H), 7.86 (t, 1H), 7.94 (m, 3H), 8.15 (dd, 1H), 8.20 (td, 1H), 8.37 (m, 1H); EI-MS 308.9 (M).

15 Example 241 4-Chloro-2-(2-trifluoromethyl-phenyl)-6,7-dihydro-5H-cyclopentapyrimidine (B16): Prepared in 22% yield.  $^1\text{H}$ NMR (500MHz, DMSO- $d_6$ )  $\delta$  2.19 (m, H), 3.01 (t, 2H), 3.08 (t, 2H), 7.49 (t, 1H), 7.55 (t, 1H), 7.62 (d, 1H), 7.71 (d, 1H). EI-MS 299.0 (M+H).

20

Example 242 4-Chloro-2-(2-chloro-phenyl)-6,7,8,9-tetrahydro-5H-cycloheptapyrimidine (B17): Prepared according to Method C in 82% yield to afford a white solid.  $^1\text{H}$ NMR (500MHz,  $\text{CDCl}_3$ )  $\delta$  1.67 (m 4H), 1.87 (m 2H),  
25 3.02 (m 4H), 7.28 (m, 2H), 7.40 (m, 1H), 7.65 (m, 1H); EI-MS 293.0 (M+1).

Example 243 4-Chloro-2-(2-trifluoromethyl-phenyl)-5,6,7,8,9,10-hexahydro-cyclooctapyrimidine (B18):  
30 Prepared in 38% yield to afford a brown oil.  $^1\text{H}$ NMR (500MHz,  $\text{CDCl}_3$ )  $\delta$  1.35 (m 2H), 1.41 (m 2H), 1.76 (m 4H), 2.96 (m, 4H), 7.48 (t, 1H), 7.56 (t, 1H), 7.66 (d, 1H), 7.70 (d, 1H); EI-MS 341.0 (M+1).

Example 244 4-Chloro-8-methoxy-2-(2-trifluoromethyl-phenyl)-quinazoline (B19): Prepared from 8-methoxy-2-(2-trifluoromethyl-phenyl)-3H-quinazolin-4-one (1.0g, 3.12mmol), triethylamine hydrochloride (472mg, 3.43mmol), and POCl<sub>3</sub>. Purification by flash chromatography afforded a white solid (89% yield). HPLC-Method A, R<sub>t</sub> 4.10 min, (98%), MS (m/z) 258.08 (M+H).

Example 245 2-(4-Chloro-quinazolin-2-yl)-benzonitrile (B20): Prepared to afford a yellow solid in 1.5% yield. <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) δ 8.47 (d, 1H), 8.24 (d, 1H), 8.16 (d, 1H), 8.07 (impurity), 7.94 (t, 1H), 7.92 (impurity), 7.86 (d, 1H), 7.68 (m, 2H), 7.65 (impurity), 7.54 (impurity), 7.49 (t, 1H), 4.2 (impurity), 1.05 (impurity) ppm; MS (LC/MS) 266.05 (M+H); HPLC-Method A, R<sub>t</sub> 3.88 min.

Example 246 6-Methyl-2-(2-trifluoromethyl-phenyl)-3H-pyrimidin-4-one (D3): Prepared to afford a yellow solid in 50% yield. <sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.7 (br s, 1H), 7.9 (m, 1H), 7.8 (m, 2H), 7.7 (m, 1H), 6.3 (s, 1H), 2.21 (s, 3H) ppm; MS (FIA) 255.0 (M+H); HPLC-Method A, R<sub>t</sub> 2.578 min.

Example 247 6-Cyclohexyl-2-(2-trifluoromethyl-phenyl)-3H-pyrimidin-4-one (D4): Prepared to afford an off-white solid in 54% yield. <sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.9 (br s, 1H), 7.9 (m, 4H), 6.3 (s, 1H), 2.5 (m, 1H), 1.9 (m, 5H), 1.4 (m, 5H) ppm; MS (FIA) 323.1 (M+H); HPLC-Method A, R<sub>t</sub> 3.842 min.

Example 248 2-(2-Chloro-5-trifluoromethyl-phenyl)-3H-quinazolin-4-one (D10): <sup>1</sup>H-NMR (500MHz, CD<sub>3</sub>OD) δ 8.32-8.25 (m, 1H), 8.01 (s, 1H), 7.91-7.72 (m, 1H), 7.66-7.55 (m,

1H). LC-MS (ES+) 325.01 (M+H). HPLC-Method D,  $R_t$  3.29 min.

Example 249 2-(4-Fluoro-2-trifluoromethyl-phenyl)-3H-

5 quinazolin-4-one (D14):  $^1\text{H}$ NMR (500MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.28 (d, 8.0Hz, 1H), 7.94-7.84 (m, 1H), 7.84-7.77 (m, 1H), 7.76-7.67 (m, 2H), 7.65-7.53 (m, 2H). LC-MS (ES+) 309.06 (M+H). HPLC-Method D,  $R_t$  2.88 min.

10 Example 250 2-(4-Nitro-2-chloro-phenyl)-3H-quinazolin-4-one (D15): LC-MS (ES+) 302.03 (M+H). HPLC-Method D,  $R_t$  2.81 min.

Example 251 2-(5-Fluoro-2-trifluoromethyl-phenyl)-3H-

15 quinazolin-4-one (D17):  $^1\text{H}$ NMR (500MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.28 (d,  $R_t$  J=8.05Hz, 1H), 7.96 (dd, J=5.05, 8.55Hz, 1H), 7.89 (t, J=7.9Hz, 1H), 7.78-7.69 (m, 1H), 7.66-7.46 (m, 3H). LC-MS (ES+) 309.14 (M+H). HPLC-Method D,  $R_t$  2.90 min.

20 Example 252 (1H-Indazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-1): Prepared by Method A in DMF to afford 70 mg (50% yield) as pale yellow solid.  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.1 (s, br, 1H), 8.48 (d, 1H), 7.91 (d, 2H), 7.76 (br, 2H), 7.45 (m, 2H), 7.36 (d, 1H), 7.20 (m, 4H),  
25 6.86 (t, 1H) ppm. MS (ES+) 338.07 (M+H); (ES-) 336.11 (M-H); HPLC-Method A,  $R_t$  2.88 min.

Example 253 (5-Methyl-2H-pyrazol-3-yl)-(2-phenyl-5,6,7,8-tetrahydroquinazolin-4-yl)-amine (III-7): Prepared

30 according to Method A.  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.1 (s, br, 1H), 8.70 (s, br, 1H), 8.37 (d, J = 6.7 Hz, 2H), 7.54 (m, 3H), 6.67 (s, 1H), 2.82 (m, 2H), 2.68 (m, 2H), 2.37 (s, 3H), 1.90 (s, br, 4H); MS 306.1 (M+H).

Example 254 (5-Methyl-2H-pyrazol-3-yl)-(2-phenyl-6,7,8,9-tetrahydro-5H-cycloheptapyrimidin-4-yl)-amine (III-8): MS 320.48 (M+H); HPLC-Method E,  $R_t$  1.124 min.

5 Example 255 (5-Methyl-2H-pyrazol-3-yl)-(2-pyridin-4-yl-quinazolin-4-yl)-amine (III-9): Yellow solid, mp 286-289°C,  $^1\text{H}$  NMR (DMSO)  $\delta$  2.35 (3H, s), 6.76 (1H, s), 7.61 (1H, m), 7.89 (2H, m), 8.32 (2H, d), 8.70 (1H, d), 8.78 (2H, d), 10.56 (1H, br s), 12.30 (1H, br s); IR (solid)  
10 1620, 1598, 1571, 1554, 1483, 1413, 1370, 1328; MS 303.2 (M+H)<sup>+</sup>

Example 256 (7-Chloro-2-pyridin-4-yl-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (III-28):  $^1\text{H}$  NMR (DMSO-d<sub>6</sub>)  $\delta$   
15 2.35 (3H, s), 6.75 (1H, s), 7.65 (1H, d), 7.93 (1H, s), 8.30 (2H, d), 8.73 (1H, d), 8.79 (2H, d), 10.69 (1H, s), 12.33 (1H, s); MS m/z 337.2 (M+H)<sup>+</sup>.

Example 257 (6-Chloro-2-pyridin-4-yl-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (III-29):  $^1\text{H}$  NMR (DMSO-d<sub>6</sub>)  $\delta$   
20 2.31 (3H, s), 6.74 (1H, s), 7.89 (1H, s), 8.30 (2H, d), 8.80 (2H, d), 8.91 (1H, s), 10.63 (1H, s), 12.29 (1H, s); MS 337.2 (M+H)<sup>+</sup>.

25 Example 258 (2-Cyclohexyl-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (III-30):  $^1\text{H}$  NMR (DMSO)  $\delta$  2.35 (3H, s), 1.70 (3H, m), 1.87 (2H, d), 1.99 (2H, d), 2.95 (1H, t), 6.72 (1H, s), 7.75 (1H, d), 7.88 (1H, s), 7.96 (1H, s), 8.83 (1H, s), 11.95 (1H, s), 12.70 (1H, s); MS 308.4  
30 (M+H)<sup>+</sup>.

Example 259 (5-Methyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-31): mp 246°C;  $^1\text{H}$  NMR (400MHz)

$\delta$  2.35 (3H, s), 6.70 (1H, br s), 7.51-7.57 (4H, m), 7.83-7.84 (2H, d), 8.47-8.50 (2H, d), 8.65 (1H, d), 10.4 (1H, s), 12.2 (1H, bs); IR (solid) 3696, 3680, 2972, 2922, 2865; MS 302.1 (M+H)+.

5

Example 260 [2-(4-Iodophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-32):  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  2.34 (3H, s), 6.72 (1H, s), 7.56 (1H, d), 7.84 (2H, d), 7.93 (2H, d), 8.23 (2H, d), 8.65 (1H, s), 10.44 (1H, s), 12.24 (1H, s); MS 428.5 (M+H)+.

10

Example 261 [2-(3,4-Dichlorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-33): A suspension of 2-(3,4-dichloro-phenyl)-3H-quinazolin-4-one (1g, 3.43 mmol) in phosphorus oxychloride (4 mL) was stirred at 110°C for 3 hours. The solvent was removed by evaporation and the residue is treated carefully with cold aqueous, saturated NaHCO<sub>3</sub>. The resulting solid was collected by filtration and washed with ether to afford 4-chloro-2-(3,5-dichloro-phenyl)-quinazoline as a white solid (993 mg, 93%). To the above compound (400mg, 1.29 mmol) in THF (30 mL) was added 3-amino-5-methyl pyrazole (396 mg, 2.58 mmol) and the resulting mixture heated at 65°C overnight. The solvents were evaporated and the residue triturated with ethyl acetate, filtered, and washed with the minimum amount of ethanol to afford compound III-33 as a white solid (311 mg 65%): mp 274°C;  $^1\text{H}$  NMR (DMSO)  $\delta$  2.34 (3H, s), 6.69 (1H, s), 7.60 (1H, m), 7.84 (1H, d), 7.96 (2H, d), 8.39 (1H, dd), 8.60 (1H, d), 8.65 (1H, d), 10.51 (1H, s), 12.30 (1H, s); IR (solid) 1619, 1600, 1559, 1528, 1476, 1449, 1376, 1352, 797, 764, 738; MS 370.5 (M+H)+.

15

20

25

30

Example 262 [2-(4-Bromophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-34): mp 262-265°C; <sup>1</sup>H NMR (DMSO) δ 2.34 (3S, s), 6.73 (1H, s), 7.55 (1H, m), 7.74 (2H, d), 7.83 (2H, m), 8.40 (2H, d), 8.65 (1H, d), 10.44 (1H, s), 12.25 (1H, s); IR (solid) 1603, 1579, 1546, 1484, 1408, 1365; MS 380.1/382.1 (M+H)+.

Example 263 [2-(4-Chlorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-35): mp >300°C; <sup>1</sup>H NMR (DMSO) δ 2.34 (3H, s), 6.74 (1H, s), 7.53-7.62 (3H, m), 7.84 (2H, d), 8.47 (2H, d), 8.65 (1H, d), 10.44 (1H, s), 12.26 (1H, s); IR (solid) 1628, 1608, 1584, 1546, 1489, 1408, 1369, 1169; MS 336.2 (M+H)+.

Example 264 [2-(3,5-Dichlorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-36): mp 228°C; <sup>1</sup>H NMR (DMSO) δ 2.34 (3H, s), 6.69 (1H, s), 7.96 (1H, d), 8.21 (3H, m), 8.56 (1H, d), 8.60 (2H, d), 10.51 (1H, s), 12.30 (1H, s); IR (solid) 1546, 1331, 802, 763, 729, 658, 652; MS 370.5 (M+H)+.

Example 265 [2-(4-Cyanophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-37): mp 263°C; <sup>1</sup>H NMR (DMSO) δ 2.34 (3H, s), 6.72 (1H, s), 7.61 (1H, d), 7.88 (2H, s), 8.04 (2H, d), 8.63 (2H, d), 8.67 (1H, s), 10.52 (1H, s), 12.27 (1H, s); IR (solid) 1739, 1436, 1366, 1229, 1217; MS 327.2 (M+H)+.

Example 266 [2-(3-Iodophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-38): mp 234-235°C; <sup>1</sup>H NMR (DMSO) δ 2.35 (3H, s), 6.73 (1H, s), 7.35 (1H, m), 7.56 (1H, m), 7.85 (3H, m), 8.47 (1H, m), 8.65 (1H, m), 8.86



(1H, s), 10.49 (1H, s), 12.28 (1H, br s); IR (solid) 1560, 1541, 1469, 1360; MS 428.1 (M+H)+.

Example 267 [2-(4-Ethylsulfanylphenyl)-quinazolin-4-yl]-

5 (5-methyl-2H-pyrazol-3-yl)-amine (III-39): mp 229-231°C; <sup>1</sup>H NMR (DMSO) δ 1.29 (3H, t), 2.35 (3H, s), 3.07 (2H, q), 6.76 (1H, s), 7.43 (2H, d), 7.51 (1H, m), 7.81 (2H, m), 8.41 (2H, d), 8.64 (1H, d), 10.38 (1H, s), 12.24 (1H, br s); IR (solid) 1587, 1574, 1555, 1531, 1484, 1412, 1369; 10 MS 362.1 (M+H)+.

Example 268 (5-Cyclopropyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-40): mp 218-219°C; <sup>1</sup>H NMR

(DMSO-d<sub>6</sub>) δ 0.70-0.80 (2H, m), 0.90-1.00 (2H, m), 6.70 15 (1H, s), 7.45-7.55 (4H, m), 7.80-7.85 (2H, m), 8.45-8.55 (2H, m), 8.65 (1H, d), 10.40 (1H, s), 12.27 (1H, s); IR (solid) 1624, 1605, 1591, 1572, 1561, 1533, 1479, 1439, 1419, 1361, 1327, 997, 828, 803, 780, 762, 710; MS 328.2 (M+H)+.

20

Example 269 [2-(4-tert-Butylphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-41): mp >300°C; <sup>1</sup>H NMR

(DMSO-d<sub>6</sub>) δ 1.35 (9H, s), 2.34 (3H, s), 6.79 (1H, s), 7.55 (3H, d), 7.85 (2H, d), 8.39 (2H, d), 8.62 (1H, d), 25 10.35 (1H, s), 12.22 (1H, s); IR (solid) 1603, 1599, 1577, 1561, 1535, 1481, 1409, 1371, 1359, 998, 841, 825, 766, 757; MS 358.3 (M+H)+.

Example 270 [2-(4-Chlorophenyl)-quinazolin-4-yl]-(5-

30 cyclopropyl-2H-pyrazol-3-yl)-amine (III-42): <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.77 (4H, br m), 2.05 (1H, m), 6.59 (1H, s), 7.60 (1H, d), 7.85 (2H, d), 7.91 (2H, d), 8.22 (2H, d), 8.65 (1H, s), 10.51 (1H, s), 12.33 (1H, s); MS 362.1 (M+H)+.

Example 271 (2-Benzo[1,3]dioxol-5-yl-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (III-43):  $^1\text{H}$  NMR (DMSO)  $\delta$  2.33 (3H, s), 6.13 (2H, s), 6.78 (1H, s), 7.11 (1H, d), 7.80 (1H, t), 7.94 (1H, s), 8.09 (3H, m), 8.25 (1H, d), 10.34 (1H, s), 12.21 (1H, s); MS 346.5 (M+H) $^+$ .

Example 272 [2-(4-Dimethylaminophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-44):  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  2.02 (6H, s), 2.39 (3H, s), 6.83 (1H, s), 7.71 (1H, d), 7.98 (2H, s), 8.04 (2H, d), 8.33 (2H, d), 8.67 (1H, s), 11.82 (1H, s), 12.72 (1H, s); MS 345.3 (M+H) $^+$ .

Example 273 [2-(3-Methoxyphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-45): mp 226°C;  $^1\text{H}$  NMR (DMSO)  $\delta$  2.34 (3H, s), 3.92 (3H, s), 6.72 (1H, s), 7.21 (1H, d), 7.57 (1H, t), 7.79 (1H, t), 8.02 (3H, m), 8.14 (1H, s), 8.79 (1H, d), 10.39 (1H, s), 12.22 (1H, s); IR (solid) 1599, 1572, 1538, 1478, 1427, 1359, 833, 761, 661; MS 332.2 (M+H) $^+$ .

Example 275 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(3,4-dichlorophenyl)-quinazolin-4-yl]-amine (III-46):  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.86 (2H, d), 1.02 (2H, d), 1.69 (1H, m), 6.56 (1H, s), 7.57 (1H, d), 7.84 (4H, m), 8.40 (1H, d), 8.58 (1H, s), 8.64 (1H, s), 10.53 (1H, s), 12.36 (1H, s); MS 396.0 (M+H) $^+$ .

Example 276 (2-Biphenyl-4-yl-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (III-47): To a mixture of [2-(4-bromo-phenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-34) (196 mg, 0.51 mmol) and phenylboronic acid (75 mg, 0.62 mmol) in THF:water (1:1, 4 mL) was added  $\text{Na}_2\text{CO}_3$  (219 mg, 2.06 mmol), triphenylphosphine (9mg, 1/15 mol%) and palladium acetate (1 mg, 1:135 mol%). The

resulting mixture was heated at 80°C overnight, the solvents were evaporated and the residue purified by flash chromatography (gradient of dichloromethane:MeOH) to afford III-21 as a yellow solid (99 mg, 51%): <sup>1</sup>H NMR (DMSO) δ 2.37 (3H, s), 6.82 (1H, s), 7.39-7.57 (4H, m), 7.73-7.87 (6H, m), 8.57 (2H, d), 8.67 (1H, d), 10.42 (1H, s), 12.27 (1H, s); MS 378.2 (M+H)<sup>+</sup>.

Example 277 [2-(4-Ethynylphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-48): To a mixture of [2-(4-bromo-phenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-34) (114 mg, 0.3 mmol), and trimethylsilylacetylene (147 mg, 1.5 mmol) in DMF (2 mL) was added CuI (1.1 mg, 1:50 mol%), Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (4.2 mg, 1:50 mol%) and triethylamine (121 mg, 0.36 mmol). The resulting mixture was heated at 120°C overnight and the solvent evaporated. The residue was triturated in ethyl acetate and the resulting precipitate collected by filtration. The collected solid was suspended in THF (3 mL) and TBAF (1M in THF, 1.1eq) was added. The reaction mixture was stirred at room temperature for 2 hours and the solvent evaporated. The residue was purified by flash chromatography (silica gel, gradient of DCM:MeOH) to afford III-48 as a white solid (68 mg, 70%): <sup>1</sup>H NMR (DMSO) δ 2.34 (3H, s), 4.36 (1H, s), 6.74 (1H, s), 7.55 (1H, m), 7.65 (2H, d), 7.84 (2H, m), 8.47 (2H, d), 8.65 (1H, d), 10.43 (1H, s), 12.24 (1H, s); MS 326.1 (M+H)<sup>+</sup>.

Example 278 [2-(3-Ethynylphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-49): mp 204-207°C; <sup>1</sup>H NMR (DMSO) δ 2.34 (3H, s), 4.28 (1H, s), 6.74 (1H, s), 7.55-7.63 (3H, m), 7.83-7.87 (2H, m), 8.49 (1H, d), 8.57 (1H, s), 8.65 (1H, d), 10.46 (1H, s), 12.27 (1H, s); IR

(solid) 1598, 1574, 1541, 1489, 1474, 1422, 1365; MS  
326.1 (M+H)+.

Example 279 [2-(3-Methylphenyl)-quinazolin-4-yl]-(5-

5 methyl-2H-pyrazol-3-yl)-amine (III-50): A suspension of  
1H-quinazoline-2,4-dione (10.0 g, 61.7 mmol) in POCl<sub>3</sub> (60  
mL, 644 mmol) and N,N-dimethylaniline (8mL, 63.1 mmol)  
was heated under reflux for 2 h. The excess POCl<sub>3</sub> was  
removed *in vacuo*, the residue poured into ice, and the  
10 resulting precipitate collected by filtration. The crude  
solid product 2,4-dichloro-quinazoline (6.5 g, 53% yield)  
was washed with water and dried under vacuum for next  
step use without further purification. To a solution of  
the 2,4-dichloro-quinazoline (3.3 g, 16.6 mmol) in  
15 anhydrous ethanol (150 mL) was added 5-methyl-1H-pyrazol-  
3-yl amine (3.2 g, 32.9 mmol) and the resulting mixture  
was stirred at room temperature for 4 hours. The  
resulting precipitate was collected by filtration, washed  
with ethanol, and dried under vacuum to afford 4.0 g (93%  
20 yield) of (2-chloro-quinazolin-4-yl)-(5-methyl-1H-  
pyrazol-3-yl)-amine which was used in the next step  
without further purification. To a solution of the (2-  
chloro-quinazolin-4-yl)-(5-methyl-1H-pyrazol-3-yl)-amine  
(50 mg, 0.19 mmol) in DMF (1.0 mL) was added *m*-tolyl  
25 boronic acid (0.38 mmol), 2M Na<sub>2</sub>CO<sub>3</sub> (0.96 mmol), and tri-  
t-butylphosphine (0.19 mmol). The flask was flushed with  
nitrogen and the catalyst PdCl<sub>2</sub>(dppf) (0.011 mmol) added  
in one portion. The reaction mixture was then heated at  
80°C for 10 hours, cooled to room temperature, and poured  
30 into water (2 mL). The resulting precipitate was  
collected by filtration, washed with water, and purified  
by HPLC to afford III-50 as a pale yellow solid (61mg,  
75%): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.3 (br s, 1H), 10.4  
(br s, 1H), 8.75 (d, 1H), 8.30 (s, 1H), 8.25 (d, 1H),

7.78 (s, 2H), 7.55 (m, 1H), 7.45 (m, 1H), 7.35 (m, 1H),  
6.80 (s, 1H), 2.47 (s, 3H), 2.30 (s, 3H); MS 316.1 (M+H).

5 Example 280 [2-(3,5-Difluorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-51): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.3 (br s, 1H), 10.8 (br s, 1H), 8.63 (d, 1H), 7.95 (d, 2H), 7.85 (m, 2H), 7.58 (t, 1H), 7.41 (t, 1H), 6.59 (s, 1H), 2.27 (s, 3H); MS 338.1 (M+H).

10 Example 281 [2-(3-Chloro-4-fluorophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-52): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.4 (br s, 1H), 10.8 (br s, 1H), 8.65 (d, 1H), 8.50 (d, 1H), 8.36 (m, 1H), 7.85 (m, 1H), 7.60 (m, 1H), 6.62 (s, 1H), 2.30 (s, 3H); MS 354.1 (M+H).

15

Example 282 (5-Methyl-2H-pyrazol-3-yl)-[2-(3-trifluoromethylphenyl)-quinazolin-4-yl]-amine (III-53): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.2 (br, 1H), 10.45 (br, 1H), 7.53 (s, 1H), 7.43 (d, J = 7.2 Hz, 1H), 7.06 (d, J = 8.2 Hz, 1H), 6.65 (d, J = 8.3 Hz, 1H), 6.57 (t, J = 7.6 Hz, 1H), 6.51 (d, J = 7.8 Hz, 1H), 6.43 (t, J = 7.8 Hz, 1H), 6.32 (t, J = 7.6 Hz, 1H), 5.51 (s, 1H), 2.03 (s, 3H); MS 370.2 (M+H).

20 Example 283 [2-(3-Cyanophenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-54): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 9.01 (s, 1H), 8.96 (m, 2H), 8.28 (d, J = 7.3 Hz, 1H), 8.16 (s, br, 2H), 8.06 (t, J = 7.8 Hz, 1H), 7.88 (m, 1H), 6.96 (s, 1H), 2.58 (s, 3H); MS 327.1 (M+H).

30

Example 284 [2-(3-Isopropylphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-55): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.89 (d, J = 7.5 Hz, 1H), 8.37 (s, 1H), 8.26

(s, 1H), 8.08 (m, 2H), 7.81 (t, br, 1H), 7.67 (m, 2H), 6.88 (s, 1H), 3.12 (m, 1H), 2.40 (s, 3H), 1.38 (d, J = 6.9 Hz, 6H); MS 344.2 (M+H).

- 5 Example 285 (5-Methyl-2H-pyrazol-3-yl)-(2-pyridin-3-yl-quinazolin-4-yl)-amine (III-56): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 9.50 (s, 1H), 8.84 (d, J = 7.3 Hz, 1H), 8.80 (d, J = 4.4 Hz, 1H), 8.66 (d, J = 8.2 Hz, 1H), 7.87 (m, 2H), 7.77 (m, 1H), 7.60 (t, J = 7.2 Hz, 1H), 6.67 (s, 1H), 2.28 (s, 10 3H); MS 303.1 (M+H).

- Example 286 [2-(3-Acetylphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-57): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.80 (s, 1H), 8.55 (d, J = 7.7 Hz, 1H), 8.42 15 (d, J = 7.6 Hz, 1H), 8.00 (d, J = 7.0 Hz, 1H), 7.76 (m, 2H), 7.58 (t, J = 7.7 Hz, 1H), 7.48 (s, br, 1H), 6.60 (s, 1H), 2.49 (s, 3H), 2.03 (s, 3H); MS 344.1 (M+H).

- Example 287 [2-(3,5-Ditrifluoromethylphenyl)-quinazolin-20 4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-58): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 10.7 (s, br, 1H), 8.95 (s, 2H), 8.63 (d, J = 8.2 Hz, 1H), 8.25 (s, 1H), 7.86 (m, 2H), 7.58 (t, J = 6.9 Hz, 1H), 6.62 (s, 1H), 2.26 (s, 3H); MS 438.1 (M+H).

- 25 Example 288 [2-(3-Hydroxymethylphenyl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-59): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.74 (d, J = 7.9 Hz, 1H), 8.33 (s, 1H), 8.17 (s, br, 1H), 7.95 (s, br, 1H), 7.89 (s, br, 1H), 30 7.62 (m, 3H), 6.72 (s, 1H), 5.53 (s, 1H), 4.60 (s, 2H), 2.28 (s, 3H); MS 332.1 (M+H).

Example 289 (5-Methyl-2H-pyrazol-3-yl)-[2-(3-phenoxyphenyl)-quinazolin-4-yl]-amine (III-60): mp 231-

232°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.21 (3H, s), 6.59 (1H, s),  
7.10-7.22 (4H, m), 7.41-7.45 (2H, m), 7.54-7.59 (2H, m),  
7.81 (2H, s), 8.09 (1H, s), 8.27 (1H, m), 8.64 (1H, m),  
10.40 (1H, s), 12.20 (1H, s); IR (solid); IR (solid)  
5 1589, 1560, 1541, 1536, 1484, 1360, 1227; MS 394.7 (M+H)<sup>+</sup>.

Example 290 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(3-  
phenoxyphenyl)-quinazolin-4-yl]-amine (III-61): mp 193-  
195°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.67 (2H, m), 0.93 (2H, m), 1.87  
10 (1H, m), 6.56 (1H, s), 7.06-7.20 (4H, m), 7.40-7.43 (2H,  
m), 7.55-7.59 (2H, m), 7.81 (2H, s), 8.11 (1H, s), 8.27  
(1H, m), 8.63 (1H, m), 10.43 (1H, s), 12.26 (1H, s); IR  
(solid); IR (solid) 1589, 1574, 1527, 1483, 1369, 1226;  
MS 420.7 (M+H)<sup>+</sup>.

15

Example 291 (5-Methyl-2H-pyrazol-3-yl)-(2-thiophen-3-yl-  
quinazolin-4-yl)-amine (III-62): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  
δ 11.78 (s, br, 1H), 8.75 (d, J = 8.1 Hz, 1H), 8.68 (s,  
1H), 7.98 (dd, J = 7.9, 7.5 Hz, 1H), 7.89 (m, 2H), 7.81  
20 (m, 1H), 7.68 (t, J = 7.5 Hz, 1H), 6.69 (s, 1H), 2.30 (s,  
3H); MS 308.1 (M+H).

Example 292 (2-Phenyl-quinazolin-4-yl)-(2H-pyrazol-3-yl)-  
amine (III-63): mp 247-249°C; <sup>1</sup>H NMR (DMSO) δ 6.99 (1H, br  
25 s), 7.49-7.58 (5H, m), 7.81 (1H, br s), 7.83 (2H, m),  
8.47-8.49 (2H, m), 8.66 (1H, d), 10.54 (1H, s), 12.59  
(1H, s); IR (solid) 3145, 2922, 1622, 1597; MS 288.2  
(M+H)<sup>+</sup>.

30 Example 293 (2H-Pyrazol-3-yl)-(2-pyridin-4-yl-quinazolin-  
4-yl)-amine (III-64): mp 285-286°C; <sup>1</sup>H NMR (DMSO) δ 6.99  
(1H, br s), 7.65 (1H, m), 7.81-7.94 (3H, m), 8.3-8.35  
(2H, m), 8.73 (1H, d), 8.84-8.90 (2H, m), 10.76 (1H, s),

12.6 (1H, s); IR (solid) 3180, 2972, 1600, 1574; MS 289.2 (M+H)<sup>+</sup>.

Example 294 5-Ethyl-2H-pyrazol-3-yl)-(2-phenyl-

5 quinazolin-4-yl)-amine (III-65): mp 221-222°C; <sup>1</sup>H NMR (DMSO) δ 1.31 (3H, t), 2.68 (2H, d), 6.80 (1H, s), 7.50-7.60 (4H, m), 8.45-8.55 (2H, m), 8.65-8.75 (1H, m), 10.44 (1H, s), 12.27 (1H, s); IR (solid) 3190, 1622, 1595, 1575, 1533, 1482, 1441, 1420, 1403, 1361, 758, 711; MS 316.2 (M+H)<sup>+</sup>.

Example 295 (2-Phenyl-quinazolin-4-yl)-(5-propyl-2H-

pyrazol-3-yl)-amine (III-66): mp 204-205°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.02 (3H, t), 1.66-1.75 (2H, m), 2.69 (2H, t), 6.80 (1H, s), 7.45-7.60 (4H, m), 7.80-7.88 (2H, m), 8.45-8.50 (2H, m), 8.65 (1H, d), 10.39 (1H, s), 12.25 (1H, s); IR (solid) 1621, 1560, 1572, 1533, 1479, 1441, 1421, 1363, 1328, 999, 827, 808, 763, 709, 697; MS 330.2 (M+H)<sup>+</sup>.

Example 296 (5-Isopropyl-2H-pyrazol-3-yl)-(2-phenyl-

quinazolin-4-yl)-amine (III-67): mp 218-219°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.36 (6H, d), 3.05 (1H, m), 6.86 (1H, s), 7.48-7.59 (4H, m), 7.80-7.88 (2H, m), 8.49-8.58 (2H, m), 8.66 (1H, d), 10.47 (1H, s), 12.30 (1H, s); IR (solid) 3173, 2968, 1619, 1593, 1573, 1533, 1478, 1438, 1413, 1398, 1363, 1329, 995, 822, 798, 761, 707, 666, 659; MS 330.2 (M+H)<sup>+</sup>.

Example 297 (5-tert-Butyl-2H-pyrazol-3-yl)-(2-phenyl-

30 quinazolin-4-yl)-amine (III-68): mp 136-137°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.38 (9H, s), 6.87 (1H, br s), 7.51-7.57 (4H, m), 7.84-7.85 (2H, m), 8.49-8.51 (2H, m), 8.65 (1H, d), 10.43 (1H, s), 12.21 (1H, br s); IR (solid) 3162, 2963, 1621, 1590, 1572; MS 344.2 (M+H)<sup>+</sup>.



Example 298 (5-*tert*-Butyl-2*H*-pyrazol-3-yl)-(2-pyridin-4-yl-quinazolin-4-yl)-amine (III-69): mp >300°C; <sup>1</sup>H NMR (DMSO) δ 1.38 (9H, s), 6.82 (1H, br s), 7.63 (1H, m), 7.86-7.91 (2H, m), 8.32-8.33 (2H, d), 8.69 (1H, d), 8.75-8.76 (2H, d), 10.60 (1H, s), 12.31 (1H, br s); IR (solid) 3683, 3149, 2963, 1621; MS 345.2 (M+H)<sup>+</sup>.

Example 299 (5-Cyclopentyl-2*H*-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-70): mp 240-241°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.68-1.89 (6H, m), 2.03-2.17 (2H, m), 3.14-3.22 (1H, m), 6.80 (1H, s), 7.50-7.60 (4H, m), 7.80-7.89 (2H, m), 8.45-8.52 (2H, m), 8.67 (1H, d), 10.52 (1H, s), 12.26 (1H, s); IR (solid) 2957, 1621, 1591, 1571, 1531, 1476, 1438, 1405, 1370, 1325, 999, 951, 801, 775, 761, 747, 710695, 668, 654; MS 356.2 (M+H)<sup>+</sup>.

Example 300 (5-Phenyl-2*H*-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-71): mp 207-209°C; <sup>1</sup>H NMR (DMSO) δ 7.38-7.40 (1H, m), 7.50-7.58 (6H, m), 7.82-7.88 (4H, m), 8.51 (2H, m), 8.67 (1H, s), 10.58 (1H, s), 13.11 (1H, br s); IR (solid) 3345, 3108, 1627, 1612; MS 364.2 (M+H)<sup>+</sup>.

Example 301 (5-Carboxy-2*H*-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-72): (5-Methoxycarbonyl-2*H*-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-73) (345mg, 1 mmole in THF, 6 mL) was treated with NaOH (1M, 4.0 mL), stirred at 50°C for 5 hours, cooled to room temperature, and neutralised with 1M HCl. The mixture was concentrated in vacuo to remove THF then diluted with water and the resulting precipitate filtered. The residual solid was dried at 80°C under vacuum to afford III-72 as an off-white solid (312 mg, 94%): mp 289-291°C

(dec.);  $^1\text{H}$  NMR (DMSO)  $\delta$  7.45 (1H, br s), 7.50-7.60 (5H, m), 7.80-7.88 (2H, m), 7.40-7.50 (2H, m), 8.60-8.70 (1H, d), 10.70 (1H, s), 13.00-13.80 (2H, br s); IR (solid) 1699, 1624, 1607, 1570, 1539, 1506, 1486, 1398, 1333, 1256, 1177, 1004, 827, 764, 705; MS 332.3 (M+H) $^+$ .

Example 302 (5-Methoxycarbonyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-73): mp 271-273°C;  $^1\text{H}$  NMR (DMSO)  $\delta$  3.95 (3H, s), 7.50-7.65 (5H, m), 7.80-7.98 (2H, m), 8.40-8.50 (2H, m), 8.65-8.73 (1H, m), 10.80 (1H, s), 13.80 (1H, s); IR (solid) 3359, 1720, 1624, 1597, 1561, 1538, 1500, 1475, 1435, 1410, 1358, 1329, 1283, 1261, 1146, 1125, 1018, 1010, 944, 827, 806, 780, 763, 703, 690, 670; MS 346.3 (M+H) $^+$ .

15

Example 303 (5-Hydroxymethyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-74): A solution of (5-Methoxycarbonyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-73) (345mg, 1mmol) in anhydrous THF (10mL) was treated with lithium borohydride (125mg, 5.75 mmol) at 65°C for 5 hours. The mixture was cooled to room temperature then combined with 2M HCl and ethyl acetate. Solid sodium hydrogen carbonate was added to achieve pH 8 and the resulting mixture extracted with ethyl acetate. The extracts were dried over magnesium sulphate and concentrated. Purification by flash chromatography (SiO<sub>2</sub>, methanol-dichloromethane gradient) afforded III-74 (95 mg, 30%) as an off-white solid: mp 238-239°C;  $^1\text{H}$  NMR (DMSO)  $\delta$  4.58 (2H, d, CH<sub>2</sub>), 5.35 (1H, s, OH), 6.94 (1H, s), 7.50-7.60 (4H, m), 7.85-7.90 (2H, m), 8.48-8.54 (2H, m), 8.69 (1H, 1H), 10.40 (1H, s), 12.48 (1H, s); IR (solid) 1652, 1621, 1603, 1575, 1558, 1539, 1532, 1480, 1373, 1320, 1276, 1175, 1057, 1037, 1007, 951, 865, 843, 793, 780, 7124; MS 318.2 (M+H) $^+$ .

30

Example 304 (5-Methoxymethyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-75): mp 190-191°C; <sup>1</sup>H NMR (DMSO) δ 3.34 (3H, s), 4.45 (2H, s), 7.00 (1H, s), 7.50-7.62 (4H, m), 7.82-7.90 (2H, m), 8.45-8.52 (2H, m), 8.65 (1H, br s), 10.50 (1H, s), 12.30 (1H, s); IR (solid) 3177, 1606, 1589, 1530, 1479, 1441, 1406, 1374, 1363, 1329, 1152, 1099, 999, 954, 834, 813, 766, 707, 691; MS 332.3 (M+H)<sup>+</sup>.

10

Example 305 [5-(3-Hydroxyprop-1-yl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-76): A solution of (5-benzyloxypropyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-78) (200mg, 0.46mmol) in toluene (4mL) and acetonitrile (8mL) was stirred with trimethylsilyl iodide (0.64ml, 4.6mmol) at 55°C for 3 hours to afford an amber coloured solution. This mixture was diluted with ethyl acetate and aqueous sodium hydrogen carbonate. The resulting layers were separated, the organic layer was dried over magnesium sulphate and concentrated in vacuo. Purification by flash chromatography (SiO<sub>2</sub>, methanol-dichloromethane gradient) affords a yellow oil (115mg). Trituration with dichloromethane affords III-76 as an off-white solid dried at 75°C under vacuum (83mg, 52%): mp 164-165°C; <sup>1</sup>H NMR (DMSO) δ 1.80-1.90 (2H, m), 2.70-2.80 (2H, m), 3.50-3.60 (2H, m), 4.59 (1H, s), 6.80 (1H, s), 7.50-7.60 (4H, m), 7.82-7.90 (2H, m), 8.48-8.53 (2H, m), 8.63 (1H, s), 10.40 (1H, s), 12.25 (1H, s); IR (solid) 1622, 1587, 1574, 1562, 1528, 1480, 1440, 1421, 1368, 1329, 1173, 1052, 1030, 1006, 952, 833, 762, 734, 706, 690, 671, 665; MS 346.0 (M+H)<sup>+</sup>.

15

20

25

30

Example 306 [5-(3-Methoxyprop-1-yl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-77): mp 169-170°C; <sup>1</sup>H

NMR (DMSO-d<sub>6</sub>)  $\delta$  1.86-1.97 (2H, m), 2.75 (2H, t), 3.30 (3H, s), 3.45 (2H, t), 6.80 (1H, s), 7.50-7.60 (4H, m), 7.80-7.90 (2H, m), 8.45-8.55 (2H, m), 8.67 (1H, d), 10.30 (1H, s), 12.25 (1H, s); IR (solid) 1620, 1591, 1572, 1532, 1476, 1425, 1408, 1373, 1326, 1117, 1003, 831, 764, 714, 695; MS 360.3 (M+H)<sup>+</sup>.

Example 307 [5-(3-Benzylloxyprop-1-yl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-78): mp 177-178°C;  
<sup>1</sup>H NMR (DMSO)  $\delta$  1.92-2.03 (2H, m), 3.76-3.85 (2H, m), 3.52-3.62 (2H, m), 4.51 (2H, s), 6.82 (1H, s), 7.28-7.40 (5H, m), 7.46-7.58 (4H, m), 7.80-7.85 (2H, m), 8.47-8.52 (2H, m), 8.66 (1H, d), 10.45 (1H, s); IR (solid) 1621, 1591, 1562, 1532, 1479, 1454, 1426, 1408, 1374, 1101, 1006, 835, 766, 738, 712, 696; MS 436.3 (M+H)<sup>+</sup>.

Example 308 [5-(3-Aminoprop-1-yl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-79): A solution of [5-(3-tert-butoxycarbonylaminoprop-1-yl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-80) (250mg, 0.56mmol), in dichloromethane (3mL) at 0°C was treated with TFA (2mL). The mixture was warmed to room temperature then concentrated *in vacuo*. The residue was triturated and concentrated from dichloromethane (3x5mL) and ether, then triturated with dichloromethane to crystallize the TFA salt. The resulting solid was collected by filtration and dissolved in a mixture of ethanol (3mL) and water (3mL). Potassium carbonate was added in portions to achieve pH 8 then the mixture allowed to crystallize. The product was collected by filtration and dried at 80°C under vacuum to afford III-79 as an off-white powder (122mg, 63%): mp 205-207°C; <sup>1</sup>H NMR (DMSO)  $\delta$  1.68-1.83 (2H, m), 2.65-2.80 (4H, m), 6.80 (1H, s), 7.50-7.60 (4H, m), 7.80-7.90 (2H, m), 8.45-8.53 (2H,

m), 8.65 (1H, d), 10.45 (1H, br s); IR (solid) 1621, 1598, 1568, 1533, 1484, 1414, 1364, 1327, 1169, 1030, 951, 830, 776, 764, 705, 677; MS 345.3 (M+H)<sup>+</sup>.

5 Example 309 [5-(3-tert-Butoxycarbonylaminoprop-1-yl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-80): mp 199-200°C; <sup>1</sup>H NMR (DMSO) δ 1.37 (9H, s), 1.71-1.82 (2H, m), 2.67 (2H, t), 3.00-3.11 (2H, m), 7.81 (1H, s), 7.99 (1H, s), 7.50-7.60 (4H, m), 7.80-7.85 (2H, m), 8.48-  
10 8.52 (2H, m), 8.63 (1H, d), 10.40 (1H, s), 12.26 (1H, m); IR (solid) 2953, 1687, 1622, 1594, 1573, 1535, 1481, 1441, 1419, 1364, 1327, 1281, 1252, 1166, 1070, 1028, 998, 951, 848, 807, 768, 740, 728, 710, 693; MS 445.3 (M+H)<sup>+</sup>.

15

Example 310 5-Isopropylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-81): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 1.20 (d, J = 6.6 Hz, 6H), 4.13 (m, 1H), 7.42 (br. s, 1H), 7.61 (dd, J = 7.0, 7.7 Hz, 2H), 7.66 (t, J =  
20 7.1 Hz, 1H), 7.71 (m, 1H), 7.99 (m, 2H), 8.39 (m, 1H), 8.42 (d, J = 7.1 Hz, 2H), 8.74 (d, J = 8.2 Hz, 1H), 11.41 (br. s, 1H); EI-MS 373.2 (M+H); HPLC-Method C, R<sub>t</sub> 14.09 min.

25 Example 311 (5-Allylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-82): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 4.02 (m, 2H), 5.15 (m, 1H), 5.23 (m, 1H), 5.94 (m, 1H), 7.45 (br. s, 1H), 7.60 (t, J = 6.9 Hz, 2H), 7.64 (m, 1H), 7.72 (m, 1H), 7.98 (m, 2H), 8.43 (m 2H), 8.72 (d, J = 8.2  
30 Hz, 1H), 8.84 (br. s, 1H), 11.34 (br. s, 1H); EI-MS 371.2 (M+H); HPLC-Method C, R<sub>t</sub> 13.67 min.

Example 312 [5-(2-Methoxyethylcarbamoyl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-83): <sup>1</sup>H NMR

(500MHz, DMSO-d<sub>6</sub>)  $\delta$  3.32 (s, 3H), 3.48 (m, 4H), 7.36 (br. s, 1H), 7.62 (m, 2H), 7.63 (m, 1H), 7.71 (m, 1H), 7.98 (m, 2H), 8.41 (dd, J = 1.4, 7.0, 2H), 8.70 (m, 2H), 11.30 (br. s, 1H); EI-MS 389.2 (M+H); HPLC-Method C, R<sub>t</sub> 12.37 min.

Example 313 (5-Benzylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-84): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>)  $\delta$  4.52 (d, J = 6.0 Hz, 2H), 7.29 (m, 1H), 7.38 (d, J = 4.2 Hz, 4H), 7.58 (t, J = 7.5 Hz, 2H), 7.63 (m, 1H), 7.72 (m, 1H), 7.98 (m, 2H), 8.43 (d, J = 7.7 Hz, 2H), 8.72 (d, J = 7.5 Hz, 1H), 9.23 (br. s, 2H), 11.34 (br. s, 1H); EI-MS 421.2 (M+H); HPLC-Method C, R<sub>t</sub> 16.76 min.

Example 314 (5-Cyclohexylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-85): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>)  $\delta$  1.16 (m, 1H), 1.34 (m, 4H), 1.62 (d, J = 2.6 Hz, 1H), 1.76 (m, 2H), 1.85 (m, 2H), 3.79 (m, 1H), 7.43 (m, 1H), 7.60 (t, J = 7.2 Hz, 2H), 7.65 (t, J = 7.1 Hz, 1H), 7.71 (ddd, J = 2.2, 5.4, 8.2 Hz, 1H), 7.98 (m, 2H), 8.35 (m, 1H), 8.43 (dd, J = 1.4, 7.2 Hz, 2H), 8.72 (d, J = 8.2 Hz, 1H), 11.34 (br. s, 1H); EI-MS 413.5 (M+H); HPLC-Method C, R<sub>t</sub> 17.18 min.

Example 315 (5-Diethylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-86): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>)  $\delta$  1.18 (br. s, 3H), 1.25 (br. s, 3H), 3.49 (br. s, 2H), 3.69 (b. s, 2H), 7.21 (s, 1H), 7.59 (t, J = 6.9 Hz, 2H), 7.62 (m, 1H), 7.70 (m, 1H), 7.96 (m, 2H), 8.39 (d, J = 7.1 Hz, 2H), 8.74 (d, J = 8.4 Hz, 1H), 11.37 (br. s, 1H); EI-MS 387.2 (M+H); HPLC-Method C, R<sub>t</sub> 14.50 min.

Example 316 [5-(Benzyl-methyl-carbamoyl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-87): <sup>1</sup>H NMR

(500MHz, DMSO-d<sub>6</sub>) δ: 3.33 (s, 3H), 4.75 (s, 2H), 7.26 (m, 1H), 7.31 (m, 1H), 7.38 (m, 4H), 7.58 (m, 2H), 7.70 (m, 1H), 7.95 (m, 3H), 8.26 (m, 1H), 8.40 (d, J = 7.8 Hz, 2H), 8.75 (m, 1H), 11.2 (br. s, 1H); EI-MS 435.2 (M+H); HPLC-Method C, R<sub>t</sub> 16.77 min.

Example 317 (2-Phenyl-quinazolin-4-yl)-(5-propylcarbamoyl-2H-pyrazol-3-yl)-amine (III-88): <sup>1</sup>H NMR

(500MHz, DMSO-d<sub>6</sub>) δ 0.94 (t, J = 7.3 Hz, 3H), 1.57 (m, 2H), 3.24 (q, J = 6.5 Hz, 2H), 7.39 (br. s, 1H), 7.60 (t, J = 7.3 Hz, 2H), 7.64 (m, 1H), 7.71 (br. t, J = 6.5 Hz, 1H), 7.98 (m, 2H), 8.42 (d, J = 7.2 Hz, 2H), 8.61 (br. s, 1H), 8.72 (d, J = 8.5 Hz, 1H), 11.34 (br. s, 1H); EI-MS 373.3 (M+H); HPLC-Method C, R<sub>t</sub> 13.51 min.

Example 318 [5-(Ethyl-isopropyl-carbamoyl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-89): <sup>1</sup>H NMR

(500MHz, DMSO-d<sub>6</sub>) δ 0.92 (t, J = 7.4 Hz, 6H), 1.52 (m, 2H), 1.59 (m, 1H), 3.79 (m, 2H), 7.53 (br. s, 1H), 7.57 (t, J = 7.5 Hz, 2H), 7.65 (t, J = 7.2 Hz, 1H), 7.71 (m, 1H), 7.99 (m, 2H), 8.23 (br. d, J = 8.8 Hz, 1H), 8.46 (d, J = 7.5 Hz, 2H), 8.74 (d, J = 8.4 Hz, 1H), 11.34 (br. s, 1H); EI-MS 401.2 (M+H); HPLC-Method C, R<sub>t</sub> 15.51 min.

Example 319 (5-Cyclopropylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-90): <sup>1</sup>H NMR (500MHz,

DMSO-d<sub>6</sub>) δ 0.60 (m, 2H), 0.74 (m, 2H), 2.86 (m, 1H), 7.34 (br. s, 1H), 7.62 (m, 3H), 7.70 (m, 1H), 7.97 (m, 2H), 8.41 (d, J = 7.9 Hz, 2H), 8.63 (br. s, 1H), 8.72 (d, J = 7.8 Hz, 1H), 11.35 (br. s, 1H); EI-MS 371.2 (M+H); HPLC-Method C, R<sub>t</sub> 12.64 min.

Example 320 (5-Isobutylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-91): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 0.94 (d, J = 6.7 Hz, 6H), 1.88 (m, 1H), 3.12 (t, J = 6.4 Hz, 2H), 7.45 (br. s, 1H), 7.58 (t, J = 7.2 Hz, 3H), 7.64 (t, J = 7.1 Hz, 1H), 7.71 (m, 1H), 7.98 (m, 2H), 8.44 (dd, J = 1.3, 7.9 Hz, 2H), 8.62 (br. s, 1H), 8.72 (d, J = 8.3 Hz, 1H), 11.33 (br. s, 1H); EI-MS 387.2 (M+H); HPLC-Method C, R<sub>t</sub> 14.70 min.

10 Example 321 {5-[(3S)-3-Methoxymethyl-pyrrolidine-1-carbonyl]-2H-pyrazol-3-yl}-(2-phenyl-quinazolin-4-yl)-amine (III-93): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 2.00 (m, 2H), 2.12 (m, 1H), 3.29 (s, 3H), 3.45 (t, J = 8.7 Hz, 1H), 3.57 (dd, J = 3.2, 9.3 Hz, 1H), 3.86 (m, 1H), 3.92 (m, 1H), 4.36 (m, 2H), 7.45 (br. s, 1H), 7.59 (t, J = 7.2 Hz, 2H), 7.63 (m, 1H), 7.69 (m, 1H), 7.97 (m, 2H), 8.40 (d, J = 7.5 Hz, 2H), 8.74 (d, J = 7.6 Hz, 1H), 11.38 (br. s, 1H); EI-MS 429.2 (M+H); HPLC-Method C, R<sub>t</sub> 13.84 min.

20 Example 322 (2-Phenyl-quinazolin-4-yl)-(5-m-tolylcarbamoyl-2H-pyrazol-3-yl)-amine (III-94): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 2.33 (s, 3H), 6.97 (d, J = 7.5 Hz, 1H), 7.27 (t, J = 7.8 Hz, 1H), 7.62 (m, 7H), 7.72 (m, 1H), 7.98 (m, 2H), 8.46 (dd, J = 2.0, 7.9 Hz, 2H), 8.71 (m, 1H), 10.29 (s, 1H), 11.31 (br. s, 1H); EI-MS 421.2 (M+H); HPLC-Method C, R<sub>t</sub> 17.11 min.

Example 323 (2-Phenyl-quinazolin-4-yl)-(5-p-tolylcarbamoyl-2H-pyrazol-3-yl)-amine (III-95): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 2.30 (s, 3H), 7.20 (d, J = 8.3 Hz, 2H), 7.62 (m, 5H), 7.68 (d, J = 8.3 Hz, 2H), 7.72 (m, 1H), 7.98 (m, 2H), 8.46 (dd, J = 1.8, 7.0 Hz, 2H), 8.72 (m, 1H), 10.31 (s, 1H), 11.36 (br. s, 1H); EI-MS 421.2 (M+H); HPLC-Method C, R<sub>t</sub> 16.95 min.



Example 324 (5-Methylcarbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-96):  $^1\text{H}$  NMR (500MHz, DMSO- $d_6$ )  $\delta$  2.82 (d,  $J$  = 4.6 Hz, 3H), 7.31 (br. s, 1H),  
5 7.62 (m, 3H), 7.69 (m, 1H), 7.97 (m, 2H), 8.42 (d,  $J$  = 7.1 Hz, 2H), 8.59 (br. s, 1H), 8.71 (d,  $J$  = 8.0 Hz, 1H), 11.30 (br. s, 1H); EI-MS 345.1 (M+H); HPLC-Method C,  $R_t$  11.02 min.

10 Example 325 [5-(Morpholine-4-carbonyl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-97):  $^1\text{H}$  NMR (500MHz, DMSO- $d_6$ )  $\delta$  3.33 (m, 4H), 3.83 (m 4H), 7.34 (br. s, 1H),  
7.53 (m, 4H), 7.86 (m, 2H), 8.43 (m, 2H), 8.67 (d,  $J$  = 8.6 Hz, 1H), 10.70 (s, 1H), 13.56 (s, 1H); EI-MS 401.2  
15 (M+H); HPLC-Method A,  $R_t$  2.68 min.

Example 326 [5-(1-Methylpiperazine-4-carbonyl)-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-98):  
 $^1\text{H}$  NMR (500MHz, DMSO- $d_6$ )  $\delta$  2.25 (s, 3H), 2.43 (m, 4H),  
20 3.87 (m 4H), 7.33 (br. s, 1H), 7.53 (m, 4H), 7.87 (m, 2H), 8.45 (m, 2H), 8.67 (d,  $J$  = 7.6 Hz, 1H), 10.70 (s, 1H), 13.30 (s, 1H); EI-MS 414.2 (M+H); HPLC-Method A,  $R_t$  2.38 min.

25 Example 327 [5-(2-Hydroxyethylcarbamoyl-2H-pyrazol-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-99):  $^1\text{H}$  NMR (500MHz, DMSO- $d_6$ )  $\delta$  3.36 (m, 2H), 3.52 (m, 2H), 4.79 (m, 1H), 7.50 (m, 5H), 7.83 (m, 2H), 8.50 (m, 4H), 10.52 (br. s, 1H), 13.25 (s, 1H); EI-MS 375.1 (M+H); HPLC-Method A,  $R_t$  2.51  
30 min.

Example 328 (5-Carbamoyl-2H-pyrazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-100): To a solution of 5-(2-phenyl-quinazolin-4-ylamino)-1H-pyrazole-3-carboxylic

acid 2,5-dioxo-pyrrolidin-1-yl ester (270 mg, 0.63 mmol) in DMF (20 ml) was added a solution of ammonia in 1,4-dioxane (0.5 M, 10 ml). The resulting mixture was stirred at room temperature for 24 h. After

5 concentration of the solvents, the residue was added to water (20 ml). The resulting precipitate was collected to afford III-100 (168 mg, 80%) as a yellow solid. <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 7.77-7.51 (m, 6H), 7.86 (br s, 2H), 8.11 (m, 1H), 8.50 (m, 2H), 8.63 (m, 1H), 10.52 (s, 1H),  
10 11.25 (s, 1H); EI-MS 331.1 (M+H); HPLC-Method A, R<sub>t</sub> 2.52 min.

Example 329 (4-Bromo-2H-pyrazol-3-yl)-(2-phenyl-

quinazolin-4-yl)-amine (III-101): Prepared according to

15 Method A to afford a yellow solid, mp 189°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.44-7.46 (3H, m), 7.58 (1H, m), 7.87 (2H, d), 8.15 (1H, s), 8.31-8.34 (2H, m), 8.49 (1H, d), 10.08 (1H, s), 13.13 (1H, s); IR (solid) 3286, 2969, 1738, 1632; MS 366.2/368.2 (M+H)<sup>+</sup>.

20

Example 330 (4-Bromo-5-methyl-2H-pyrazol-3-yl)-(2-phenyl-

quinazolin-4-yl)-amine (III-102): mp 183-185°C; <sup>1</sup>H NMR

(DMSO) δ 2.33 (3H, br s), 7.44-7.46 (3H, m), 7.57 (1H, m), 7.84-7.87 (2H, m), 8.31-8.34 (2H, m), 8.48 (1H, d),  
25 10.05 (1H, s), 12.91 (1H, br s); IR (solid) 3362, 3065, 2831, 1619, 1578; MS 380.2/382.2 (M+H)<sup>+</sup>.

Example 331 (4-Cyano-2H-pyrazol-3-yl)-(2-phenyl-

quinazolin-4-yl)-amine (III-103): mp >250°C; <sup>1</sup>H NMR (DMSO)

30 δ 7.47-7.49 (3H, m), 7.64 (1H, m), 7.91 (2H, m), 8.40-8.43 (2H, m), 8.53 (1H, d), 8.71 (1H, d), 10.61 (1H, s), 13.60 (1H, s); IR (solid) 3277, 3069, 2855, 2231, 1625; MS 313.2 (M+H)<sup>+</sup>.

Example 332 (5-Methyl-2H-pyrazol-3-yl)-(2-morpholin-4-yl-quinazolin-4-yl)-amine (III-104): mp 223-224°C; <sup>1</sup>H NMR (DMSO) δ 2.26(3H, s), 3.65(4H, m), 3.75(4H, m), 6.44(1H, s), 7.12(1H, d), 7.33(1H, d), 7.56(1H, t), 8.37(1H, d), 10.01(1H, s), 12.13(1H, br s); IR (solid) 1621, 1578, 1537, 1475, 1434, 1385; MS 311.0 (M+H)<sup>+</sup>.

Example 333 (5-Methyl-2H-pyrazol-3-yl)-(2-piperazin-1-yl-quinazolin-4-yl)-amine (III-105): mp 179-181°C; <sup>1</sup>H NMR (DMSO) δ 2.26(3H, s), 2.74 (4H, br s), 3.71(4H, br s), 6.43(1H, s), 7.08(1H, t), 7.30(1H, d), 7.53(1H, t), 8.34(1H, d), 9.50(1H, s), 12.08(1H, br s); IR (solid) 2853, 1619, 1603, 1566, 1549, 1539; MS 310.0 (M+H)<sup>+</sup>

Example 334 [2-(4-Methylpiperidin-1-yl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-106): mp 148-150°C; <sup>1</sup>H NMR (DMSO) δ 1.06(3H, d), 1.03(2H, m), 1.51-1.70(3H, m), 2.26(3H, s), 2.86(2H, m), 4.73(2H, d), 6.44(1H, s), 7.06(1H, d), 7.29(1H, d), 7.52(1H, t), 8.32(1H, d), 9.92(1H, s), 12.09(1H, br s); IR (solid) 2917, 2840, 1629, 1593, 1562, 1546, 1486; MS 323.0 (M+H)<sup>+</sup>.

Example 335 [2-(4-Methylpiperazin-1-yl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-107): mp 105-107°C; <sup>1</sup>H NMR (DMSO) δ 2.21(3H, s), 2.26(3H, s), 2.34(4H, m), 3.75(4H, m), 6.45(1H, s), 7.09(1H, t), 7.31(1H, d), 7.54(1H, t), 8.34(1H, d), 9.96(1H, s), 12.12(1H, br s); IR (solid) 2934, 2844, 2804, 1620, 1593, 1572, 1536, 1476; MS 324.0 (M+H)<sup>+</sup>.

Example 336 (5-Methyl-2H-pyrazol-3-yl)-(2-piperidin-1-yl-quinazolin-4-yl)-amine (III-108): mp 294°C; <sup>1</sup>H NMR (DMSO) δ 1.45-1.58 (4H, m), 1.63 (2H, m), 2.26 (3H, s), 3.79

(4H, m), 6.45 (1H, br s), 7.06 (1H, t), 7.29 (1H, d), 7.52 (1H, t), 8.33 (1H, d), 9.92 (1H, s), 12.11 (1H, br s); IR (solid) 2929, 2847, 1632, 1591, 1500, 1482, 1437, 1382; MS 309.3 (M+H)<sup>+</sup>.

5

Example 337 [2-(4-(2-Hydroxyethylpiperidin-1-yl)-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (III-109): mp 269°C; <sup>1</sup>H NMR (DMSO) δ 1.50 (4H, br s), 1.76 (4H, br s), 2.25 (3H, s), 3.78 (4H, t), 6.55 (1H, br s), 7.03 (1H, t), 7.28 (1H, d), 7.50 (1H, t), 8.33 (1H, d), 9.92 (1H, s), 12.09 (1H, br s); IR (solid) 3427, 2963, 2927, 2909, 2872, 2850, 1623, 1595, 1586, 1568, 1504, 1486, 1468, 1386, 1427; MS 323.3 (M+H)<sup>+</sup>.

Example 338 [2-(4-(2-Hydroxyethylpiperidin-1-yl)-quinazolin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (III-110): mp 175°C; <sup>1</sup>H NMR (DMSO) δ 1.08 (2H, m), 1.38 (2H, m), 1.57-1.83 (3H, m), 2.26 (3H, s), 2.85 (2H, t), 3.47 (2H, m), 4.38 (1H, t), 4.75 (2H, d), 6.45 (1H, br s), 7.06 (1H, t), 7.29 (1H, d), 7.52 (1H, t), 8.32 (1H, d), 9.93 (1H, s), 12.12 (1H, br s); IR (solid) 3365, 3073, 2972, 2868, 1622, 1604, 1586, 1568, 1486, 1463, 1440, 1394; MS 353.2 (M+H)<sup>+</sup>.

Example 339 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(4-methylpiperidin-1-yl)-quinazolin-4-yl]-amine (III-111): To a solution of (5-cyclopropyl-1H-pyrazol-3-yl)-(2-chloro-quinazolin-4-yl)-amine (118 mg, 0.41 mmol) in tert-butanol (3.0 mL) was added 4-methylpiperidine (0.49 mL, 4.1 mmol) and the reaction mixture heated at reflux overnight. The reaction mixture was concentrated in vacuo and the residue dissolved in a mixture EtOH:water (1:3, 4 mL). Potassium carbonate (57mg, 0.41 mmol) was added and the mixture stirred at room temperature for 2

hours. The resulting suspension was filtered, washed with water (x2), and rinsed with Et<sub>2</sub>O (x2) to afford III-111 as a white solid (123mg, 85%): mp 190°C; <sup>1</sup>H NMR (DMSO) δ 0.66 (2H, s), 0.93 (5H, br s), 1.07 (2H, d), 1.66 (3H, s), 1.91 (1H, s), 2.85 (2H, t), 4.72 (2H, d), 6.33 (1H, s), 7.06 (1H, t), 7.29 (1H, d), 7.52 (1H, t), 8.31 (1H, d), 9.95 (1H, s), 12.18 (1H, br s); IR (solid) 2925, 2852, 1622, 1590, 1581, 1558, 1494, 1481, 1453, 1435, 1394; MS 349.2 (M+H)<sup>+</sup>.

10

Example 340 [2-(1,4-Dioxo-8-aza-spiro[4,5]dec-8-yl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-112): mp 191°C; <sup>1</sup>H NMR (DMSO) δ 1.65 (4H, s), 2.26 (3H, s), 3.90 (4H, s), 3.93 (4H, s), 6.43 (1H, br s), 7.09 (1H, t), 7.32 (1H, d), 7.54 (1H, t), 8.35 (1H, d), 9.99 (1H, br s), 12.13 (1H, br s); IR (solid) 3069, 2964, 2927, 2868, 1618, 1581, 1568, 1540, 1495, 1481, 1435, 1390; MS 367.3 (M+H)<sup>+</sup>.

15

Example 341 [2-(4-Cyclopentylamino-piperidin-1-yl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-113): mp 191°C; <sup>1</sup>H NMR (DMSO) δ 1.33 (2H, d), 1.65 (4H, s), 1.87 (2H, d), 2.20 (1H, s), 2.26 (3H, s), 2.49 (2H, s), 3.00 (2H, t), 3.36 (2H, s), 4.61 (2H, d), 6.45 (1H, br s), 7.07 (1H, s), 7.31 (1H, d), 7.52 (1H, s), 8.33 (1H, d), 9.94 (1H, br s), 12.12 (1H, br s); IR (solid) 3371, 2943, 1622, 1600, 1581, 1545, 1509, 1463, 1440, 1390; MS 378.2 (M+H)<sup>+</sup>.

25

Example 342 [2-(4-Hydroxypiperidin-1-yl)-quinazolin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (III-114): mp 123°C; <sup>1</sup>H NMR (DMSO) δ 1.34 (2H, d), 1.80 (2H, d), 2.26 (3H, s), 3.24 (2H, t), 3.72 (1H, br s), 4.39 (2H, d), 4.70 (1H, d), 6.44 (1H, br s), 7.07 (1H, t), 7.30 (1H, d), 7.53

30

(1H, t), 8.33 (1H, d), 9.94 (1H, br s), 12.11 (1H, br s); IR (solid) 3265, 3151, 2927, 2863, 1622, 1600, 1572, 1540, 1504, 1476, 1440, 1390, 1349, 1066, 1098; MS 325.3 (M+H)<sup>+</sup>.

5

Example 343 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(4-hydroxy-4-phenylpiperidin-1-yl)-quinazolin-4-yl]-amine

(III-115): mp 131°C; <sup>1</sup>H NMR (DMSO) δ 0.64 (2H, q), 0.93 (2H, q), 1.68 (2H, d), 1.83-1.97 (3H, m), 3.20-3.45 (2H, m), 4.69 (2H, d), 5.11 (1H, s), 6.37 (1H, br s), 7.08 (1H, t), 7.20 (1H, t), 7.31 (3H, t), 7.49 (2H, d), 7.53 (1H, t), 8.33 (1H, d), 9.98 (1H, br s), 12.18 (1H, br s); IR (solid) 3362, 2952, 2934, 2911, 2870, 2825, 1618, 1584, 1570, 1559, 1536, 1481, 1459, 1431, 1372, 1336, 1213, 994; MS 427.6 (M+H)<sup>+</sup>.

Example 344 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(1,3-dihydro-isoindol-2-yl)-quinazolin-4-yl]-amine (III-116):

Prepared according to Method E-I to afford an off-white solid, mp 237°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.79 (2H, s), 1.00 (2H, d), 1.99 (1H, m), 4.92 (4H, d), 6.72 (1H, br s), 7.13 (1H, t), 7.33 (2H, s), 7.30-7.48 (3H, m), 7.58 (1H, t), 8.40 (1H, d), 10.12 (1H, s), 12.17 (1H, s); IR (solid) 3449, 3318, 2850, 1623, 1595, 1577, 1541, 1509, 1482, 1432, 1391, 1359, 1141, 1027, 877, 814; MS 369.4 (M+H)<sup>+</sup>.

Example 345 (2-Azepan-1-yl)-quinazolin-4-yl]-(5-cyclopropyl-2H-pyrazol-3-yl)-amine (III-117): mp 199-

200°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.60-0.70 (2H, m), 0.90-1.00 (2H, m), 1.45-1.57 (4H, m), 1.70-1.85 (4H, m), 1.88-1.97 (1H, m), 3.75-3.87 (4H, m), 6.42 (1H, s), 7.02 (1H, t), 7.27 (1H, d), 7.49 (1H, t), 8.29 (1H, d), 9.91 (1H, s), 12.19 (1H, br s); IR (solid) 2929, 1624, 1595, 1581,

1563, 1542, 1498, 1482, 1440, 1426, 1397, 1356, 1305,  
1000, 825, 754; MS 349.2 (M+H)<sup>+</sup>.

Example 346 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(3,4-

5 dihydro-1H-isoquinolin-2-yl)-quinazolin-4-yl]-amine (III-  
118): mp 182-184°C; <sup>1</sup>H NMR (DMSO) δ 0.75 (2H, d), 1.02  
(2H, d), 1.96 (1H, m), 2.89 (2H, m), 4.05 (2H, m), 4.94  
(2H, s), 6.46 (1H, s), 7.10 (1H, t), 7.21 (4H, d), 7.37  
(1H, d), 7.55 (1H, d), 8.36 (1H, d), 10.05 (1H, s), 12.23  
10 (1H, br s); IR (solid) 1621, 1581, 1560, 1537, 1479,  
1456, 1426, 1396, 1374, 1341, 1222; MS 383.3 (M+H)<sup>+</sup>.

Example 347 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(2,3-

dihydro-indol-1-yl)-quinazolin-4-yl]-amine (III-119): mp  
15 150-153°C; <sup>1</sup>H NMR (DMSO) δ 0.74 (2H, d), 0.98 (2H, d),  
1.96 (1H, m), 3.15 (2H, t), 4.25 (2H, t), 6.45 (1H, br  
s), 6.88 (1H, t), 7.09 (1H, t), 7.20 (2H, m), 7.53 (1H,  
d), 7.65 (1H, t), 8.43 (2H, br s), 10.09 (1H, s), 12.28  
(1H, br s); IR (solid) 1621, 1588, 1577, 1564, 1537,  
20 1487, 1455, 1425, 1386, 1259; MS 369.3 (M+H)<sup>+</sup>.

Example 348 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(4-

hydroxymethylpiperidin-1-yl)-quinazolin-4-yl]-amine (III-  
120): mp 142°C; <sup>1</sup>H NMR (DMSO) δ 0.67 (2H, d), 0.96 (2H,  
25 d), 1.10 (2H, q), 1.55-1.70 (3H, m), 1.91 (1H, m), 2.85  
(2H, t), 3.28 (2H, s), 4.48 (1H, s), 4.76 (2H, d), 6.34  
(1H, s), 7.06 (1H, t), 7.30 (1H, d), 7.52 (1H, t), 8.31  
(1H, d), 9.96 (1H, s), 12.19 (1H, s); IR (solid) 3363,  
3000, 2927, 2854, 1618, 1604, 1573, 1536, 1509, 1477,  
30 1436, 1395, 1354, 1314, 1241, 1186, 1091, 995, 941, 823;  
MS 365.8 (M+H)<sup>+</sup>.

Example 349 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(3,4-

dihydro-2H-quinolin-1-yl)-quinazolin-4-yl]-amine (III-

121): mp 137-145°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.55 (2H, d), 0.88 (2H, d), 1.78 (1H, m), 1.92 (2H, t), 2.75 (2H, t), 4.04 (2H, t), 6.20 (1H, br s), 6.97 (1H, t), 7.14 (1H, m), 7.19 (1H, t), 7.42 (1H, d), 7.61 (1H, t), 7.67 (1H, d), 8.43 (1H, d), 10.04 (1H, s), 12.21 (1H, br s); IR (solid) 1622, 1572, 1539, 1493, 1454, 1420, 1373, 1249; MS 383.3 (M+H)<sup>+</sup>.

Example 350 (5-Methoxycarbonyl-2H-pyrazol-3-yl)-[2-(piperidine-1-yl)-quinazolin-4-yl]-amine (III-122): <sup>1</sup>H NMR (500MHz, CDCl<sub>3</sub>) δ 1.7-1.8 (6H, m), δ 3.8 (4H, m), δ 3.9 (3H, s), δ 5.5 (1H, s), δ 7.15 (1H, t), δ 7.4 (1H, d), δ 7.6 (1H, t), δ 8.0 (1H, d). HPLC-Method B, (starting with 95% H<sub>2</sub>O) R<sub>t</sub> 7.4 min; MS (ES+) 353.24 (M+H).

15

Example 351 [5-(Piperidine-1-carbonyl)-2H-pyrazol-3-yl]-[2-(piperidine-1-yl)-quinazolin-4-yl]-amine (III-123): HPLC-Method B, (starting with 95% H<sub>2</sub>O:0.1% TFA) R<sub>t</sub> 8.0 min; MS (ES+) 406.30, (ES-) 404.30.

20

Example 352 (5-Hydroxymethyl-2H-pyrazol-3-yl)-[2-(piperidin-1-yl)-quinazolin-4-yl]-amine (III-124): To a solution of III-122 (10.0 mg, 0.028 mmol) in THF (6 mL) at ambient temperature was slowly added a 1M solution of LiAlH<sub>4</sub> in THF (0.05 mL, 0.05 mmol). After 15 minutes the solution was quenched with water and 1N HCl. The product was extracted from the aqueous layer with EtOAc. The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The residue was purified by preparatory HPLC to afford III-124 (4.0 mg, 44%). HPLC-Method B, (starting with 95% H<sub>2</sub>O:0.1% TFA) R<sub>t</sub> 6.1 min; MS (ES+) 325.13 (M+H), (ES-) 323.13 (M-H).



Example 353 (5-Carbamoyl-2H-pyrazol-3-yl)-[2-(piperidin-1-yl)-quinazolin-4-yl]-amine (III-125): A solution of

III-122 (1.5 g, 4.3 mmol) in 2.0 M NH<sub>3</sub>/MeOH (100 mL) was heated at 110°C for 2 days. The dark brown reaction

mixture was concentrated in vacuo to afford a viscous oil which was purified by column chromatography to yield 0.7 g (50%) of III-125. <sup>1</sup>H NMR (500MHz, CD<sub>3</sub>OD-d<sub>3</sub>) δ 1.6 (4H, m), δ 1.7 (2H, m), δ 3.3 (1H, s), δ 3.8 (4H, m), δ 5.5 (1H, s), δ 7.15 (1H, t), δ 7.45 (1H, d), δ 7.55 (1H, t), δ 8.0 (1H, d); HPLC-Method B, (starting with 95% H<sub>2</sub>O:0.1% TFA) R<sub>t</sub> 5.9 min; MS (ES+) 338.13, (ES-) 336.15.

Example 354 (5-Carbamoyl-2H-pyrazol-3-yl)-[2-(4-methylpiperidin-1-yl)-quinazolin-4-yl]-amine (III-126):

HPLC-Method B, (starting with 95% H<sub>2</sub>O:0.1% TFA) R<sub>t</sub> 6.4 min; MS (ES+) 352.19, (ES-) 350.20.

Example 355 (5,7-Difluoro-1H-indazol-3-yl)-(2-phenyl-5,6,7,8-tetrahydroquinazolin-4-yl)-amine (III-127): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.7 (s, 1H), 10.3 (s, br, 1H), 7.90 (d, 2H), 7.52 (t, 1H), 7.45 (m, 3H), 7.26 (d, 1H), 2.99 (m, 2H), 2.75 (m, 2H), 1.95 (br, 4H) ppm; MS (ES+) 378.24 (M+H); (ES-) 376.23 (M-H); HPLC-Method A, R<sub>t</sub> 3.04 min.

Example 356 (2-Phenyl-5,6,7,8-tetrahydroquinazolin-4-yl)-(5-trifluoromethyl-1H-indazol-3-yl)-amine (III-128): <sup>1</sup>H

NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.4 (s, 1H), 10.2 (s, br, 1H), 8.13 (s, 1H), 7.86 (d, 2H), 7.78 (d, 1H), 7.69 (d, 1H), 7.50 (t, 1H), 7.35 (dd, 2H), 2.89 (m, 2H), 2.72 (m, 2H), 1.90 (s, br, 4H) ppm; MS (ES+) 410.24 (M+H); (ES-) 408.23 (M-H); HPLC-Method A, R<sub>t</sub> 3.19 min.

Example 357 (7-Fluoro-1H-indazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-129): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.6 (s, 1H), 11.1 (s, br, 1H), 8.65 (d, 1H), 8.03 (d, 2H), 7.95 (s, 2H), 7.67 (m, 1H), 7.45 (m, 2H), 7.33 (t, 2H), 7.22 (dd, 1H), 6.99 (td, 1H) ppm. MS (ES+): m/e= 356.20 (M+H); HPLC-Method A R<sub>t</sub> 3.00 min.

Example 358 (5-Fluoro-1H-indazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-130): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.2 (s, 1H), 11.3 (s, br, 1H), 8.67 (d, 1H), 8.04 (d, 2H), 7.96 (s, 2H), 7.70 (m, 1H), 7.58 (dd, 1H), 7.43 (m, 4H), 7.28 (td, 1H) ppm. MS (ES+) 356.20 (M+H); HPLC-Method A, R<sub>t</sub> 3.00 min.

Example 359 (5,7-Difluoro-1H-indazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-131): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.7 (s, 1H), 8.65 (d, 1H), 8.04 (d, 2H), 7.95 (s, 2H), 7.68 (m, 1H), 7.45 (m, 1H), 7.35 (m, 4H) ppm. MS (ES+): m/e= 374.17 (M+H); HPLC-Method A, R<sub>t</sub> 3.07 min.

20

Example 360 (1H-Indazol-3-yl)-[2-(3-trifluoromethyl-phenyl)-quinazolin-4-yl]-amine (III-132): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 7.06 (t, 1H), 7.42 (t, 1H), 7.59 (d, 1H), 7.63 (t, 1H), 7.66 (d, 1H), 7.71 (m, 1H), 7.80 (d, 1H), 7.98 (m, 2H), 8.33 (s, 1H), 8.46 (d, 1H), 8.71 (d, 1H), 11.04 (br. s, 1H), 12.97 (s, 1H); EI-MS 406.1 (M+1); HPLC-Method A, R<sub>t</sub> 3.15 min.

Example 361 (2-Phenyl-quinazolin-4-yl)-(1H-pyrazolo[4,3-b]pyridin-3-yl)-amine (III-133): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.3 (s, br, 1H), 11.4 (s, br, 1H), 8.78 (d, 1H), 8.58 (dd, 1H), 8.24 (d, 1H), 8.10 (m, 2H), 7.95 (d, 2H), 7.86

(t, 1H), 7.56 (m, 2H), 7.44 (t, 2H) ppm. MS (ES+) 339.11 (M+H); HPLC-Method A, R<sub>t</sub> 2.63 min.

Example 362 [5-(3-Methoxy-phenyl)-6-oxo-5,6-dihydro-1H-

5 pyrazolo[4,3-c]pyridazin-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-134): <sup>1</sup>H NMR (500 MHz, MeOH-d<sub>4</sub>) δ8.65 (d, 1H), 8.17 (m, 3H), 8.10 (d, 1H), 7.90 (t, 1H), 7.75 (t, 1H), 7.58 (m, 2H), 7.25 (t, 1H), 6.95 (m, 2H), 6.85 (d, 1H), 6.80 (s, 1H), 3.64 (s, 3H) ppm. MS (ES+): m/e= 10 462.2 (M+H).

Example 363 (6-Oxo-5-phenyl-5,6-dihydro-1H-pyrazolo[4,3-c]pyridazin-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (III-

15 135): <sup>1</sup>H NMR (500 MHz, MeOH-d<sub>4</sub>) δ8.61 (d, 1H), 8.13 (m, 3H), 8.05 (d, 1H), 7.85 (t, 1H), 7.70 (t, 1H), 7.58 (m, 2H), 7.32 (m, 5H), 6.79 (s, 1H) ppm. MS (ES+): m/e= 432.2 (M+H).

Example 364 [5-(4-Methoxy-phenyl)-6-oxo-5,6-dihydro-1H-

20 pyrazolo[4,3-c]pyridazin-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-136): MS (ES+) 462.2 (M+H).

Example 365 [5-(2,4-Dichloro-phenyl)-6-oxo-5,6-dihydro-1H-pyrazolo[4,3-c]pyridazin-3-yl]-(2-phenyl-quinazolin-4-

25 yl)-amine (III-137): <sup>1</sup>H NMR (500 MHz, MeOH-d<sub>4</sub>) δ8.63 (d, 1H), 8.17 (m, 4H), 7.89 (t, 1H), 7.73 (t, 1H), 7.61 (t, 2H), 7.57 (d, 1H), 7.32 (m, 1H), 7.21 (d, 1H), 6.84 (s, 1H) ppm. MS (ES+): m/e= 500.1 (M+H).

30 Example 366 [6-Oxo-5-(3-trifluoromethyl-phenyl)-5,6-dihydro-1H-pyrazolo[4,3-c]pyridazin-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-138): <sup>1</sup>H NMR (500 MHz, MeOH-d<sub>4</sub>) δ8.55 (d, 1H), 8.19 (d, 2H), 7.92 (m, 2H), 7.65 (m,

3H), 7.45 (t, 2H), 7.25 (t, 1H), 7.13 (t, 1H), 7.05 (t, 1H), 6.75 (s, 1H) ppm. MS (ES+): m/e= 500.2 (M+H).

Example 367 [6-Oxo-5-(4-Phenoxy-phenyl)-5,6-dihydro-1H-pyrazolo[4,3-c]pyridazin-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-139): MS (ES+) 524.3 (M+H).

Example 368 [5-(4-Chloro-phenyl)-6-oxo-5,6-dihydro-1H-pyrazolo[4,3-c]pyridazin-3-yl]-(2-phenyl-quinazolin-4-yl)-amine (III-140): MS (ES+) 466.2 (M+H).

Example 369 (2-imidazol-1-yl-quinazolin-4-yl)-(1H-indazol-3-yl)-amine (III-141): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 7.10 (t, 1H), 7.44 (t, 1H), 7.50 (br. s, 1H), 7.60 (d, 1H), 7.72 (m, 2H), 7.77 (m, 1H), 7.88 (d, 1H), 7.98 (t, 1H), 8.73 (d, 1H), 8.96 (s, 1H), 11.23 (s, 1H), 13.06 (s, 1H); EI-MS 328.1 (M+1); HPLC-Method A, R<sub>t</sub> 2.93 min.

Example 370 (1H-Indazol-3-yl)-[2-(2-methyl-imidazol-1-yl-quinazolin-4-yl)-amine (III-142): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 2.48 (s, 3H), 7.10 (t, 1H), 7.43 (t, 1H), 7.57 (d, 1H), 7.60 (d, 1H), 7.67 (d, 1H), 7.76 (td, 1H), 7.86 (d, 1H), 7.91 (d, 1H), 8.01 (td, 1H), 8.72 (d, 1H), 11.15 (s, 1H), 13.10 (s, 1H); EI-MS 342.1 (M+1); HPLC-Method A, R<sub>t</sub> 3.06 min.

Example 371 (1H-Indazol-3-yl)-(2-piperidin-1-yl-quinazolin-4-yl)-amine (III-143): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 1.48 (m, 6H), 3.60 (m, 4H), 7.11 (t, 1H), 7.52 (t, 1H), 7.55 (d, 1H), 7.64 (d, 1H), 7.69 (d, 1H), 7.75 (d, 1H), 7.90 (t, 1H), 8.58 (d, 1H), 11.82 (br. s, 1H), 13.25 (s, 1H); EI-MS 345.1 (M+1); HPLC-Method A, R<sub>t</sub> 3.03 min.

Example 372 (1*H*-Indazol-3-yl)-[2-(octahydro-quinolin-1-yl)-quinazolin-4-yl]-amine (III-144): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 0.6-1.9 (m, 13 H), 3.15 (m, 1H), 3.25 (m, 1H), 4.0 (m, 1H), 7.10 (t, 0.5H), 7.12 (t, 0.5H), 7.55 (m, 2H), 7.66 (d, 0.5 H), 7.69 (d, 0.5 H), 7.77 (d, 1H), 7.91 (t, 1H), 8.55 (d, 0.5 H), 8.59 (d, 0.5 H), 11.46 (s, 0.5 H), 11.54 (s, 0.5 H), 11.78 (s, 0.5 H), 11.84 (s, 0.5 H), 13.10 (s, 0.5 H), 13.12 (s, 0.5 H); EI-MS 399.3 (M+1); HPLC-Method A, R<sub>t</sub> 3.37 min.

10

Example 373 (1*H*-Indazol-3-yl)-[2-(2,6-dimethyl-morpholin-4-yl)-quinazolin-4-yl]-amine (III-145): <sup>1</sup>H NMR (500MHz, DMSO-d<sub>6</sub>) δ 1.0 (m, 6H), 4.0 (m, 6H), 7.12 (t, 1H), 7.41 (td, 1H), 7.56 (t, 1H), 7.58 (d, 1H), 7.68 (dd, 1H), 7.77 (t, 1H), 7.93 (t, 1H), 8.60 (d, 1H), 11.69 (s, 1H), 13.16 (s, 1H); EI-MS 375.3 (M+1); HPLC-Method A, R<sub>t</sub> 2.93 min.

15

Example 374 (5-Methyl-2*H*-pyrazol-3-yl)-(2-phenyl-pyrimidin-4-yl)-amine (IV-1): mp 245-246°C; <sup>1</sup>H NMR (DMSO) δ 2.26 (3H, s), 6.32 (1H, br s), 7.07 (1H, br s), 7.48-7.54 (3H, m), 8.33-8.39 (3H, m), 9.87 (1H, s), 12.03 (1H, s); IR (solid) 1628, 1589, 1579, 1522, 1479, 1441, 1393, 1336; MS 252.2 (M+H)<sup>+</sup>.

20

Example 375 [6-(4-Acetamidophenylsulfanyl)-2-phenyl-pyrimidin-4-yl]-(5-methyl-2*H*-pyrazol-3-yl)-amine (IV-3): A suspension of Fenclozime (4,6-dichloro-2-phenylpyrimidine) (0.1g, 0.44 mmol), 3-amino-5-methylpyrazole (0.045 g, 0.47 mmol), *N,N*-diisopropylethylamine (0.08 ml, 0.47 mmol) and sodium iodide (0.067 g, 0.44 mmol) in *n*-butanol (5 ml) were heated at 117 °C for 18 hours. The solvent was removed in vacuo and the crude product purified by flash chromatography (silica gel, 3:2 Petrol:EtOAc) to afford

30

0.037 g (29 % yield) of (6-Chloro-2-phenyl-pyrimidin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine as a off-white solid. A suspension of the above pyrimidine (0.037 g, 0.13 mmol) and thioacetamidothiophenol (0.108 g, 0.64 mmol) in *tert*-butanol was heated at 85 °C under nitrogen for 2 days. The reaction mixture was cooled to room temperature and the solvent removed *in vacuo*. The concentrate was dissolved in EtOAc, and washed with NaHCO<sub>3</sub> (sat, aq.). The organic layer is concentrated in *vacuo*, and the crude product by preparative HPLC. The residual disulfide that still remained in the mixture after HPLC may be removed by precipitation from EtOAc and filtration. The mother liquor was concentrated to afford IV-3 (7mg, 13 % yield) as an off-white solid: mp 235-236°C; <sup>1</sup>H NMR (DMSO) δ 2.10 (3H, s), 2.21 (3H, s), 6.33 (1H, br s), 7.50 (3H, m), 7.7-7.59 (2H, m), 7.76-7.78 (2H, m), 8.25 (2H, m), 9.72, 10.26 and 11.93 (3 H, 3 x br s); IR (solid) 1669, 1585, 1551, 1492, 1392, 1372, 1312, 1289, 1259, 1174, 1102, 1089, 1027, 1015, 984; MS 417.3 (M+H)<sup>+</sup>.

Example 376 [2-(4-Methylpiperidin-1-yl)-pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-4): mp 215-216°C; <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 0.96 (3H, d), 1.16 (2H, m), 1.66 (3H, m), 2.27 (3H, s), 2.86 (2H, t), 4.58 (2H, m), 4.78 (2H, exch.protons), 6.13 (2H, m), 7.83 (1H, d); IR (solid) 1593, 1550, 1489, 1436, 1331, 1246, 1231; MS 273.1 (M+H)<sup>+</sup>.

Example 377 [2-(4-Methylpiperidin-1-yl)-5-nitropyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-5): mp 185-187°C; <sup>1</sup>H NMR (DMSO) δ 0.93 (3H, d), 1.06-1.18 (2H, m), 1.68-1.80 (3H, m), 2.26 (3H, s), 3.01-3.12 (2H, m), 4.63 (1H, d), 4.80 (1H, d), 6.39 (1H, s), 9.00 (1H, s), 10.41

(1H, s), 12.36 (1H, s); IR (solid) 1589, 1517, 1479, 1446, 1346, 1317, 1246, 1222, 1055; MS 318.2 (M+H)<sup>+</sup>.

Example 378 [5-Amino-2-(4-Methylpiperidin-1-yl)-

5 pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-6):

To a solution of IV-5 (48 mg, 0.151 mmol) in ethanol (2.0 mL) was added tin dichloride dihydrate (171 mg, 0.756 mmol) and the resulting mixture heated at reflux for 3 hours. The reaction was cooled to room temperature and

10 poured onto a mixture of 1M NaOH:dichloromethane:propanol

(18:8:4mL) and stirred for 15 minutes. The layers were separated and the aqueous layer extracted twice with

dichloromethane. The combined organic layers were

concentrated in vacuo and the residue purified by flash

15 chromatography (silica gel, gradient

dichloromethane:MeOH) to afford IV-6 as a grey solid

(27mg, 63%): <sup>1</sup>H NMR (DMSO) δ 0.88-1.04 (5H, m), 1.55-1.62

(3H, m), 2.21 (3H, s), 2.70 (2H, m), 3.36 (2H, m), 4.40

(2H, m), 6.37 (1H, s), 7.49 (1H, s), 8.40 (1H, s), 11.92

20 (1H, br s); MS 288.2 (M+H)<sup>+</sup>.

Example 379 [5-Amino-6-methyl-2-(4-methylpiperidin-1-yl)-pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-7):

mp 172-175°C; <sup>1</sup>H NMR (DMSO) δ 0.90 (3H, d), 1.03 (2H, m),

25 1.52-1.62 (3H, m), 2.13 (3H, s), 2.20 (3H, s), 2.69 (2H,

m), 3.92 (2H, br s), 4.44 (2H, d), 6.35 (1H, s), 8.41

(1H, s), 11.85 (1H, br s); IR (solid) 1612, 1589, 1489,

1446, 1317; MS 302.5 (M+H)<sup>+</sup>.

30 Example 380 [6-Methyl-2-(4-methyl-phenyl)-pyrimidin-4-yl]-(5-phenyl-2H-pyrazol-3-yl)-amine (IV-10): MS 342.34

(M+H); HPLC-Method E, R<sub>t</sub> 1.334 min.

Example 381 [2-(4-Chloro-phenyl)-6-methyl-pyrimidin-4-yl]-(5-furan-2-yl-2H-pyrazol-3-yl)-amine (IV-11): MS 352.11 (M+H); HPLC Method E, R<sub>t</sub> 1.194 min.

5 Example 382 5-Furan-2-yl-2H-pyrazol-3-yl)-(6-methyl-2-phenyl-pyrimidin-4-yl)-amine (IV-12): MS 318.21 (M+H); HPLC-Method E, 1.192 min.

Example 383 [6-Methyl-2-(4-trifluoromethyl-phenyl)-pyrimidin-4-yl]-(5-phenyl-2-yl-2H-pyrazol-3-yl)-amine (IV-13): MS 396.24 (M+H); HPLC-Method E, R<sub>t</sub> 1.419 min.

Example 384 (5-Furan-2-yl-2H-pyrazol-3-yl)-[6-methyl-2-(4-trifluoromethyl-phenyl)-pyrimidin-4-yl]-amine (IV-14): MS 386.08 (M+H); HPLC-Method E 1.347 min.

Example 385 [2-(2,3-Dihydro-benzo[1,4]dioxin-2-yl)-6-methyl-pyrimidin-4-yl]-(5-furan-2-yl-2H-pyrazol-3-yl)-amine (IV-15): MS 376.18 (M+H); HPLC-Method E, R<sub>t</sub> 1.181 min.

Example 386 [2-(2,3-Dihydro-bezo[1,4]dioxin-2-yl)-6-ethyl-pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-16): MS 338.17 (M+H); HPLC-Method E, R<sub>t</sub> 1.082 min.

Example 387 (6-Ethyl-2-phenyl-pyrimidin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (IV-17): MS 280.18 (M+H); HPLC-Method E, R<sub>t</sub> 1.024 min.

30 Example 388 (6-Methyl-2-phenyl-pyrimidin-4-yl)-(5-phenyl-2H-pyrazol-3-yl)-amine (IV-19): MS 328.51 (M+H); HPLC-Method E, R<sub>t</sub> 1.192 min.



Example 389 [6-Ethyl-2-(4-trifluoromethyl-phenyl)-pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-20): MS 348.5 (M+H); HPLC-Method E,  $R_t$  1.224 min.

- 5 Example 390 (5-Furan-2-yl-2H-pyrazol-3-yl)-[6-methyl-2-(4-methyl-phenyl)-pyrimidin-4-yl]-amine (IV-21): MS 332.23 (M+H); HPLC-Method E,  $R_t$  1.139 min.

- 10 Example 391 (6-Methoxymethyl-2-phenyl-pyrimidin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (IV-22): MS 296.31 (M+H); HPLC-Method E,  $R_t$  0.971 min.

- 15 Example 392 (5,6-Dimethyl-2-phenyl-pyrimidin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (IV-23): MS 280.2 (M+H); HPLC-Method E,  $R_t$  0.927 min.

- 20 Example 393 (6-Methyl-2-phenyl-pyrimidin-4-yl)-(5-methyl-2H-pyrazol-3-yl)-amine (IV-24): MS 266.18 (M+H); HPLC-Method E,  $R_t$  0.925 min.

Example 394 [6-Ethyl-2-(4-methyl-phenyl)-pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-25): MS 294.46 (M+H); HPLC-Method E,  $R_t$  1.174 min.

- 25 Example 395 [2-(4-Chloro-phenyl)-6-ethyl-pyrimidin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-26): MS 314.42 (M+H); HPLC-Method E  $R_t$  1.213 min.

- 30 Example 396 (5-Methyl-1H-pyrazol-3-yl)-(6-methyl-2-p-tolyl-pyrimidin-4-yl)-amine (IV-27): MS 280.45 (M+H); HPLC-Method E,  $R_t$  1.135 min.

Example 397 (1H-Indazol-3-yl)-(6-methoxymethyl-2-phenyl-pyrimidin-4-yl)-amine (IV-28):  $^1\text{H}$  NMR (500 MHz, DMSO)  $\delta$

3.57 (3H, s), 4.65 (2H, s), 7.23 (1H, J=7.5 Hz, t), 7.52 (1H, J=7.6 Hz, t), 7.63 (4H, m), 7.75 (1H, br), 8.13 (1H, J=5.5 Hz, br d), 8.44 (1H, J=5.7 Hz, br d), 10.6 (1H, br), 12.8 (1H, br s) ppm; HPLC-Method A,  $R_t$  2.944 min; MS (FIA) 332.1 (M+H)<sup>+</sup>.

Example 398 (5-Methyl-2H-pyrazol-3-yl)-(2-pyridin-4-yl-thieno[3,2-d]pyrimidin-4-yl)-amine (IV-29): <sup>1</sup>H NMR (DMSO)  $\delta$  2.34 (3H, s), 6.66 (1H, s), 7.53 (1H, d), 7.84 (1H, d), 8.32 (2H, d), 8.70 (2H, d); MS 309.6 (M+H)<sup>+</sup>.

Example 399 (5-Methyl-2H-pyrazol-3-yl)-(2-phenyl-pyrido[3,4-d]pyrimidin-4-yl)-amine (IV-30): mp 225°C; <sup>1</sup>H NMR (DMSO)  $\delta$  2.35 (3H, s), 6.81 (1H, s), 7.50-7.63 (3H, m), 8.45-8.52 (2H, m), 8.54 (1H, d), 8.62 (1H, d), 9.20 (1H, s), 10.79 (1H, s), 12.38 (1H, br s); IR (solid) 2958, 2917, 2852, 1593, 1565, 1524, 1467, 1450; MS 303.2 (M+H)<sup>+</sup>.

Example 400 (5-Methyl-2H-pyrazol-3-yl)-(2-phenyl-pyrido[2,3-d]pyrimidin-4-yl)-amine (IV-31):  
To a solution of 4-chloro-2-phenyl-pyrido[2,3-d]pyrimidine (J. Pharm. Belg., 29, 1974, 145-148) (109mg, 0.45 mmol) in THF (15 mL) was added 3-amino-5-methylpyrazole (48 mg, 0.5 mmol) and the resulting mixture heated at 65 °C overnight. The mixture was cooled to room temperature and the resulting suspension was filtered and washed with Et<sub>2</sub>O. The solid was dissolved in a mixture EtOH:water and the pH adjusted to pH 7. The aqueous was extracted twice with ethyl acetate and the combined organic layers were dried (MgSO<sub>4</sub>), filtered, and concentrated in vacuo. The residue was purified by flash chromatography (SiO<sub>2</sub>, DCM-MeOH gradient) to afford IV-31 as an off-white solid (69 mg, 50%): mp 234°C; <sup>1</sup>H NMR

(DMSO)  $\delta$  2.14 (3H, s), 5.99 (1H, s), 7.20-7.40 (3H, m), 7.40-7.50 (3H, m), 8.60 (1H, d), 8.79 (1H, d), 12.82 (1H, br s); IR (solid) 2957, 2921, 2857, 1644, 1560, 1459, 1427; MS 303.2 (M+H)<sup>+</sup>.

5

Example 401 (5-Cyclopropyl-2H-pyrazol-3-yl)-(2-phenyl-pyrido[3,4-d]pyrimidin-4-yl)-amine (IV-32): off-white solid, mp 232-233°C; <sup>1</sup>H NMR (DMSO)  $\delta$  0.70-0.85 (2H, m), 0.90-1.05 (2H, m), 1.05-2.07 (1H, m), 6.75 (1H, s), 7.50-10 7.75 (3H, m), 8.40-8.70 (4H, m), 9.20 (1H, s), 10.80 (1H, s), 12.41 (1H); IR (solid) 3178, 1601, 1573, 1532, 1484, 1452, 1409, 1367, 1328, 802, 781, 667; MS 329.2 (M+H)<sup>+</sup>.

Example 402 [2-(4-Methylpiperidin-1-yl)-purin-4-yl]-(5-methyl-2H-pyrazol-3-yl)-amine (IV-33): To a suspension of 15 2,4-dichloro-purine (2.0 g, 10.6 mmol) in anhydrous ethanol (10 mL) was added 5-methyl-1H-pyrazol-3-yl amine (2.05 g, 21.2 mmol). The resulting mixture was stirred at room temperature for 48 h. The resulting precipitate 20 was collected by filtration, washed with ethanol, and dried under vacuum to afford 1.524 g (58% yield) of (2-chloro-purin-4-yl)-(5-methyl-1H-pyrazol-3-yl)-amine which was used in the next step without further purification. To a solution of (2-chloro-purin-4-yl)-(5-methyl-1H-25 pyrazol-3-yl)-amine (200 mg, 0.80 mmol) was added 4-methylpiperidine (4 mL, 8.01 mmol) and the reaction mixture heated at reflux overnight. The solvent was evaporated and the residue dissolved in a mixture EtOH:water (1:3, 4 mL). Potassium carbonate (57mg, 0.41 30 mmol) was added and the mixture was stirred at room temperature for 2 hours. The resulting suspension was filtered, washed with water (x2) and rinsed with Et<sub>2</sub>O (x2) to afford IV-33 as a white solid (225mg, 90%): mp >300°C; <sup>1</sup>H NMR (DMSO)  $\delta$  0.91 (3H, d), 1.10 (2H, m), 1.65 (3H, m),

2.24 (3H, s), 2.84 (2H, m), 4.60 (2H, m), 6.40 (1H, s), 7.87 (1H, m), 9.37-9.59 (1H, m), 12.03-12.39 (2H, m); IR (solid) 1651, 1612, 1574, 1484, 1446, 1327, 1317, 1255, 1203; MS 313.3 (M+H)<sup>+</sup>.

5

Example 403 (5-Cyclopropyl-2H-pyrazol-3-yl)-[2-(4-methylpiperidin-1-yl)-pyrrolo[3,2-d]pyrimidin-4-yl]-amine (IV-34): white solid; <sup>1</sup>H NMR (DMSO) δ 0.65 (2H, m), 0.91-0.96 (5H, m), 1.08 (2H, m), 1.58-1.64 (3H, m), 1.89 (1H, m), 2.77 (2H, t), 4.57 (2H, d), 6.09 (1H, s), 6.38 (1H, s), 7.33 (1H, s), 9.42 (1H, s), 10.65 (1H, s), 12.02 (1H, br s); MS 338.3 (M+H)<sup>+</sup>.

Example 404 [6-Benzyl-2-phenyl-5,6,7,8-tetrahydro-pyrido[4,3-d]pyrimidin-4-yl]-(5-fluoro-1H-indazol-3-yl)-amine (IV-35): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.0 (s, 1H), 10.4 (s, br, 1H), 9.73 (s, 1H, TFA-OH), 8.00 (d, 2H), 7.64 (m, 2H), 7.59 (dd, 1H), 7.52 (m, 3H), 7.41 (t, 1H), 7.31 (m, 3H), 7.14 (dd, 1H), 4.58 (s, 2H), 4.35 (br, 2H), 3.74 (m, 2H), 3.17 (s, 2H) ppm. MS (ES<sup>+</sup>): m/e = 451.30 (M+H); HPLC-Method A, T<sub>ret</sub> 2.96 min.

Example 405 (5-Fluoro-1H-indazol-3-yl)-(2-phenyl-5,6,7,8-tetrahydro-pyrido[4,3-d]pyrimidin-4-yl)-amine (IV-36): Prepared from IV-35 (0.13 mmol) by treatment with an equal weight of Pd/C (10%) in 4.4% HCOOH in MeOH at room temperature for 12 h. The mixture was filtered through celite, the filtrate was evaporated, and crude product was purified by HPLC to afford IV-36 as yellow solid in 35% yield. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1H), 9.06 (s, 1H), 7.99 (d, 2H), 7.57 (dd, 1H), 7.34 (m, 1H), 7.28 (m, 3H), 7.22 (d, 1H), 3.83 (s, 2H), 3.05 (m, 2H), 2.72 (m, 2H) ppm. MS (ES<sup>+</sup>): m/e = 361.20 (M+H); HPLC-Method A, T<sub>ret</sub> 2.68 min.

Example 406 (5-Methyl-2H-pyrazol-3-yl)-(3-phenyl-

isoquinolin-1-yl)-amine (V-1): To a solution of 1-chloro-3-phenylisoquinoline (J. Het. Chem., 20, 1983, 121-128) (0.33g, 1.37 mmol) in DMF (anhydrous, 5 mL) was added 3-amino-5-methylpyrazole (0.27g, 2.74 mmol) and potassium carbonate (0.57g, 4.13 mmol) and the resulting mixture was heated at reflux for 6 hours. The reaction mixture was then cooled and solvent removed in vacuo. The residue was extracted twice with ethyl acetate and the combined organic layers washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated in vacuo. The crude product was purified by flash chromatography (SiO<sub>2</sub>, gradient DCM-MeOH) to afford V-1 as a colourless oil; <sup>1</sup>H NMR (MeOD) δ 2.23 (3H, s), 5.61 (1H, s), 7.41 (1H, m), 7.52 (2H, m), 7.62 (1H, m), 7.81 (1H, m), 8.07 (1H, d), 8.19 (2H, m), 8.29 (1H, s), 8.54 (1H, d); MS 301.2 (M+H)<sup>+</sup>.

Example 407 (1H-Indazol-3-yl)-[3-(2-trifluoromethyl-

phenyl)-isoquinoline-1-yl]-amine (V-2): A solution of 1-chloro-3-(2-trifluoromethyl-phenyl)-isoquinoline (100 mg, 0.326 mmol) and 1H-indazol-3-ylamine (86 mg, 0.651 mmol) in ethanol (3 mL) was heated at 160 C and the solvent evaporated with a stream of nitrogen. The remaining oil was then heated at 160 C for 18 hours under nitrogen. The resulting melt was dissolved in 5% methanol:dichloromethane (50 mL), washed with saturated aqueous sodium bicarbonate (1 x 25 mL) then dried over magnesium sulfate. Purification by silica gel chromatography (25% to 50% hexane:ethyl acetate) afforded V-2 as a yellow solid (35 mg, 27%). <sup>1</sup>H NMR (500 MHz, d<sub>6</sub>-DMSO) δ 9.78 (br s, 1H), 8.62 (d, 1H), 7.9-7.85 (m, 1H), 7.78-7.72 (m, 1H), 7.70-7.68 (m, 1H), 7.65-7.62 (m, 1H), 7.60-7.55 (m, 1H), 7.52-7.45 (m, 3H), 7.41-7.38 (m, 1H),

7.28-7.25 (m, 1H), 7.18 (s, 1H), 6.95-6.92 (m, 1H), 5.76 (s, 1H); LC-MS (ES+) m/e= 405.18 (M+H); HPLC-Method D R<sub>t</sub> 2.74 min.

- 5 Example 408 (5,7-Difluoro-1H-indazol-3-yl)-[3-(2-trifluoromethyl-phenyl)-isoquinolin-1-yl]-amine (V-3): Prepared from 5,7-difluoro-1H-indazol-3-ylamineto afford compound V-3 as a yellow solid (90 mg, 63%). <sup>1</sup>H NMR (500 MHz, d<sub>6</sub>-DMSO) δ 13.25 (s, 1H), 9.92 (br s, 1H), 8.61 (d, 1H), 7.9 (d, 1H), 7.81-7.49 (m, 6H), 7.26-7.2 (m, 2H), 7.12-7.10 (m, 1H); LC-MS (ES+) m/e= 441.16 (M+H); HPLC-Method D, R<sub>t</sub> 3.58 min.

- Example 409 (5-Methyl-2H-pyrazol-3-yl)-(2-phenyl-quinolin-4-yl)-amine (V-4): To a mixture of 4-chloro-2-phenylquinoline (J. Het. Chem., 20, 1983, 121-128) (0.53g, 2.21 mmol) in diphenylether (5 mL) was added 3-amino-5-methylpyrazole (0.43g, 4.42 mmol) and the resulting mixture heated at 200°C overnight with stirring. The reaction mixture was cooled to ambient temperature then petroleum ether (20 mL) was added and the resulting precipitate was isolated by filtration. The crude solid was purified by flash chromatography (SiO<sub>2</sub>, gradient DCM-MeOH) to afford V-4 as a white solid: mp 242-244°C; <sup>1</sup>H NMR (DMSO) δ 2.27(3H, s), 6.02(1H, s), 7.47(2H, d), 7.53-7.40(2H, br m), 7.67(1H, m), 7.92(1H, m), 8.09(2H, d), 8.48(2H, m), 9.20(1H, s), 12.17(1H, br s); IR (solid) 1584, 1559, 1554, 1483, 1447, 1430, 1389; MS 301.2 (M+H)<sup>+</sup>.

- 30 Example 410 (1H-Indazol-3-yl)-(2-phenyl-quinolin-4-yl)-amine (V-5): <sup>1</sup>H NMR (500 MHz, d<sub>6</sub>-DMSO) δ 12.78 (s, 1H), 9.50 (s, 1H), 8.65 (d, 1H), 8.15 (s, 1H), 8.04-7.98 (m, 3H), 7.94 (s, 1H), 7.78-7.75 (m, 1H), 7.60-7.40 (m, 6H),

7.15-7.10 (m, 1H). LC-MS (ES+) m/e= 337.11 (M+H); HPLC-Method D, R<sub>t</sub> 2.10 min.

Example 411 (2-Phenyl-quinolin-4-yl)-(1H-pyrazolo[4,3-b]pyridin-3-yl)-amine (V-6): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.6 (s, 1H), 11.4 (s, 1H), 8.94 (d, 1H), 8.61 (dd, 1H), 8.23 (d, 1H), 8.16 (dd, 1H), 8.12 (t, 1H), 7.89 (t, 1H), 7.86 (d, 1H), 7.65 (m, 4H), 7.54 (s, 1H), 7.52 (dd, 1H) ppm. MS (ES+): m/e= 338.11 (M+H); HPLC-Method A, HPLC-Method D, R<sub>t</sub> 2.91 min.

Example 412 (1H-Indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinolin-4-yl]-amine (V-7): <sup>1</sup>H NMR (500 MHz, d<sub>6</sub>-DMSO) δ 12.68 (s, 1H), 9.51 (s, 1H), 8.7 (d, 1H), 7.95-7.89 (m, 2H), 7.83-7.70 (m, 3H), 7.68-7.62 (m, 2H), 7.60 (s, 1H), 7.55-7.52 (m, 1H), 7.49-7.45 (m, 1H), 7.40-7.37 (m, 1H), 7.12-7.09 (m, 1H); LC-MS (ES+) m/e= 405.15 (M+H); HPLC-Method D R<sub>t</sub> 2.25 min.

Example 413 (5,7-Difluoro-1H-indazol-3-yl)-[2-(2-trifluoromethyl-phenyl)-quinolin-4-yl]-amine (V-8): <sup>1</sup>H NMR (500 MHz, d<sub>6</sub>-DMSO) δ 13.31 (s, 1H), 9.49 (s, 1H), 8.70-8.67 (m, 1H), 7.96-7.92 (m, 1H), 7.85-7.66 (m, 7H), 7.63-7.60 (m, 1H), 7.42-7.40 (m, 1H). LC-MS (ES+) m/e= 441.18 (M+H); HPLC-Method D R<sub>t</sub> 2.39 min.

Example 414 [2-(2-trifluoromethyl-phenyl)-quinolin-4-yl]-(1H-pyrazolo[4,3-b]pyridin-3-yl)-amine (V-9): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.6 (s, 1H), 11.6 (s, br, 1H), 8.98 (d, 1H), 8.57 (dd, 1H), 8.12 (m, 3H), 7.97 (m, 2H), 7.86 (m, 3H), 7.49 (dd, 1H), 7.23 (s, 1H) ppm. MS (ES+): m/e= 406.20 (M+H); HPLC-Method A R<sub>t</sub> 2.91 min.

Example 415 (2-Phenyl-quinazolin-4-yl)-(2H-

[1,2,4]triazol-3-yl)-amine (IX-154): off-white solid, mp 266-267°C; <sup>1</sup>H NMR (DMSO) δ 7.50-7.70 (4H, m), 7.85-8.00 (2H, m), 8.15-8.25 (2H, m), 8.37-8.45 (2H, m), 8.58 (1H, d), 13.90 (1H, br s); IR (solid) 3344, 3059, 1630, 1609, 1570, 1557, 1543, 1501, 1495, 1445, 1411, 1355, 1326, 1267, 1182, 1053, 1038, 760, 676, 667, 654; MS 289.2 (M+H)<sup>+</sup>.

10 Example 416 (5-Methyl-2H-[1,2,4]triazol-3-yl)-(2-phenyl-quinazolin-4-yl)-amine (IX-155): <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.59 (s, 1H), 8.42 (d, J = 6.7 Hz, 2H), 7.79 (m, 4H), 8.03 (m, 2H), 7.74 (m, 4H), 2.51 (s, 3H) ppm. MS (ES<sup>+</sup>): m/e = 303.08 (M+H); HPLC-Method A, R<sub>t</sub> 2.64 min.

15

Example 417 (2H-[1,2,4]-Triazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (IX-47): Pale yellow solid (52% yield). <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.54 (s, 1H), 8.15 (s, br, 1H), 7.91 (t, 1H), 7.85 (m, 2H), 7.76 (m, 3H), 7.66 (t, 1H) ppm. MS (ES<sup>+</sup>): m/e = 357.13 (M+H); (ES<sup>-</sup>): m/e = 355.15 (M-H); HPLC-Method A, R<sub>t</sub> 2.81 min.

20

Example 418 (5-Methyl-2H-[1,2,4]triazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (IX-38): Pale yellow solid (54% yield). <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.44 (s, br, 1H), 7.92 (m, 3H), 7.84 (m, 1H), 7.77 (m, 2H), 7.68 (t, 1H), 2.28 (s, 3H) ppm. MS (ES<sup>+</sup>): m/e = 371.14 (M+H); (ES<sup>-</sup>): m/e = 369.18 (M-H); HPLC-Method A, R<sub>t</sub> 2.89 min.

25

30

Example 419 (5-Methylsulfanyl-2H-[1,2,4]triazol-3-yl)-[2-(2-trifluoromethylphenyl)-quinazolin-4-yl]-amine (IX-156): Pale yellow solid (65% yield). <sup>1</sup>H NMR (500 MHz,



DMSO-d<sub>6</sub>)  $\delta$  8.56 (br, 1H), 7.90 (t, 1H), 7.84 (m, 2H), 7.78 (m, 2H), 7.67 (m, 2H), 2.51 (s, 3H, buried by DMSO) ppm.  
MS (ES<sup>+</sup>): m/e = 403.12 (M+H); (ES<sup>-</sup>): m/e = 401.16 (M-H);  
HPLC-Method A, R<sub>t</sub> 3.20 min.

5

Example 420 (1H-[1,2,4]Triazol-3-yl)-[3-(2-trifluoromethyl-phenyl)-isoquinolin-1-yl]-amine (IX-175):

A solution of 1-chloro-3-(2-trifluoromethyl-phenyl)-isoquinoline (0.326 mmol) and 1H-[1,2,4]triazol-3-ylamine  
10 (0.651 mmol) in ethanol (3 mL) was heated at 160°C and the solvent evaporated with a stream of nitrogen. The remaining oil was then heated at 160°C for 18 hours under nitrogen. The resulting melt was dissolved in 5% methanol/dichloromethane (50 mL), washed with saturated  
15 aqueous sodium bicarbonate (1 x 25 mL) then dried over magnesium sulfate. Purification by silica gel chromatography afforded IX-175 as a colorless oil (4% yield). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.18 (d, 1H), 8.82 (s, 1H), 7.90 (d, 1H), 7.85-7.75 (m, 3H), 7.71-7.62 (m, 3H),  
20 7.60-7.55 (m, 2H), 4.42-4.35 (m, 1H). LC-MS (ES<sup>+</sup>) 356.16 (M+H); HPLC-Method D, R<sub>t</sub> 3.55 min.

Example 421 (2-Phenyl-quinolin-4-yl)-(1H-[1,2,4]triazol-3-yl)-amine (IX-176): Pale yellow solid (30% yield). <sup>1</sup>H

25 NMR (500 MHz, d<sub>6</sub>-DMSO)  $\delta$  13.82 (s, 1H), 9.91 (s, 1H), 8.80 (s, 1H), 8.70-8.65 (m, 1H), 8.55 (s, 1H), 8.15-8.12 (m, 2H), 8.03-7.98 (m, 1H), 7.75-7.72 (m, 1H), 7.57-7.49 (m, 3H). LC-MS (ES<sup>+</sup>) m/e = 288.11 (M+H); HPLC-Method D, R<sub>t</sub> 1.55 min.

30

Example 422 (1H-[1,2,4]triazol-3-yl)-[2-(2-

trifluoromethyl-phenyl)-quinolin-4-yl]-amine (IX-177): Pale yellow solid (46% yield). <sup>1</sup>H NMR (500 MHz, d<sub>6</sub>-DMSO)  $\delta$  13.70 (s, 1H), 9.98 (s, 1H), 8.70 (d, 1H), 8.49 (s,

1H), 8.30 (s, 1H), 7.94-7.88 (m, 2H), 7.80-7.68 (m, 3H), 7.64-7.56 (m, 2H). LC-MS (ES+) m/e= 356.18 (M+H); HPLC-Method D, R<sub>t</sub> 1.68 min.

- 5 Example 423 (1-H-Indazol-3-yl)-[5-methyl-6-morpholin-4-yl-2-(2-trifluoromethyl-phenyl)-pyrimidin-4-yl]-amine (II-251): Colorless film; 2 % yield; <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD) δ 7.84 (m, 2H), 7.71 (m, 3H), 7.41 (t, 2H), 7.14 (m, 1H), 3.74 (m, 4H), 3.69 (m, 4H), 1.24 (s, 3H) ppm;
- 10 HPLC-Method A R<sub>t</sub> 3.26 min; MS (FIA) 455.1 (M+H).

### BIOLOGICAL TESTING

The activity of the compounds as protein kinase inhibitors may be assayed *in vitro*, *in vivo* or in a cell

15 line. *In vitro* assays include assays that determine inhibition of either the phosphorylation activity or ATPase activity of the activated protein kinase.

Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to the protein kinase. Inhibitor

20 binding may be measured by radiolabelling the inhibitor prior to binding, isolating the inhibitor/protein kinase complex and determining the amount of radiolabel bound.

Alternatively, inhibitor binding may be determined by running a competition experiment where new inhibitors are

25 incubated with the protein kinase bound to known radioligands.

### BIOLOGICAL TESTING EXAMPLE 1

#### K<sub>i</sub> DETERMINATION FOR THE INHIBITION OF GSK-3

30 Compounds were screened for their ability to inhibit GSK-3β (AA 1-420) activity using a standard coupled enzyme system (Fox et al. (1998) *Protein Sci.* 7, 2249). Reactions were carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM MgCl<sub>2</sub>, 25 mM NaCl,

300  $\mu$ M NADH, 1 mM DTT and 1.5% DMSO. Final substrate concentrations in the assay were 20  $\mu$ M ATP (Sigma Chemicals, St Louis, MO) and 300  $\mu$ M peptide (HSSPHQS(PO<sub>3</sub>H<sub>2</sub>)EDEEE, American Peptide, Sunnyvale, CA).

5 Reactions were carried out at 30 °C and 20 nM GSK-3 $\beta$ . Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 300  $\mu$ M NADH, 30  $\mu$ g/ml pyruvate kinase and 10  $\mu$ g/ml lactate dehydrogenase.

10 An assay stock buffer solution was prepared containing all of the reagents listed above with the exception of ATP and the test compound of interest. The assay stock buffer solution (175  $\mu$ l) was incubated in a 96 well plate with 5  $\mu$ l of the test compound of interest  
15 at final concentrations spanning 0.002  $\mu$ M to 30  $\mu$ M at 30 °C for 10 min. Typically, a 12 point titration was conducted by preparing serial dilutions (from 10 mM compound stocks) with DMSO of the test compounds in daughter plates. The reaction was initiated by the  
20 addition of 20  $\mu$ l of ATP (final concentration 20  $\mu$ M). Rates of reaction were obtained using a Molecular Devices Spectramax plate reader (Sunnyvale, CA) over 10 min at 30 °C. The K<sub>i</sub> values were determined from the rate data as a function of inhibitor concentration.

25 The following compounds were shown to have K<sub>i</sub> values less than 0.1  $\mu$ M for GSK-3: compounds II-1, II-105, II-33, II-34, II-36, II-39, II-38, II-39, II-40, II-41, II-42, II-46, II-57, II-59, II-60, II-61, II-62, II-63, II-64, II-66, II-67, II-69, II-70, II-53, II-71, II-30 99, II-73, II-74, II-75, II-76, II-77, II-7, II-8, II-9, II-10, II-24, II-19, II-78, II-54, II-79, II-80, II-81, II-82, II-83, II-84, II-56, II-86, II-20, II-25, II-26, II-85, II-21, II-27, II-28, II-87, II-88, II-29, II-11, II-12, II-30, II-31, II-13, II-14, II-15, II-16, II-17,

II-18, II-79, II-23, II-2, II-90, II-91, II-92, II-93,  
II-3, II-4, II-5, II-6, II-94, II-95, II-96, II-107, II-  
108, II-109, II-110, II-124, II-125, II-111, II-112, II-  
113, II-114, II-115, II-116, II-117, II-118, II-119, II-  
5 120, II-121, II-208, III-8, III-7, III-9, III-37, III-38,  
III-39, III-40, III-42, III-45, III-46, III-47, III-48,  
III-49, III-51, III-52, III-53, III-54, III-55, III-56,  
III-57, III-58, III-59, III-60, III-61, III-62, III-63,  
III-30, III-65, III-66, III-67, III-70, III-73, III-31,  
10 III-75, III-76, III-77, III-33, III-34, III-106, III-108,  
III-109, III-111, III-35, III-116, III-117, III-118, III-  
119, III-120, III-121, III-127, III-128, III-141, III-  
130, III-131, IV-15, IV-16, IV-17, IV-20, IV-25, IV-26,  
IV-30, IV-34, V-3, and IX-47.

15           The following compounds were shown to have  $K_i$   
values between 0.1 and 1.0  $\mu\text{M}$  for GSK-3: compounds II-  
103, II-104, II-35, II-44, II-45, II-49, II-50, II-97,  
II-101, II-22, II-32, III-41, III-43, III-44, III-28,  
III-50, III-29, III-64, III-71, III-74, III-78, III-82,  
20 III-88, III-90, III-102, III-105, III-107, III-110, III-  
112, III-114, III-115, III-122, III-124, III-124, IV-1,  
III-1, III-138, III-140, III-142, III-129, III-132, III-  
134, III-135, III-136, IV-1, IV-10, IV-11, IV-12, IV-13,  
IV-14, IV-19, IV-21, IV-22, IV-23, IV-24, IV-3, IV-4, IV-  
25 6, IV-7, IV-8, IV-29, IV-31, IV-32, IV-33, IV-36, V-2, V-  
7, IX-38, IX-154, and IX-177.

          The following compounds were shown to have  $K_i$   
values between 1.0 and 20  $\mu\text{M}$  for GSK-3: compounds II-43,  
II-65, II-48, II-47, II-51, II-68, II-52, II-72, II-100,  
30 II-98, II-89, III-68, III-81, III-83, III-91, III-94,  
III-95, III-96, III-97, III-98, III-99, III-100, III-101,  
III-103, III-123, III-137, III-139, III-143, III-145,  
III-146, V-4, V-8, IX-156, and IX-176.

BIOLOGICAL TESTING EXAMPLE 2K<sub>i</sub> DETERMINATION FOR THE INHIBITION OF AURORA-2

Compounds were screened in the following manner for their ability to inhibit Aurora-2 using a standard coupled enzyme assay (Fox et al (1998) *Protein Sci* 7, 2249).

To an assay stock buffer solution containing 0.1M HEPES 7.5, 10 mM MgCl<sub>2</sub>, 1 mM DTT, 25 mM NaCl, 2.5 mM phosphoenolpyruvate, 300 mM NADH, 30 mg/ml pyruvate kinase, 10 mg/ml lactate dehydrogenase, 40 mM ATP, and 800 μM peptide (LRRASLG, American Peptide, Sunnyvale, CA) was added a DMSO solution of a compound of the present invention to a final concentration of 30 μM. The resulting mixture was incubated at 30 °C for 10 min. The reaction was initiated by the addition of 10 μL of Aurora-2 stock solution to give a final concentration of 70 nM in the assay. The rates of reaction were obtained by monitoring absorbance at 340 nm over a 5 minute read time at 30 °C using a BioRad Ultramark plate reader (Hercules, CA). The K<sub>i</sub> values were determined from the rate data as a function of inhibitor concentration.

The following compounds were shown to have K<sub>i</sub> values less than 0.1 μM for Aurora-2: compounds II-33, II-34, II-36, II-37, II-40, II-41, II-55, III-7, III-9, III-37, III-38, III-39, III-40, III-41, III-42, III-44, III-45, III-46, III-47, III-48, III-49, III-50, III-51, III-52, III-53, III-54, III-55, III-56, III-57, III-59, III-60, III-61, III-63, III-30, III-65, III-66, III-67, III-70, III-31, III-76, III-77, III-78, III-80, III-32, III-33, III-34, III-106, III-108, III-109, III-110, III-111, III-112, III-114, III-35, III-115, III-116, III-117, III-118, III-119, III-120, III-121, IV-7, IV-30, IV-32, and IV-34.

The following compounds were shown to have  $K_i$  values between 0.1 and 1.0  $\mu\text{M}$  for Aurora-2: compounds II-1, II-105, II-35, II-38, II-39, II-42, II-64, II-70, II-53, II-99, II-77, II-79, II-86, II-20, II-93, II-94, III-28, III-58, III-64, III-71, III-73, III-74, III-75, III-102, III-105, III-107, III-113, III-124, III-1, III-130, IV-1, IV-3, IV-4, IV-6, IV-29, IV-33, and V-4.

The following compounds were shown to have  $K_i$  values between 1.0 and 20  $\mu\text{M}$  for Aurora-2: compounds II-103, II-104, II-57, II-59, II-61, II-63, II-67, II-69, II-75, II-76, II-10, II-19, II-78, II-54, II-80, II-82, II-21, II-90, II-91, II-96, II-107, III-68, III-79, III-82, III-101, III-103, III-127, III-141, III-129, III-132, IV-31, V-2, IX-47, IX-154, and IX-177.

### BIOLOGICAL TESTING EXAMPLE 3

#### CDK-2 INHIBITION ASSAY

Compounds were screened in the following manner for their ability to inhibit CDK-2 using a standard coupled enzyme assay (Fox et al (1998) *Protein Sci* 7, 2249).

To an assay stock buffer solution containing 0.1M HEPES 7.5, 10 mM  $\text{MgCl}_2$ , 1 mM DTT, 25 mM NaCl, 2.5 mM phosphoenolpyruvate, 300 mM NADH, 30 mg/ml pyruvate kinase, 10 mg/ml lactate dehydrogenase, 100 mM ATP, and 100  $\mu\text{M}$  peptide (MAHHHRSPRKRAKKK, American Peptide, Sunnyvale, CA) was added a DMSO solution of a compound of the present invention to a final concentration of 30  $\mu\text{M}$ . The resulting mixture was incubated at 30 °C for 10 min.

The reaction was initiated by the addition of 10  $\mu\text{L}$  of CDK-2/Cyclin A stock solution to give a final concentration of 25 nM in the assay. The rates of reaction were obtained by monitoring absorbance at 340 nm over a 5-minute read time at 30 °C using a BioRad

Ultramark plate reader (Hercules, CA). The  $K_i$  values were determined from the rate data as a function of inhibitor concentration.

5

#### BIOLOGICAL TESTING EXAMPLE 4

##### ERK INHIBITION ASSAY

Compounds were assayed for the inhibition of ERK2 by a spectrophotometric coupled-enzyme assay (Fox et al (1998) *Protein Sci* 7, 2249). In this assay, a fixed  
10 concentration of activated ERK2 (10 nM) was incubated with various concentrations of the compound in DMSO (2.5 %) for 10 min. at 30°C in 0.1 M HEPES buffer, pH 7.5, containing 10 mM  $MgCl_2$ , 2.5 mM phosphoenolpyruvate, 200  $\mu$ M NADH, 150  $\mu$ g/mL pyruvate kinase, 50  $\mu$ g/mL lactate  
15 dehydrogenase, and 200  $\mu$ M erktide peptide. The reaction was initiated by the addition of 65  $\mu$ M ATP. The rate of decrease of absorbance at 340 nM was monitored. The  $IC_{50}$  was evaluated from the rate data as a function of inhibitor concentration.

20

The following compounds were shown to have a  $K_i$  value of  $<1\mu$ M for ERK-2: III-109, III-111, III-115, III-117, III-118, III-120, and IV-4.

25

The following compounds were shown to have a  $K_i$  value of between  $1\mu$ M and  $12\mu$ M for ERK-2: III-63, III-40, and III-108.

#### BIOLOGICAL TESTING EXAMPLE 5

##### AKT INHIBITION ASSAY

Compounds were screened for their ability to  
30 inhibit AKT using a standard coupled enzyme assay (Fox et al., *Protein Sci.*, (1998) 7, 2249). Assays were carried out in a mixture of 100 mM HEPES 7.5, 10 mM  $MgCl_2$ , 25 mM NaCl, 1 mM DTT and 1.5% DMSO. Final substrate concentrations in the assay were 170  $\mu$ M ATP (Sigma

Chemicals) and 200  $\mu$ M peptide (RPRAATF, American Peptide, Sunnyvale, CA). Assays were carried out at 30 °C and 45 nM AKT. Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 300  $\mu$ M NADH, 30  $\mu$ g/ML pyruvate kinase and 10  $\mu$ g/ml lactate dehydrogenase.

An assay stock buffer solution was prepared containing all of the reagents listed above, with the exception of AKT, DTT, and the test compound of interest. 56  $\mu$ l of the stock solution was placed in a 384 well plate followed by addition of 1  $\mu$ l of 2 mM DMSO stock containing the test compound (final compound concentration 30  $\mu$ M). The plate was preincubated for about 10 minutes at 30 °C and the reaction initiated by addition of 10  $\mu$ l of enzyme (final concentration 45 nM) and 1 mM DTT. Rates of reaction were obtained using a BioRad Ultramark plate reader (Hercules, CA) over a 5 minute read time at 30 °C. Compounds showing greater than 50% inhibition versus standard wells containing the assay mixture and DMSO without test compound were titrated to determine IC<sub>50</sub> values.

#### BIOLOGICAL TESTING EXAMPLE 6

##### SRC INHIBITION ASSAY

The compounds were evaluated as inhibitors of human Src kinase using either a radioactivity-based assay or spectrophotometric assay.

##### Src Inhibition Assay A: Radioactivity-based Assay

The compounds were assayed as inhibitors of full length recombinant human Src kinase (from Upstate Biotechnology, cat. no. 14-117) expressed and purified from baculo viral cells. Src kinase activity was monitored by following the incorporation of <sup>33</sup>P from ATP into the tyrosine of a random poly Glu-Tyr polymer substrate of composition, Glu:Tyr = 4:1 (Sigma, cat. no.



P-0275). The following were the final concentrations of the assay components: 0.05 M HEPES, pH 7.6, 10 mM MgCl<sub>2</sub>, 2 mM DTT, 0.25 mg/ml BSA, 10  $\mu$ M ATP (1-2  $\mu$ Ci <sup>33</sup>P-ATP per reaction), 5 mg/ml poly Glu-Tyr, and 1-2 units of recombinant human Src kinase. In a typical assay, all the reaction components with the exception of ATP were pre-mixed and aliquoted into assay plate wells. Inhibitors dissolved in DMSO were added to the wells to give a final DMSO concentration of 2.5%. The assay plate was incubated at 30 °C for 10 min before initiating the reaction with <sup>33</sup>P-ATP. After 20 min of reaction, the reactions were quenched with 150  $\mu$ l of 10% trichloroacetic acid (TCA) containing 20 mM Na<sub>3</sub>PO<sub>4</sub>. The quenched samples were then transferred to a 96-well filter plate (Whatman, UNI-Filter GF/F Glass Fiber Filter, cat no. 7700-3310) installed on a filter plate vacuum manifold. Filter plates were washed four times with 10% TCA containing 20 mM Na<sub>3</sub>PO<sub>4</sub> and then 4 times with methanol. 200 $\mu$ l of scintillation fluid was then added to each well. The plates were sealed and the amount of radioactivity associated with the filters was quantified on a TopCount scintillation counter. The radioactivity incorporated was plotted as a function of the inhibitor concentration. The data was fitted to a competitive inhibition kinetics model to get the K<sub>i</sub> for the compound.

#### Src Inhibition Assay B: Spectrophotometric Assay

The ADP produced from ATP by the human recombinant Src kinase-catalyzed phosphorylation of poly Glu-Tyr substrate was quantified using a coupled enzyme assay (Fox et al (1998) *Protein Sci* 7, 2249). In this assay one molecule of NADH is oxidised to NAD for every molecule of ADP produced in the kinase reaction. The

disappearance of NADH can be conveniently followed at 340 nm.

The following were the final concentrations of the assay components: 0.025 M HEPES, pH 7.6, 10 mM MgCl<sub>2</sub>,  
5 2 mM DTT, 0.25 mg/ml poly Glu-Tyr, and 25 nM of recombinant human Src kinase. Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 200  $\mu$ M NADH, 30  $\mu$ g/ml pyruvate kinase and 10  $\mu$ g/ml lactate dehydrogenase.

10 In a typical assay, all the reaction components with the exception of ATP were pre-mixed and aliquoted into assay plate wells. Inhibitors dissolved in DMSO were added to the wells to give a final DMSO concentration of 2.5%. The assay plate was incubated at 30°C for 10 min  
15 before initiating the reaction with 100  $\mu$ M ATP. The absorbance change at 340 nm with time, the rate of the reaction, was monitored on a molecular devices plate reader. The data of rate as a function of the inhibitor concentration was fitted to competitive inhibition  
20 kinetics model to get the K<sub>i</sub> for the compound.

The following compounds were shown to have a K<sub>i</sub> value of <100nM on SRC: III-31, III-32, III-33, III-34, III-35, III-47, III-65, III-66, III-37, III-38, III-39, III-40, III-42, III-44, III-48, III-49, III-70, III-45,  
25 III-78, III-76, and IV- 32.

The following compounds were shown to have a K<sub>i</sub> value of between 100nM and 1 $\mu$ M for SRC: III-63, III-71, III-75, III-73, III-72, III-74, III-80, III-50, IV-30.


The following compounds were shown to have a K<sub>i</sub>  
30 value of between 1 $\mu$ M and 6 $\mu$ M for SRC: III-79, IV-1, and IV-31.

While we have hereinbefore presented a number of embodiments of this invention, it is apparent that our basic construction can be altered to provide other

embodiments which utilize the compounds and methods of  
this invention. Therefore, it will be appreciated that  
the scope of this invention is to be defined by the  
appended claims rather than by the specific embodiments  
5 which have been represented by way of example.

We claim:

1. A compound of formula VIII:



VIII

or a pharmaceutically acceptable derivative or prodrug thereof, wherein:

$Z^1$  is N or C- $R^9$ ,  $Z^2$  is N or CH, and  $Z^3$  is N or C- $R^x$ , provided that one of  $Z^1$  and  $Z^3$  is nitrogen;

G is Ring C or Ring D;

Ring C is selected from a phenyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, or 1,2,4-triazinyl ring, wherein said Ring C has one or two ortho substituents independently selected from  $-R^1$ , any non-ortho carbon position on Ring C is optionally and independently substituted by  $-R^5$ , and two adjacent substituents on Ring C are optionally taken together with their intervening atoms to form a fused, unsaturated or partially unsaturated, 5-6 membered ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, said fused ring being optionally substituted by halo, oxo, or  $-R^8$ ;

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein Ring D is

substituted at any substitutable ring carbon by halo, oxo, or  $-R^5$ , and at any substitutable ring nitrogen by  $-R^4$ , provided that when Ring D is a six-membered aryl or heteroaryl ring,  $-R^5$  is hydrogen at each ortho carbon position of Ring D;

$R^1$  is selected from -halo, -CN,  $-NO_2$ , T-V- $R^6$ , phenyl, 5-6 membered heteroaryl ring, 5-6 membered heterocyclyl ring, or  $C_{1-6}$  aliphatic group, said phenyl, heteroaryl, and heterocyclyl rings each optionally substituted by up to three groups independently selected from halo, oxo, or  $-R^8$ , said  $C_{1-6}$  aliphatic group optionally substituted with halo, cyano, nitro, or oxygen, or  $R^1$  and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

$R^x$  is T- $R^3$ ;

T is a valence bond or a  $C_{1-4}$  alkylidene chain;

$R^2$  and  $R^{2'}$  are independently selected from -R, -T-W- $R^6$ , or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable carbon on said fused ring formed by  $R^2$  and  $R^{2'}$  is substituted by halo, oxo, -CN,  $-NO_2$ ,  $-R^7$ , or -V- $R^6$ , and any substitutable nitrogen on said ring formed by  $R^2$  and  $R^{2'}$  is substituted by  $R^4$ ;

$R^3$  is selected from -R, -halo, -OR,  $-C(=O)R$ ,  $-CO_2R$ , -COCOR,  $-COCH_2COR$ ,  $-NO_2$ , -CN,  $-S(O)R$ ,  $-S(O)_2R$ , -SR,  $-N(R^4)_2$ ,  $-CON(R^7)_2$ ,  $-SO_2N(R^7)_2$ ,  $-OC(=O)R$ ,  $-N(R^7)COR$ ,  $-N(R^7)CO_2$ (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^7)CON(R^7)_2$ ,  $-N(R^7)SO_2N(R^7)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^7)_2$ ;

each R is independently selected from hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, C<sub>6-10</sub> aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each R<sup>4</sup> is independently selected from -R<sup>7</sup>, -COR<sup>7</sup>, -CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -CON(R<sup>7</sup>)<sub>2</sub>, or -SO<sub>2</sub>R<sup>7</sup>, or two R<sup>4</sup> on the same nitrogen are taken together to form a 5-8 membered heterocyclyl or heteroaryl ring;

each R<sup>5</sup> is independently selected from -R, halo, -OR, -C(=O)R, -CO<sub>2</sub>R, -COCOR, -NO<sub>2</sub>, -CN, -S(O)R, -SO<sub>2</sub>R, -SR, -N(R<sup>4</sup>)<sub>2</sub>, -CON(R<sup>4</sup>)<sub>2</sub>, -SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -OC(=O)R, -N(R<sup>4</sup>)COR, -N(R<sup>4</sup>)CO<sub>2</sub>(optionally substituted C<sub>1-6</sub> aliphatic), -N(R<sup>4</sup>)N(R<sup>4</sup>)<sub>2</sub>, -C=NN(R<sup>4</sup>)<sub>2</sub>, -C=N-OR, -N(R<sup>4</sup>)CON(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>N(R<sup>4</sup>)<sub>2</sub>, -N(R<sup>4</sup>)SO<sub>2</sub>R, or -OC(=O)N(R<sup>4</sup>)<sub>2</sub>, or R<sup>5</sup> and an adjacent substituent taken together with their intervening atoms form said ring fused to Ring C;

V is -O-, -S-, -SO-, -SO<sub>2</sub>-, -N(R<sup>6</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -N(R<sup>6</sup>)CO-, -N(R<sup>6</sup>)C(O)O-, -N(R<sup>6</sup>)CON(R<sup>6</sup>)-, -N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, -N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(O)N(R<sup>6</sup>)-, -OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-, -C(R<sup>6</sup>)=N-O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, or -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-;

W is -C(R<sup>6</sup>)<sub>2</sub>O-, -C(R<sup>6</sup>)<sub>2</sub>S-, -C(R<sup>6</sup>)<sub>2</sub>SO-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>-, -C(R<sup>6</sup>)<sub>2</sub>SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)-, -CO-, -CO<sub>2</sub>-, -C(R<sup>6</sup>)OC(O)-, -C(R<sup>6</sup>)OC(O)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CO-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)C(O)O-, -C(R<sup>6</sup>)=NN(R<sup>6</sup>)-, -C(R<sup>6</sup>)=N-O-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)SO<sub>2</sub>N(R<sup>6</sup>)-, -C(R<sup>6</sup>)<sub>2</sub>N(R<sup>6</sup>)CON(R<sup>6</sup>)-, or -CON(R<sup>6</sup>)-;

each R<sup>6</sup> is independently selected from hydrogen, an optionally substituted C<sub>1-4</sub> aliphatic group, or two R<sup>6</sup> groups on the same nitrogen atom are taken together with

the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;

each  $R^7$  is independently selected from hydrogen or an optionally substituted  $C_{1-6}$  aliphatic group, or two  $R^7$  on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring;

each  $R^8$  is independently selected from an optionally substituted  $C_{1-4}$  aliphatic group,  $-OR^6$ ,  $-SR^6$ ,  $-COR^6$ ,  $-SO_2R^6$ ,  $-N(R^6)_2$ ,  $-N(R^6)N(R^6)_2$ ,  $-CN$ ,  $-NO_2$ ,  $-CON(R^6)_2$ , or  $-CO_2R^6$ ; and  $R^9$  is selected from  $-R$ , halo,  $-OR$ ,  $-C(=O)R$ ,  $-CO_2R$ ,  $-COCOR$ ,  $-NO_2$ ,  $-CN$ ,  $-S(O)R$ ,  $-SO_2R$ ,  $-SR$ ,  $-N(R^4)_2$ ,  $-CON(R^4)_2$ ,  $-SO_2N(R^4)_2$ ,  $-OC(=O)R$ ,  $-N(R^4)COR$ ,  $-N(R^4)CO_2$  (optionally substituted  $C_{1-6}$  aliphatic),  $-N(R^4)N(R^4)_2$ ,  $-C=NN(R^4)_2$ ,  $-C=N-OR$ ,  $-N(R^4)CON(R^4)_2$ ,  $-N(R^4)SO_2N(R^4)_2$ ,  $-N(R^4)SO_2R$ , or  $-OC(=O)N(R^4)_2$ .

2. The compound according to claim 1, wherein said compound has one or more features selected from the group consisting of:

(a) Ring C is an optionally substituted ring selected from phenyl or pyridinyl, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a naphthyl, quinolinyll or isoquinolinyll ring, and  $R^1$  is -halo, an optionally substituted  $C_{1-6}$  aliphatic group, phenyl,  $-COR^6$ ,  $-OR^6$ ,  $-CN$ ,  $-SO_2R^6$ ,  $-SO_2NH_2$ ,  $-N(R^6)_2$ ,  $-CO_2R^6$ ,  $-CONH_2$ ,  $-NHCOR^6$ ,  $-OC(O)NH_2$ , or  $-NHCO_2R^6$ ; or Ring D is an optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyll, 1,2,3,4-tetrahydroquinolinyll, 2,3-dihydro-1H-isoindolyl, 2,3-

dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^x$  is  $T-R^3$  wherein T is a valence bond or a methylene; and

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring.

3. The compound according to claim 2, wherein:

(a) Ring C is an optionally substituted ring selected from phenyl or pyridinyl, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is selected from a naphthyl, quinolinyl or isoquinolinyl ring, and  $R^1$  is -halo, an optionally substituted  $C_{1-6}$  aliphatic group, phenyl,  $-COR^6$ ,  $-OR^6$ ,  $-CN$ ,  $-SO_2R^6$ ,  $-SO_2NH_2$ ,  $-N(R^6)_2$ ,  $-CO_2R^6$ ,  $-CONH_2$ ,  $-NHCOR^6$ ,  $-OC(O)NH_2$ , or  $-NHSO_2R^6$ ; or Ring D is an optionally substituted ring selected from a phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, thienyl, azepanyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl ring;

(b)  $R^x$  is  $T-R^3$  wherein T is a valence bond or a methylene; and

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, heteroaryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a



substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring.

4. The compound according to claim 2, wherein said compound has one or more features selected from the group consisting of:

(a) Ring C is an optionally substituted ring selected from phenyl or pyridinyl, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1H-isoindolyl, 2,3-dihydro-1H-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^x$  is  $T-R^3$  wherein T is a valence bond or a methylene and  $R^3$  is CN, -R or -OR;

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring; and

(d) each  $R^5$  is independently selected from -halo, -CN, -NO<sub>2</sub>, -N( $R^4$ )<sub>2</sub>, an optionally substituted  $C_{1-6}$  aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH( $R^4$ ), -N( $R^4$ )COR, -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, or -N( $R^4$ )SO<sub>2</sub>R.

5. The compound according to claim 4, wherein:

(a) Ring C is an optionally substituted ring selected from phenyl or pyridinyl, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-6}$  haloaliphatic group, a  $C_{1-6}$  aliphatic group, phenyl, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 2,3-dihydro-1*H*-isoindolyl, 2,3-dihydro-1*H*-indolyl, isoquinolinyl, quinolinyl, or naphthyl;

(b)  $R^x$  is  $T-R^3$  wherein T is a valence bond or a methylene and  $R^3$  is CN, -R or -OR;

(c)  $R^{2'}$  is hydrogen and  $R^2$  is hydrogen or a substituted or unsubstituted group selected from aryl, or a  $C_{1-6}$  aliphatic group, or  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a substituted or unsubstituted benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring; and

(d) each  $R^5$  is independently selected from -halo, -CN, -NO<sub>2</sub>, -N( $R^4$ )<sub>2</sub>, an optionally substituted  $C_{1-6}$  aliphatic group, -OR, -C(O)R, -CO<sub>2</sub>R, -CONH( $R^4$ ), -N( $R^4$ )COR, -SO<sub>2</sub>N( $R^4$ )<sub>2</sub>, or -N( $R^4$ )SO<sub>2</sub>R.

6. The compound according to claim 4, wherein said compound has one or more features selected from the group consisting of:

(a)  $R^x$  is  $T-R^3$  wherein T is a valence bond or a methylene and  $R^3$  is -R or -OR wherein R is an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring;

(b) Ring C is an optionally substituted ring selected from phenyl or pyridinyl, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or naphthyl;

(c)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring optionally substituted with -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4} \text{ alkyl})$ ,  $-CO_2(C_{1-4} \text{ alkyl})$ , -CN,  $-SO_2(C_{1-4} \text{ alkyl})$ ,  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4} \text{ alkyl})$ ,  $-NHC(O)(C_{1-4} \text{ alkyl})$ ,  $-C(O)NH_2$ , or  $-CO(C_{1-4} \text{ alkyl})$ , wherein the  $(C_{1-4} \text{ alkyl})$  is a straight, branched, or cyclic alkyl group;

(d) each  $R^5$  is independently selected from -Cl, -F, -CN,  $-CF_3$ ,  $-NH_2$ ,  $-NH(C_{1-4} \text{ aliphatic})$ ,  $-N(C_{1-4} \text{ aliphatic})_2$ ,  $-O(C_{1-4} \text{ aliphatic})$ ,  $C_{1-4} \text{ aliphatic}$ , and  $-CO_2(C_{1-4} \text{ aliphatic})$ ; and

(e)  $R^9$  is R, OR, or  $N(R^4)_2$ .

7. The compound according to claim 6, wherein:

(a)  $R^x$  is  $T-R^3$  wherein T is a valence bond or a methylene and  $R^3$  is -R or -OR wherein R is an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, or a 5-6 membered heteroaryl or heterocyclyl ring;

(b) Ring C is an optionally substituted ring selected from phenyl or pyridinyl, wherein when Ring C and two adjacent substituents thereon form a bicyclic ring

system, the bicyclic ring system is a naphthyl ring, and  $R^1$  is -halo, a  $C_{1-4}$  aliphatic group optionally substituted with halogen, or -CN; or Ring D is an optionally substituted ring selected from phenyl, pyridinyl, piperidinyl, piperazinyl, pyrrolidinyl, morpholinyl, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, isoquinolinyl, quinolinyl, or naphthyl;

(c)  $R^2$  and  $R^{2'}$  are taken together with their intervening atoms to form a benzo, pyrido, pyrimido or partially unsaturated 6-membered carbocyclo ring optionally substituted with -halo,  $-N(R^4)_2$ ,  $-C_{1-4}$  alkyl,  $-C_{1-4}$  haloalkyl,  $-NO_2$ ,  $-O(C_{1-4} \text{ alkyl})$ ,  $-CO_2(C_{1-4} \text{ alkyl})$ , -CN,  $-SO_2(C_{1-4} \text{ alkyl})$ ,  $-SO_2NH_2$ ,  $-OC(O)NH_2$ ,  $-NH_2SO_2(C_{1-4} \text{ alkyl})$ ,  $-NHC(O)(C_{1-4} \text{ alkyl})$ ,  $-C(O)NH_2$ , or  $-CO(C_{1-4} \text{ alkyl})$ , wherein the  $(C_{1-4} \text{ alkyl})$  is a straight, branched, or cyclic alkyl group;

(d) each  $R^5$  is independently selected from -Cl, -F, -CN,  $-CF_3$ ,  $-NH_2$ ,  $-NH(C_{1-4} \text{ aliphatic})$ ,  $-N(C_{1-4} \text{ aliphatic})_2$ ,  $-O(C_{1-4} \text{ aliphatic})$ ,  $C_{1-4} \text{ aliphatic}$ , and  $-CO_2(C_{1-4} \text{ aliphatic})$ ; and

(e)  $R^9$  is R, OR, or  $N(R^4)_2$ .

8. The compound according to claim 1, wherein said compound is selected from Table 7.

9. A composition comprising a compound according to any of claims 1-8 and a pharmaceutically acceptable carrier.

10. The composition according to claim 9 further comprising a second therapeutic agent.

11. A method of inhibiting GSK-3 or Aurora activity in a patient comprising the step of administering to said patient a therapeutically effective amount of the composition according to claim 9.

12. The method according to claim 11, wherein said method inhibits GSK-3 activity in a patient.

13. A method of inhibiting GSK-3 or Aurora activity in a biological sample comprising contacting said biological sample with the compound according to claim 1.

14. A method of treating a disease that is alleviated by treatment with an GSK-3 inhibitor, said method comprising the step of administering to a patient in need of such a treatment a therapeutically effective amount of the composition according to claim 9.

15. The method according to claim 14 further comprising the step of administering to said patient a second therapeutic agent.

16. The method according to claim 14, wherein said disease is diabetes.

17. The method according to claim 14, wherein said disease is Alzheimer's disease.

18. The method according to claim 14, wherein said disease is schizophrenia.

19. A method of enhancing glycogen synthesis in a patient in need thereof, which method comprises the step of administering to said patient a therapeutically effective amount of the composition according to claim 9.

20. A method of lowering blood levels of glucose in a patient in need thereof, which method comprises the step of administering to said patient a therapeutically effective amount of the composition according to claim 9.

21. A method of inhibiting the production of hyperphosphorylated Tau protein in a patient in need thereof, which method comprises the step of administering to said patient a therapeutically effective amount of the composition according to claim 9.

22. A method of inhibiting the phosphorylation of  $\beta$ -catenin in a patient in need thereof, which method comprises the step of administering to said patient a therapeutically effective amount of the composition according to claim 9.

23. A method of treating a disease that is alleviated by treatment with an aurora inhibitor, which method comprises the step of administering to said patient in need of such a treatment a therapeutically effective amount of the composition according to claim 9.

24. The method according to claim 23, further comprising the step of administering to said patient a second therapeutic agent.

25. The method according to claim 23 wherein said disease is cancer.

# INTERNATIONAL SEARCH REPORT

In tional Application No  
F.../US 01/28793

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D403/12 C07D401/14 C07D409/14 A61K31/497 A61K31/53  
A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 21955 A (PASQUET GEORGES ;HENNEQUIN LAURENT FRANCOIS AND (FR); ZENECA PHARM) 20 April 2000 (2000-04-20) examples 16-20 ---	1, 16, 23, 25
A	WO 00 39101 A (BREault GLORIA ANNE ;PEASE JANET ELIZABETH (GB); ASTRAZENECA UK LT) 6 July 2000 (2000-07-06) example 50 ---	1, 23, 25
A	WO 95 15758 A (HSU CHIN YI JENNY ;ZILBERSTEIN ASHER (US); JOHNSON SUSAN E (US); M) 15 June 1995 (1995-06-15) page 15, line 22 -----	1, 23, 25

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

28 November 2001

Date of mailing of the international search report

10/12/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B



## INTERNATIONAL SEARCH REPORT

Int lional Application No

PCT/US 01/28793

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0021955	A	20-04-2000	AU 6112899 A	01-05-2000
			BR 9914326 A	26-06-2001
			CN 1322202 T	14-11-2001
			EP 1119567 A1	01-08-2001
			WO 0021955 A1	20-04-2000
			NO 20011739 A	07-06-2001
WO 0039101	A	06-07-2000	AU 1874300 A	31-07-2000
			BR 9916590 A	23-10-2001
			EP 1140860 A1	10-10-2001
			WO 0039101 A1	06-07-2000
			NO 20013038 A	22-08-2001
WO 9515758	A	15-06-1995	US 5480883 A	02-01-1996
			US 5710158 A	20-01-1998
			AU 1305095 A	27-06-1995
			EP 0871448 A1	21-10-1998
			SG 54172 A1	16-11-1998
			WO 9515758 A1	15-06-1995
			US 5795889 A	18-08-1998
			US 5646153 A	08-07-1997
			US 5721237 A	24-02-1998
			US 5714493 A	03-02-1998
			US 6057320 A	02-05-2000
			US RE36256 E	20-07-1999

This Page Blank (uspto)